TEXTE

94/2020

Development of an OECD Guidance Document for the Application of OECD Test Guideline 236 (Acute Fish Embryo Toxicity Test) Final Report



TEXTE 94/2020

Ressortforschungsplan of the Federal Ministry for the Enviroment, Nature Conservation and Nuclear Safety

Project No. (FKZ) 3715674190 Report No. (FB) FB000024/ENG

Development of an OECD Guidance Document for the Application of OECD Test Guideline 236 (Acute Fish Embryo Toxicity Test)

Final report

by

Prof. Dr. Thomas Braunbeck, Dr. Svenja Böhler, Dr. Susanne Knörr, Ann-Kathrin Lörracher, Dr. Katharina Pelka and Dr. Britta Kais

Aquatic Ecology and Toxicology, Center for Organismal Studies, University of Heidelberg, Im Neuenheimer Feld 504, D-69120 Heidelberg

On behalf of the German Environment Agency

Imprint

Publisher

Umweltbundesamt Wörlitzer Platz 1 06844 Dessau-Roßlau Tel: +49 340-2103-0 Fax: +49 340-2103-2285 <u>buergerservice@uba.de</u> Internet: <u>www.umweltbundesamt.de</u>

✔/<u>umweltbundesamt.de</u>
 ✔/<u>umweltbundesamt</u>

Report performed by:

Aquatic Ecology and Toxicology, Center for Organismal Studies, University of Heidelberg Im Neuenheimer Feld 504 D-69120 Heidelberg Federal Republic of Germany

Report completed in:

August 2019

Edited by:

Section IV 1.1 – International Chemicals Management Walter-Rohde, Susanne

Publication as pdf: http://www.umweltbundesamt.de/publikationen

ISSN 1862-4804

Dessau-Roßlau, June 2020

The responsibility for the content of this publication lies with the author(s).

In July 2013, the Fish Embryo Acute Toxicity Test (FET) was adopted by OECD as Test Guideline 236. The FET thus represents the first fully validated alternative method within the OECD Test Guidelines Programme in the ecotoxicology sector. However, the practical applicability of the FET has ever since been discussed with respect to a series of aspects, which are addressed in this report: With respect to the potential barrier function of the zebrafish chorion, the molecular size for a free passage of non-charged chemical substances across the chorion of the zebrafish could be set at 3,000 - 4,000 Da. The limit for the passage of charged molecules is lower than that of non-charged ones. DMSO concentrations of ≥ 0.1 % decrease the barrier function of the chorion. Zebrafish embryos, juveniles and adults have been studied to a different extent with respect to their biotransformation capacities. Our knowledge on biotransformation capacities in zebrafish is fragmentary. On this basis, transfer of biotransformation data from juvenile to adult fish is not possible. In zebrafish embryos, biotransformation could be documented whenever studied in more detail; only for rare exceptions, lack of biotransformation was demonstrated. Gene activation cannot be translated into biochemical functionality; mammalian data cannot be extrapolated to fish. The analysis of historical short-term (acute) Fish Toxicity Test (AFT) data produced according to OECD TG 203 within a dataset used for a corresponding FET study by ECHA started with 2936 studies on a total of 1842 substances. Applications of filters identical to those used for the FET analysis resulted in exclusion of 62.3 % of the AFT studies and 63.0 % of the substances. Data of the present report will be integrated into OECD project no. 2.54: "Guidance Document on an Integrated Approach on Testing and Assessment for Fish Acute Toxicity Testing" – Integration of the Fish Embryo Test into the Threshold Approach".

Kurzbeschreibung: Chorionstruktur und Biotransformationskapazität des Zebrabärblings als Randparameter für die OECD-Richtlinie 236 (Akuter Fischembryotest)

Im Juli 2013 wurde der akute Fischembryotest (FET) als OECD-Prüfrichtlinie 236 anerkannt. Der FET stellt damit die erste voll validierte Alternativmethode im Rahmen des OECD-Prüfrichtlinienprogramms im Bereich der Ökotoxikologie dar. Die praktische Anwendbarkeit wurde jedoch seitdem hinsichtlich mehrerer Aspekte diskutiert, die im vorliegenden Bericht adressiert werden: Im Hinblick auf die potentielle Barrierefunktion der Eihülle des Zebrabärblings konnte eine Molekulargröße von 3000 – 4000 Da für die freie Passage ungeladener Moleküle identifiziert werden. Das Limit für die Passage geladener Moleküle ist geringer. DMSO-Konzentrationen $\geq 0,1$ % reduzieren die Barrierefunktion weiter. Embryonen, juvenile und erwachsene Zebrabärblinge sind im Hinblick auf ihre Biotransformation unterschiedlich gut untersucht; insgesamt ist unser Wissen fragmentiert. Auf dieser Basis ist die Übertragung von Daten von juvenilen Fischen auf erwachsene und umgekehrt nicht möglich. Wann immer die Biotransformation in Embryonen genauer betrachtet wurde, konnte sie zumindest qualitativ nachgewiesen werden. Nur für seltene Ausnahmen, konnte eine nicht vorhandene Biotransformation im Embryo demonstriert werden. Genaktivierung kann nicht in biologische Funktionalität übersetzt werden; Säugerdaten können nicht auf Fisch extrapoliert werden. Die Analyse eines historischen Datensatzes zur akuten Fischtoxizität, der bereits als Grundlage für die Bewertung der Eignung des FET gedient hatte, ging von 2936 Studien mit Daten zu 1842 Substanzen aus. Die Anwendung der gleichen Filter, die von der ECHA für den FET angesetzt wurden, führte zum Ausschluss von 62,3 % der Studien bzw. 63,0 % der Substanzen. Die Daten des vorliegenden Berichts werden in das OECD-Projekt Nr. 254 ("Guidance Document on an Integrated Approach on Testing and Assessment for Fish Acute Toxicity Testing" – Integration of the Fish Embryo Test into the Threshold Approach") integriert.

Table of content

List of figures							
List of tables 10							
Li	List of abbreviations						
Sι	Summary 12						
Zι	Zusammenfassung1						
1 The zebrafish (<i>Danio rerio</i>) chorion as a potential barrier for the uptake of xenobiotics into							
	th	e zebrafish embryo	20				
	1.1	Introduction	20				
	1.2	Development of the zebrafish chorion2	20				
	1.3	Structure of the zebrafish chorion	23				
	1.4	Dechorionation as a tool to improve the fish embryo toxicity test (FET) with the zebrafish (Danio rerio)	27				
	1.5	Fluorescent dyes as markers for the permeability of the chorion in zebrafish (<i>Danio rerio</i>) embryos	29				
	1.6	Polyethylene glycols of different molecular size as markers of chorion permeability in the zebrafish (<i>Danio rerio</i>) embryo	39				
	1.7	Conclusions: Barrier function of the zebrafish chorion	13				
2	Bie a l	otransformation and bioactivation capacities in early life stages of zebrafish (<i>Danio rerio</i>) – iterature review	15				
	2.1	Introduction	15				
	2.2	Metabolism of xenobiotics	15				
	2.3	The role of metabolism in activation and detoxification of xenobiotics	16				
	2.4	Cytochrome P450-dependent monooxygenases (CYPs)	16				
	2.5	Number of CYP isoforms in zebrafish	16				
	2.6	Expression patterns of CYPs during zebrafish development	17				
	2.7	Spatial expression patterns of CYPs in zebrafish6	52				
	2.8	CYP-dependent activities6	58				
	2.9	Inducibility of cytochrome P450s in zebrafish	73				
	2.10	Bioactivation capacity of early-life stages: functional confirmation of CYP activity	73				
	2.11	Metabolism capacity – chemical analysis	74				
	2.12	Conclusions and recommendations for future research into the biotransformation capacities of zebrafish (<i>Danio rerio</i>)	30				
3	Ar Of	nalysis of the relevance and adequateness of the Fish Acute Toxicity Test (AFT) according to ECD TG 203 to fulfil the information requirements and addressing concerns under REACH &	32				

	3.1	Summary	
	3.2	Introduction	83
	3.3	Treatment of datasets	83
	3.4	Results of the data filtering procdure	85
	3.4.1	Filtering steps I and II: Unsuitable LC_{50} values and duplicate studies	. 85
	3.4.2	Filtering steps III and IV: LC_{50} and water solubility	. 86
	3.4.3	Elimination step V: Analysis of log Pow data	88
	3.4.4	Further filtering: Inorganic substances	91
	3.4.5	Further filtering: AFT studies with zebrafish (Danio rerio) only	91
	3.4.6	Further filtering: Interspecies and intraspecies comparisons	91
	3.5	Conclusions drawn from the analysis of AFT data from the ECHA database and	
		recommendations for future AFT studies	93
	3.5.1	Comparison of data handling in the present AFT study and the FET by (Scholz et al.,	
		2016)	93
	3.5.2	Conclusions and recommendations for future AFT testing	. 95
4	Cont	tributions to OECD project no. 2.54: "Guidance Document on an Integrated Approach on	
	Test	ing and Assessment for Fish Acute Toxicity Testing" – Integration of the Fish Embryo Test	
	into	the Threshold Approach (OECD Guidance Document 126)	96
5	Refe	rences	97

List of figures

Figure 1:	Oocytes of zebrafish (Danio rerio)21
Figure 2:	Zebrafish (Danio rerio) oocytes in maturation
Figure 3:	The zebrafish chorion23
Figure 4:	Surface of the chorion of zebrafish (Danio rerio) eggs24
Figure 5:	Layers of the zebrafish (Danio rerio) chorion
Figure 6:	Surface of the zebrafish (Danio rerio) chorion
Figure 7:	Inner surface of the zebrafish (Danio rerio) chorion25
Figure 8:	Granular layer of the zebrafish (Danio rerio) chorion26
Figure 9:	Pore canals in the chorion of zebrafish (Danio rerio)
Figure 10:	Dechorionation of zebrafish (Danio rerio) embryos27
Figure 11:	Correlation of zebrafish (Danio rerio) age and survival after
	dechorionation28
Figure 12:	Mortalities of non-dechorionated zebrafish (Danio rerio)
	embryos after exposure to Luviquat28
Figure 13:	Mortalities of dechorionated and non-dechorionated zebrafish
	(Danio rerio) embryos after exposure to Luviquat
Figure 14:	Rhodamine and sulforhodamine transport across the zebrafish
	(<i>Danio rerio</i>) chorion31
Figure 15:	Time-dependent passage of the fluorescent marker
	dichlorofluorescein across the zebrafish (Danio rerio)
	chorion31
Figure 16:	Impact of DMSO on substance transfer across the chorion of
	zebrafish (Danio rerio)32
Figure 17:	Impact of DMSO on substance uptake into zebrafish (Danio
	<i>rerio</i>) embryos32
Figure 18:	Epifluorescence images of zebrafish (Danio rerio) embryos33
Figure 19:	Cross-sections and surface images of zebrafish (Danio rerio)
	embryos34
Figure 20:	Transmission, cross-section and 3D images of zebrafish (Danio
	<i>rerio</i>) embryos35
Figure 21:	Composite 3D surface confocal laser images of zebrafish (Danio
	<i>rerio</i>) embryos36
Figure 22:	Cross-section and composite 3D surface images of zebrafish
	(Danio rerio) embryos after exposure to 2,7-
	dichlorofluorescein I37
Figure 23:	Cross-section and composite 3D surface images of zebrafish
	(Danio rerio) embryos after exposure to 2,7-
	dichlorofluorescein II38
Figure 24:	Shrinkage of the chorion of zebrafish (Danio rerio) embryos
	after transfer into polyethylene glycols of different molecular
	mass

Figure 25:	Sublethal effects of PEG 6000 in zebrafish (<i>Danio rerio</i>) embryos
Figure 26:	Chorion deformation of zebrafish (<i>Danio rerio</i>) eggs after
	exposure to PEGs41
Figure 27:	Chronological sequence of deformation of the zebrafish (<i>Danio rerio</i>) chorion following exposure to differently sized
	nolvethylene glycols (PEGs)
Eiguro 29	Changes in the area of the chorion of zohrafich (Dania raria)
Figure 20.	embryos exposed to polyethylene glycols (PEGs) of different
	molecular weight43
Figure 29:	CYP3A65 gene expression trends in early-life stages of
	zebrafish (Danio rerio)47
Figure 30:	Temporal expression of CYP450 in embryonic and juvenile
	zebrafish (Danio rerio)48
Figure 31:	In vivo localization of EROD activity in 8 h old zebrafish (Danio
	<i>rerio</i>) embryos68
Figure 32:	EROD induction patterns in zebrafish (Danio rerio) embryos
	after 3 h pulse and continuous long-term exposure
Figure 33:	In vivo localization of EROD activities in zebrafish (Danio rerio)
	embryos after exposure to chlorpyrifos in various scenarios I 70
Figure 34:	In vivo localization of EROD activities in zebrafish (Danio rerio)
	embryos after exposure to chlorpyrifos in various scenarios
	II71
Figure 35:	In vivo localization of EROD activities in zebrafish (Danio rerio)
	embryos after exposure to chlorpyrifos in various scenarios
	III72
Figure 36:	Overview on the filtering steps in the ECHA dataset
Figure 37:	Elimination of unclear entries and duplicate studies in the
-	ECHA dataset
Figure 38:	Entries in the ECHA database with LC_{50} > water solubility, no
	data on water solubility and water solubility given as range
	with the LC_{50} lying within this range
Figure 39:	Ratio of the studies in the ECHA database with use of analytics
-	and vehicle in studies with LC_{50} tested above water solubility 87
Figure 40:	Solvent use and analytical verification in substances from the
_	ECHA database tested with LC ₅₀ over water solubility87
Figure 41:	Analytical verification of the test concentrations of substances
	tested within the ECHA database, when log Pow > 4
Figure 42:	Summary of study elimination per filtering step relative to
-	original data (%)90
Figure 43:	Range of deviation factors in intra- and interspecies
-	comparisons within the ECHA database

List of tables

Table 1:	Spatial and temporal patterns of CYP enzyme expression in
	zebrafish (Danio rerio)58
Table 2:	CYP isoforms in adult, juvenile, and embryonic life-stages of
	zebrafish (Danio rerio)61
Table 3:	Spatial expression of CYP3C1 in developmental stages of
	zebrafish (Danio rerio)63
Table 4:	Spatial expression patterns of CYP 450 in zebrafish (Danio
	rerio)64
Table 5:	Spatial expression patterns of cytochrome P450 in adult
	zebrafish (Danio rerio)66
Table 6:	LC_{50} , EC_{50} and teratogenicity index (TI) of selected
	proteratogens in 3 d old zebrafish (Danio rerio) embryos74
Table 7:	Phase I and phase II metabolites identified in zebrafish (Danio
	rerio)75
Table 8:	Testosterone metabolite profiles zebrafish (Danio rerio) at
	various developmental stages80
Table 9:	Overview of the search criteria in the ECHA database for
	historical AFT data83
Table 10:	Absolute number of studies/substances removed or
	considered for final analysis89
Table 11:	FET and AFT sets and numbers of substances/studies
	considered in final dataset94

List of abbreviations

AFT	Acute Fish Toxicity (OECD TG 203)			
CLSM	Confocal Laser Scanning Microscopy			
COS	Center for Organismal Studies (University of Heidelberg)			
DMSO	Dimethyl sulfoxide			
СҮР	Cytochrome P450			
3D	Three-dimensional			
Da	Dalton			
ECHA	European Chemicals Agency, Helsinki			
EC ₅₀	Test concentration with 50 % effect			
ECOD	7-Ethoxycoumarin-O-deethylase			
EDTA	Ethylene diamine tetraacetic acid			
EROD	7-Ethoxyresorufin-O-deethylase			
FET	Fish Embryo Acute Toxicity Test (OECD TG 236)			
GD	(OECD) Guidance Document			
hpf	hours post-fertilization			
h	Hours			
LC ₅₀	Test concentration with 50 % mortality			
LOEC	Lowest Observed Effect Concentration			
NOEC	No Observed Effect Concentration			
mRNA	Messenger ribonucleic acid			
OECD	Organization for Economic Co-operation and Development			
P450	Cytochrome P450			
PEG	Polyethylene glycol			
POW	Octanol/water partitioning coefficient			
REACH	Registration, Evaluation, Authorization and Restriction of Chemicals			
TG	(OECD) Test Guideline			
ті	Teratogenicity index			
UBA	German Environment Agency, Dessau			
VMG-eco	(OECD) Validation Management Group for Ecotoxicity Testing			
WNT	Working Group of National Coordinators for the OECD Test Guidelines Programme			
For abbreviations in figures, see figures legends.				

Summary

In July 2013, the Fish Embryo Acute Toxicity Test (FET; OECD 2013) validated by Germany together with the OECD Ad hoc Expert Group on the Fish Embryo Toxicity Test and the OECD Validation Management Group for Ecotoxicity Testing (VMG-eco) was adopted by the Working Group of National Coordinators for the OECD Test Guidelines Programme (WNT) as OECD Test Guideline (TG) 236 under the title "Fish Embryo Acute Toxicity (FET) Test". Given that nonfeeding stages of zebrafish (Danio rerio) development do not fall under the auspices of the current EU Directive 2010/63/EU on the protection of animals used for scientific purposes (EU, 2010) until the age of 5 days (Strähle et al., 2011), the procedure of the Fish Embryo Acute Toxicity (FET) Test represents the first fully validated alternative method within the OECD Test Guideline Programme in the ecotoxicology section. However, since its adoption by OECD in 2013, the practical applicability of the FET has been discussed with respect to a series of aspects including (1) the suspected, but poorly understood barrier function of the chorion and (2) the speculated, but hardly documented biotransformation potential of zebrafish embryos. With respect to the applicability of the FET within the REACH regulation, (3) the adequateness of the FET has been much debated. Only recently, Sobanska et al. (2018) published an analysis of the FET protocol and database, which, however, gave rise to criticism and will, thus, be critically questioned in the present publication. Finally, the suitability of the FET (4) to contribute to other OECD projects such as OECD project no. 2.54 ("Guidance Document on an Integrated Approach on Testing and Assessment for Fish Acute Toxicity Testing" - Integration of the Fish Embryo Test into the Threshold Approach; OECD Guidance Document 126) is being discussed. All of these issues are addressed in the present UBA report.

The potential barrier function of the zebrafish chorion

For long, a major objection to the acceptance of the Fish Embryo Test (FET) according to OECD TG 236 has been the putative protecting role of the chorion (eggshell) of the embryo. Whenever fish embryo toxicity tests produced results different from data published for the conventional Acute Fish Toxicity test (AFT) according to OECD TG 203, impaired permeability of the zebrafish chorion has been speculated. Starting as early as in 2008, the Federal Environmental Protection Agency (UBA) has funded investigations into the permeability of the zebrafish chorion at the University of Heidelberg (cf. UBA report FKZ 37080650400). Since then, more-in-depth experiments have been carried out. The present report provides an integration of all investigations conducted so far at the Aquatic Ecology and Toxicology Group in the Center for Organismal Studies at the University of Heidelberg and elsewhere.

With respect to the potential barrier function of the zebrafish chorion, further evidence was provided by means of fluorescent markers and differently sized polyethylene glycols to conclude that the molecular size for a free passage of non-charged chemical substances across the chorion of the zebrafish could be set at approx. 3,000 - 4,000 Da. The rate of passage of charged molecules across the chorion of zebrafish is substantially lower than that of non-charged molecules. As a conclusion, the passage of most uncharged industrial chemicals across the zebrafish molecule can be assumed to not be impaired by a systematic size limitation.

Dimethyl sulfoxide (DMSO) concentrations of ≥ 0.1 % decrease the barrier function of the chorion. Since dimethyl sulfoxide (DMSO) does not affect zebrafish embryo survival up to concentrations well above 2 %, DMSO may be used as a detergent to facilitate the passage of substances that might be critical in terms of their molecular size to pass the zebrafish chorion. Since DMSO does not affect zebrafish embryo survival up to concentrations well above 2 %, DMSO may be used as a solvent to facilitate the passage of substances that might be critical for handling in aqueous solutions due to their high lipophilicity. So far, there is no convincing

evidence that – after appropriate solution by the aid of solvents – highly lipophilic substances are not taken up by zebrafish embryos. See also advice given in OECD Guidance Document (GD) no. 23 on aqueous-phase aquatic toxicity testing of difficult test chemicals.

For highly lipophilic substances, neither the Acute Fish Toxicity test (AFT, OCED TG 203) nor the Fish Embryo Toxicity test (FET, OECD TG 236) appear to be suitable protocols, since highly lipophilic substances might accumulate over extended periods of time and might not have reached a steady-state after an exposure time restricted to 96 h. Rather, for highly lipophilic substances, longer-term tests such as a Fish Early Life-Stage test according to OECD TG 210 should be conducted.

In case of substances, the passage of which across the zebrafish chorion might be critical, dechorionation of zebrafish embryos may be applied: With appropriate modifications of the FET protocol, embryos can reproducibly be dechorionated at 24 h post-fertilization (hpf) with reliably high survival rates of \geq 90 %. Dechorionation of younger stages (< 24 hpf) is generally possible, however with lower survival rates.

Biotransformation and bioactivation capacities in early life stages of zebrafish (*Danio rerio*)

Major objections to the use of the fish embryo test (FET) with the zebrafish (Danio rerio) have been based on the assumption of limited biotransformation capacities in early life stages of fish. Repeatedly, doubts have been expressed concerning the comparability of metabolic capacity between early juvenile and adult life stages of zebrafish. In fact, a limited biotransformation or bioactivation capacity could lead to underestimations of toxic or teratogenic potency of a xenobiotic, which would be bioactivated in other life stages. On the other hand, given the *ex utero* development, early-life stages of (zebra)fish depend on their intrinsic metabolic capacity for elimination or transformation of xenobiotics; therefore, it seems reasonable to assume at least qualitatively appropriate biotransformation capacities.

As a consequence, in order to strengthen the acceptance of the Fish Embryo Toxicity (FET) test, it seemed essential to attempt to better understand the metabolic capacities of zebrafish embryos across different life-stages. However, while literature concerning spatial and temporal expression patterns of xenobiotic metabolizing enzymes has been growing fast at the genome and RNA levels, only few investigations addressed protein function and catalytic activities, although these are ecologically much more relevant parameters than DNA- and RNA-based characteristics. The database concerning zebrafish metabolism, especially in early-life stages, is much less complete than that available for other model organisms (e.g. rainbow trout) and is scattered over more than 100 studies.

So far, zebrafish embryos, juveniles and adults have been studied to a very different extent with respect to their biotransformation capacities. Overall, our knowledge about xenobiotic transformation capacities in zebrafish can – at best – be called fragmentary. Since especially juveniles have largely been neglected, transfer of biotransformation data from juvenile fish to adult fish does not appear justified. In zebrafish embryos, however, biotransformation could be documented whenever studied in more detail. So far, only for rare exceptions such as allyl alcohol and albendazol, a lack of biotransformation could be demonstrated. Studies on biotransformation gene expression (induction) is not as conclusive as studies in enzyme biosynthesis (transcriptomics, proteomics) or – even better – biochemical activity of enzymes. Since our knowledge on the spectrum of substances accepted as substrate(s) by various cytochrome P450 isoforms is quite limited, there is no way to extrapolate gene activation to biochemical functionality.

There is no adequate database to extrapolate observations from mammalian studies to fish and fish embryos. For many P450 isoforms, we do not even have conclusive evidence of congruence in cytochrome P450 terminology between mammalian studies and studies in lower vertebrates. Therefore, transfer of a link between nomenclature and metabolic function from mammals to fish ("same nomenclature means same function") or vice versa is not necessarily justified. With respect to biotransformation phase II reactions, our knowledge is even more fragmentary than it is for phase I metabolization.

As a gold standard for studies into biotransformation capacities in (zebra)fish embryos, juveniles and adults, gene expression studies need to be linked to proteome analysis data and chemical analytical analyses into the metabolites formed by cytochrome P450 enzymatic activity. Such studies, however, are most challenging in terms of expertise, resources, time and funding. Given the limitations of resources and the multitude of potential substrates for cytochrome P450 conversion, studies providing indirect evidence of the existence of active biotransformation seem more promising for screening purposes. Such indirect evidence may come from (1) studies designed to visualize the formation of fluorescent or colored metabolites, (2) studies based on chemical analyses of emerging metabolites, and (3) studies based on the detection of biological effects by substances activated by cytochrome P450 action.

An analysis of the relevance and adequateness of the Fish Acute Toxicity Test (AFT) according to OECD TG 203 to fulfil the information requirements and addressing concerns under REACH

The analysis of historical short-term (acute) Fish Toxicity Test (AFT) data produced according to OECD TG 203 within a dataset already used for the ECHA FET study (Scholz et al., 2016) started with 2936 studies on a total of 1842 substances from the ECHA database. Exclusion of studies with not precisely defined LC_{50} entries (e.g. only approximate information on LC_{50} range given) eliminated 54 % of all studies and 50 % of the test substances originally contained in the database. Exclusion of studies on substances that have only been tested in a single species only resulted in a remainder of 34.6 % of the original dataset and 18.8 % of substances, respectively.

Both intra- and interspecies comparisons of AFT data revealed ranges for a given substance between 1- and 1000fold. For the majority of studies, the deviation was minor, i.e. between a factor of 1 and 10. In the interspecies comparison, 72.4 % of the AFT studies produced results with deviations within a factor of 10. However, deviations by a factor between 10 and 100 (22 %) or even a factor between 100 and 1000 (5.6 %) could also be observed. Finally, only 5.8 % of the studies/substances of the intraspecies comparison gave exactly the same LC_{50} value.

Given that approx. 60 % of the existing AFT data would not fulfil the requirements set in the ECHA FET study (Scholz et al., 2016; Sobanska et al., 2018), massive re-testing of acute fish toxicity would be required. Especially study results for substances tested at concentrations close or above their water solubility, but without analytical verification should be interpreted with care or considered for revision.

Data on the AFT analysis are preliminary; analyses into effects of the application of further filtering criteria are ongoing.

Contributions to OECD project no. 2.54: "Guidance Document on an Integrated Approach on Testing and Assessment for Fish Acute Toxicity Testing" – Integration of the Fish Embryo Test into the Threshold Approach (Revision of OECD short Guidance no. 126)"

Both the literature review on biotransformation capacities in various developmental stages of zebrafish (chapter 2) and the analysis of the relevance and adequateness AFT data according to OECD TG 203 to fulfil the information requirements and addressing concerns under REACH

(chapter 3) will be integrated as major contributions to OECD project no. 2.54: "Guidance Document on an Integrated Approach on Testing and Assessment for Fish Acute Toxicity Testing" – Integration of the Fish Embryo Test into the Threshold Approach (OECD Short Guidance Document 126).

Zusammenfassung

Im Juli 2013 wurde der mit der OECD Ad hoc Expert Group on the Fish Embryo Toxicity Test und der und OECD Validation Management Group for Ecotoxicity Testing (VMG-eco) validierte akute Fischembryotoxizitätstest (FET; OECD 2013) von der Working Group of National Coordinators for the OECD Test Guidelines Programme (WNT) als OECD Prüfrichtline (TG) 236 mit dem Titel "Fish Embryo Acute Toxicity (FET) Test" verabschiedet. Da nach allgemeinem Rechtsverständnis zumindest in der Bundesrepublik Deutschland nicht selbständig fressende Entwicklungsstadien des Zebrabärblings (Danio rerio) bis zu einem Alter von 5 Tagen (Strähle et al., 2011) nicht unter den Schutz von Organsimen im Sinne der gültigen EU Directive 2010/63/EU on the protection of animals used for scientific purposes (EU, 2010) fallen, stellt das Protokoll für den Fischembryotest (FET) das erste vollständig validierte Alternativverfahren innerhalb des OECD Testrichtlinienprogramms im Bereich der Ökotoxikolgie dar. Seit seiner Akzeptanz durch die OED im Jahr 2013 ist die praktische Anwendbarkeit des Fischembryotoxizitätstests im Hinblick auf eine Reihe von Gesichtspunkten jedoch immer wieder diskutiert worden: zunächst (1) die immer wieder vermutete, aber ausgesprochen schlecht verstandene potentielle Barrierefunktion des Chorions sowie (2) die kaum untersuchte Biotransformationskapazität junger Entwicklungsstadien des Zebrabärblings. Im Hinblick auf die Anwendbarkeit des Fischembryotests z.B. im Rahmen der REACH-Richtlinien der EU wurde (3) die Eignung des FET intensiv diskutiert. Erst 5 Jahre nach der Akzeptanz des Fischembryotests als OECD-Richtlinie publizierten Sobanska et al. (2018) eine Analyse des FET-Protokolls und der zugrundeliegenden FET-Datenbank, die Anlass zur Kritik auslöste und daher in der vorliegenden Studie kritisch beleuchtet wird. Schließlich wird (4) die Eignung des Fischembryotests zur Integration in andere laufende OECD-Aktivitäten wie Projekt Nr. 2.54 ("Guidance Document on an Integrated Approach on Testing and Assessment for Fish Acute Toxicity Testing" - Integration of the Fish Embryo Test into the Threshold Approach; OECD Guidance Document 126) diskutiert. Diese vier Aspekte im Zusammenhang mit dem Fischembryotest werden in der vorliegenden Zusammenfassung adressiert.

Die potentielle Barrierefunktion des Chorions des Zebrabärblings

Seit langem bildet die Existenz des Chorions, einer extrazellulären Eischale um den Embryo des Zebrabärblings, die den Embryo auch gegen den adversen Einfluss von Chemikalien schützen soll, die Grundlage für eine Ablehnung des Fischembryotests nach OECD TG 236. Wann immer Fischembryotests Daten lieferten, die sich von jenen korrespondierender konventioneller akuter Fischtoxizitätsdaten (OECD TG 203) unterschieden, wurde der Verdacht auf eine eingeschränkte Permeabilität des Chorions des Zebrabärblings geäußert. Bereits seit dem Jahr 2008 hat das Umweltbundesamt Untersuchungen an der Universität Heidelberg zur Permeabilität des Chorions gefördert (siehe UBA-Bericht zu Projekt FKZ 37080650400); Seitdem wurden zahlreiche weitere Studien durchgeführt. Der vorliegende Bericht fasst alle Untersuchungen zusammen, die seitdem von der Arbeitsgruppe Aquatische Ökologie und Toxikologie an der Universität Heidelberg und anderen Arbeitsgruppen fertig gestellt wurden.

Im Hinblick auf die potentielle Barrierefunktion des Chorions des Zebrabärblings erbrachten Untersuchungen mit fluoreszierenden Markersubstanzen und unterschiedlich großen Polyethylenglykolen weitere Hinweise, die den Schluss zulassen, dass ein Molekulargewicht von 3000 – 4000 Dalton als Grenze für die freie Passage von ungeladenen Substanzen über das Chorion des Zebrabärblings angesehen werden kann. Die Transferrate bzw. -geschwindigkeit von geladenen Molekülen ist deutlich geringer als der Transfer von ungeladenen Molekülen. Es darf der Schluss gezogen werden, dass das Chorion des Zebrabärblings schon aufgrund der Molekulargröße für die meisten (ungeladenen) Industriechemikalien keine Barriere darstellt. In Konzentrationen ab 0,1 % reduziert Dimethylsulfoxid (DMSO) die Barrierefunktion des Chorions weiter. Da DMSO bis zu einer Konzentration von deutlich > 2 % die Überlebensrate von Embryonen des Zebrabärblings nicht beeinflusst, kann DMSO als Detergenz eingesetzt werden, um die Transferrate von Substanzen zu erhöhen, die sonst aufgrund ihrer Molekülgröße das Chorion nur langsam überwinden würden. Des Weiteren kann DMSO als Lösungsvermittler für Substanzen empfohlen werden, die aufgrund ihrer Lipophilität als in wässrigen Lösungen als schlecht löslich eingestuft werden müssten. Bei adäquater Nutzung geeigneter Lösungsvermittler gibt es bisher keine Hinweise darauf, dass hoch lipophile Substanzen nicht im Fischembryotest (FET) getestet werden könnten. Siehe hierzu auch die Empfehlungen des OECD Guidance Document (GD) Nr. 23 für schwierige Prüfsubstanzen (GD on aqueous-phase aquatic toxicity testing of difficult test chemicals).

Grundsätzlich erscheinen jedoch wegen der geringen Testdauer weder der Akute Fischtest (AFT; OECD Guideline Nr. 203) noch der Fischembryotest (FET; OECD TG 236) als geeignete Protokolle zur Prüfung hoch lipophiler Substanzen, da diese aufgrund ihrer geringen Wasserlöslichkeit und langsamen, jedoch lange anhaltenden Bioakkumulation innerhalb eines Expositionszeitraums von 4 Tagen keinen Gleichgewichtszustand erreichen dürften. Vielmehr sollte für hoch lipophile Substanzen gleich ein länger andauernder Test wie ein Early Life-Stage Test (OECD TG 210) durchgeführt werden.

Für Substanzen, die hinsichtlich ihres Durchtritts durch das Chorion des Zebrabärblings als kritisch betrachtet werden, kann die Dechorionierung in das Protokoll des Fischembryotests integriert werden: Mit nur geringen Modifikationen des FET-Protokolls können Embryonen des Zebrabärblings reproduzierbar und mit hoher Überlebenswahrscheinlichkeit (> 90 %) ab einem Alter von 24 h nach der Befruchtung dechorioniert werden. Die Dechorionierung jüngerer Entwicklungsstadien (< 24 h) ist grundsätzlich möglich, führt jedoch zu geringeren Überlebensraten.

Biotransformation und Bioaktivierungskapazität früher Entwicklungsstadien des Zebrabärblings (*Danio rerio*)

Viele Einsprüche gegen die Anwendbarkeit des Fischembryotests mit dem Zebrabärbling basieren auf der Annahme, dass frühe Entwicklungsstadien des Fisches nur über eine verminderte Biotransformationskapazität verfügen. Immer wieder wurden Zweifel geäußert, dass die metabolische Kapazität von frühen juvenilen und adulten Stadien des Zebrabärblings vergleichbar sei. Tatsächlich könnte eine begrenzte Biotransformationskapazität theoretisch zu einer Unterschätzung von Toxizität und Teratogenität von Substanzen führen, die der Bioaktivierung bedürfen. Andererseits hängen frühe Lebensstadien von Fischen in Anbetracht ihrer *Ex utero*-Entwicklung von einer intrinsischen Befähigung zur Transformation und Elimination von (exogenen und endogenen) toxischen Stoffen ab. Es scheint daher plausibel, zumindest qualitativ vom Vorhandensein einer entsprechenden Biotransformationskapazität auszugehen.

Um weiter Argumente für die Akzeptanz des Fischembryotests (FET) zu liefern, erschien zumindest der Versuch unabdingbar, die metabolischen Kapazitäten unterschiedlich alter Entwicklungsstadien des Zebrabärblings besser zu verstehen. Während unser Wissen über örtliche und zeitliche Expressionsmuster von Biotransformationsenzymen auf der Ebene der mRNA in den letzten Jahren stark zugenommen hat, befassten sich nur relativ wenige Studien mit Proteinfunktion und enzymatischer Aktivität, obwohl diese von ungleich größerer ökologischer Bedeutung als die Expression auf mRNA-Ebene sind. Grundsätzlich ist die Datenbank zum Metabolismus des Zebrabärblings auch heute noch insbesondere im Hinblick auf frühe Entwicklungsstadien deutlich unvollständiger als für andere Modellorganismen wie z.B. die Regenbogenforelle und auf über 100 wissenschaftliche Publikationen verstreut.

Im Hinblick auf ihre Biotransformationskapazitäten sind Embryonen, Juvenilstadien und Adulte des Zebrabärblings sehr unterschiedlich gut untersucht; trotz der rasanten Entwicklung des Zebrabärblings als Modellorganismus in den letzten Jahren kann unser Wissen über den Metabolismus von Xenobiotika beim Zebrabärbling bestenfalls als fragmentarisch bezeichnet werden. Da insbesondere Juvenilstadien weitgehend vernachlässigt wurden, erscheint der Transfer vor allem von Erkenntnissen bei juvenilen auf adulte Zebrabärblinge nicht statthaft. In Embryonen des Zebrabärblings konnte Biotransformation jedoch in allen Studien, die genauer ins Detail gingen, zumindest qualitativ belegt werden. Bislang konnte das Fehlen einer Biotransformation nur für seltene Ausnahmen wie Allylalkohol und Albendazol belegt werden. Grundsätzlich sind Studien zur Expression (Induktion) von Biotransformationsgenen weniger aufschlussreich als Untersuchungen zur Biosynthese der aktiven Enzyme (Transkriptomics, Proteomics) oder – noch besser – Untersuchungen zur biochemischen Aktivität der Enzyme. Da unsere Kenntnisse zum Spektrum der Substanzen, die als potentielle Substrate verschiedener Isoformen von Cytochrom P450 dienen können, sehr beschränkt sind, besteht auch nicht die Möglichkeit, von der Genaktivierung auf biochemische Funktionalität zu schließen.

Auch für den Transfer von Erkenntnissen zu Säugetieren auf Fische bzw. Fischembryonen fehlt die entsprechende Datenbasis. Für viele Isoformen von Cytochrom P450 gibt es keine ausreichenden Hinweise auf eine Kongruenz wenigstens der Terminologie von Isoformen von Cytochrom P450 bei Säugetieren und niederen Wirbeltieren. Aus diesem Grund ist ein Transfer von Nomenklatur und metabolischer Funktion ("gleicher Name bedeutet gleiche Funktion") von Säugetieren auf Fische (oder umgekehrt) nicht zulässig. Im Hinblick auf Enzyme der Biotransformation Phase II ist unser Wissen mit noch größeren Lücken behaftet als für Enzyme der Phase I.

Als Goldstandard für Biotransformationsstudien mit Embryonen, Juvenilstadien und Adulten von Fischen müssen Genexpressionsstudien mit Daten der Proteomanalyse und Analysen der gebildeten Metaboliten, die bei den biochemischen Reaktionen gebildet werden, verlinkt werden. Solche Studien stellen jedoch im Hinblick auf Expertise, Ressourcen, Zeit und Finanzierung eine erhebliche Herausforderung dar. In Anbetracht der Beschränkungen von Ressourcen und der Vielzahl potentieller Substrate für Cytochrom P450 erscheinen Studien zum indirekten Nachweis der Existenz aktiver Biotransformation für Screenings mehr zu versprechen und schneller zum Ziel zu führen als der Nachweis von Genaktivierung, Proteinsynthese und -aktivität für eine Vielzahl einzelner Isoformen von Cytochrom P450. Derartige indirekte Hinweise können sich ergeben aus (1) dem optischen (mikroskopischen) Nachweis der Bildung fluoreszierender oder gefärbter Metaboliten, (2) dem chemischanalytischen Nachweis der gebildeten Metaboliten sowie (3) dem Nachweis der gebildeten Metaboliten über ihre biologische Wirkung nach entsprechender Bioaktivierung von zunächst untoxischen Substraten. Die vorliegende Studie stellt Beispiele für diese indirekten Methoden vor.

Analyse der Relevanz und Eignung von Daten aus akuten Fischtests (AFT) nach OECD TG 203 für die Erfüllung der Informationsanforderungen unter REACH

Die Analyse eines historischen Datensatzes zur akuten Fischtoxizität (AFT) auf der Basis von Tests nach der OECD TG Nr. 203, der bereits die Grundlage für die Analyse in einer von der Europäischen Chemikalienagentur (ECHA) in Auftrag gegebenen Studie zur Eignung des Fischembryotests (OECD TG 236) gebildet hatte (Scholz et al., 2016), ging ursprünglich von 2936 Studien mit insgesamt 1842 Substanzen aus. Der Ausschluss von Studien, die keine präzisen Angaben zum LC₅₀ (z.B. nur näherungsweise Angaben zu einem ungefähren LC₅₀-Bereich) machten, führte zu einer Elimination von 54 % aller Studien sowie 50 % der Testsubstanzen, die ursprünglich im Datensatz enthalten waren. Der Ausschluss von Studien zu Substanzen, die nur in einer einzelnen Fischspezies durchgeführt worden waren, beließ noch 34,6 % bzw. 18.8 % der ursprünglich im Datensatz enthaltenen Studien bzw. Substanzen in der weiteren Untersuchung.

Sowohl Intra- als auch Interspeziesvergleiche von AFT-Daten ergaben für bestimmte Substanzen Schwankungen der Daten um Faktoren bis zu 1000. Für die Mehrheit der Substanzen waren die Abweichungen marginal, d.h. im Bereich von Faktoren zwischen 1 und 10. So ergab der Interspeziesvergleich für 72,4 % der Fälle Abweichungen mit einem Faktor bis zu 10. Jedoch konnten für 22 bzw. 5,6 % auch Abweichungen um Faktoren zwischen 10 und 100 bzw. zwischen 1000 und 10000 beobachtet werden. Letztendlich konnte im Rahmen des Intraspeziesvergleichs für nur 5,8 % der Studien bzw. Substanzen der gleiche LC₅₀-Wert ermittelt werden.

Die Tatsache, dass ca. 60 % der existierenden AFT-Daten die Anforderungen nicht erfüllen, die von Scholz et al. (2016) sowie Sobanska et al. (2018) vor dem Hintergrund von REACH an den Fischembryotest (FET) gestellt wurden, scheint eine umfangreiche Neutestung von Substanzen für REACH – auch mit dem akuten Fischtest nach OECD TG 203 – erforderlich zu machen. Insbesondere Studien im Bereich von Substanzkonzentrationen nahe oder über der Löslichkeitsgrenze der Substanzen erfordern besondere Aufmerksamkeit und sollten notfalls erneut überprüft werden.

Die Daten zur Analyse der AFT-Daten sind vorläufig; die Studie wird weitergeführt.

Beiträge des Fischembryotests (FET) zu OECD-Projekt Nr. 2.54: "Guidance Document on an Integrated Approach on Testing and Assessment for Fish Acute Toxicity Testing" – Integration of the Fish Embryo Test into the Threshold Approach (OECD Guidance Document 126)"

Sowohl die Literaturstudie zur Biotransformationskapazität in verschiedenen Entwicklungsstadien des Zebrabärblings (Kapitel 2) als auch die Analyse der Relevanz und Eignung des akuten Fischtests (AFT) nach OECD TG 203 für die Erbringung der notwendigen Informationen und die Behandlung von Bedenken unter REACH (Kapitel 3) werden als wichtiger Beitrag in das OECD-Projekt "Guidance Document on an Integrated Approach on Testing and Assessment for Fish Acute Toxicity Testing" – Integration of the Fish Embryo Test into the Threshold Approach (OECD Guidance Document 126)" integriert.

1 The zebrafish (*Danio rerio*) chorion as a potential barrier for the uptake of xenobiotics into the zebrafish embryo

1.1 Introduction

For long, a major objection to the acceptance of the Fish Embryo Test (FET) according to Organization for Economic Co-operation and Development (OECD) Test Guideline (TG) 236 has been the putative protecting role of the chorion (eggshell) of the embryo. Whenever fish embryo toxicity tests produced results different from data published for the conventional Acute Fish Toxicity test (AFT) according to OECD TG 203, impaired permeability of the zebrafish chorion has been speculated. Starting as early as in 2008, the German Environmental Agency (UBA) has funded investigations into the permeability of the zebrafish chorion at the University of Heidelberg (cf. UBA report FKZ 37080650400). Since then, more-in-depth experiments have been carried out. The present report provides an integration of all investigations conducted so far at the Aquatic Ecology and Toxicology Group in the Center for Organismal Studies at the University of Heidelberg and elsewhere.

1.2 Development of the zebrafish chorion

Since in most teleosts fertilization of eggs takes place in the free water body, fish eggs are shed during the spawning act into the free water spaces, where they are fertilized by simultaneously released sperm (Fedderwitz, 2008; Hisaoka, 1958; Kais, 2009). As a consequence, the eggs of non-live-bearing fish are protected by an outer shell, the chorion, against mechanical damage, infection by parasites and potentially also against chemical insult. Genetically and developmentally important, the chorion protects against polyspermy (Donovan and Hart, 1983). In zebrafish, sperm lack acrosomal structures to penetrate the chorion; in contrast, the egg chorion has a preformed permeation site named micropyle (Mold et al., 2001). In unfertilized zebrafish eggs, the micropyle area has an average diameter of $2.1 - 2.5 \mu m$ (Donovan and Hart, 1983), and the area for penetration of the sperm is limited to approx. 100 μm , where cortical granules have accumulated under the plasmalemma.

So far, the chorion of teleosts is thought to be a primary sheath (Brivio et al., 1991), i.e. the primary chorion is formed by the oocyte itself (Yamagami et al., 1992). Secondary shells are produced by follicular cells, and tertiary shells are produced by other reproductive organs such as the oviduct (Yamagami et al., 1992). As mentioned, the teleost chorion prevents multiple fertilization (Gilkey et al., 1978). The morphological structure and chemical composition of the fish chorion significantly varies with species (Brivio et al., 1991), most likely as a consequence to adaption to various environments (Hart et al., 1984).

In the zebrafish, the chorion has an average diameter of $1.5 - 2.5 \mu m$ (Rawson et al., 2000) and consists of three layers: an external electron-dense layer (Zona radiata externa), an intermediate fibrillary layer (upper Zona radiate interna) and an inner layer made up of 16 horizontal electron-dense sublayers interspersed with 15 sublayers of lower electron-density (deep Zona radiate interna) (Hart et al., 1984). However, the innermost layer of the eggshell has also been given various synonymous names: translucent membrane, vitelline membrane, vitelline sheath, zona pellucida, chorion, primary, secondary and tertiary shell (Arukwe and Goksoyer, 2003). The outer, intermediate and inner layers of the zebrafish chorion measures about 0.2 - 0.3, 0.3 - 0.6 and 1.0 - 1.6 μm in thickness, respectively (Rawson et al., 2000).

The chorion of fish is produced during oocyte maturation within the female gonads (Selman et al., 1993):

(1) In oocyte maturation stage IA (pre-follicular, primary growth stage), oocytes are arranged in nests of oocytes and measure 7 - 20 μ m in diameter; a chorion cannot yet be discerned. Stage IB is characterized by interdigitating microvilli formed by both the oocytes and the follicular cells as well as secretion of electron-dense materials secreted from either side. This first manifestation of a chorion shows only one single layer and reaches a thickness of approx. 0.15 μ m (Fig. 1a).

(2) In stage II (cortical alveolus stage), the oocyte diameter reaches $0.14 - 0.34 \mu m$ and is surrounded by a translucent layer of flattened to cubic follicular cells extending long microvilli towards the oocytes (Fig. 1b). The chorion itself has reached almost 6 μm in thickness and consists of 2 - 3 layers built up by several sublayers. The second layer has formed between the primary layer and the oolemma (cell membrane of the oocyte), whereas the third layer is being formed alongside the oolemma. Pores across all of these layers are kept open by microvilli from both the oocyte and the follicular cells.

(3) During oocyte maturation stage III (vitellogenesis), zebrafish oocytes may reach diameters of $340 - 690 \,\mu\text{m}$ (Fig. 2). The cubic translucent follicular cells surround a chorion of at least three layers; however, the chorion starts to attenuate, but microvilli still keep the chorion pore canals open.

(4) During stage IV (maturation), the follicular cells withdraw their microvilli from the pore canals, which then close (Donovan and Hart, 1983). The chorion diameter further declines, and oocyte have reached $690 - 730 \,\mu$ m in diameter.

(5) In stage V (mature oocyte), the chorion has reduced its diameter further to < $300 \mu m$. Under the electron microscope, the pores appear empty; however, as can be visualized under the light microscope, the distal endings of the pore canals are obstructed by some dense materials. Follicular cells can no longer be discerned, and the eggshell has become translucent. Upon spawning and contact to the external medium water, the vitelline membrane detaches from the oocyte plasma membrane and forms the final inner chorion layer, the perivitelline space (Coward et al., 2002).



Figure 1: Oocytes of zebrafish (Danio rerio)

Non-fertilized oocytes of zebrafish (Danio rerio) in stage I (primary growth; left) and stage II (cortical alveolus; right).

Given its structure, the zebrafish chorion should be permeable to water, electrolytes and smaller hydrophilic organic substances (Hisaoka, 1958; Hisaoka and Firlit, 1960). In fact, water permeability strongly depends on the age of the oocytes: After fertilization and a subsequent phase of water hardening, the permeability for water rapidly declines (Hagedorn et al., 1997a). Correspondingly, as early as 1 h after fertilization, zebrafish eggs show a significantly reduced sensitivity to effluents (Gellert and Heinrichsdorff, 2001). Using deuterium-labelled water, embryos up to the 2-cell stage were found to take up water much more slowly than subsequent stages, i.e. water permeability is significantly increased after water hardening of the chorion (Adams et al., 2005).



Figure 2: Zebrafish (Danio rerio) oocytes in maturation

Non-fertilized zebrafish (*Danio rerio*) oocyte in maturation stage III (vitellogenesis). The chorion consists of three layers and starts to attenuate.

As a consequence of hardening, the mechanical stability of zebrafish eggs also increases: Penetration of the chorion by means of a micropipette requires much higher forces in the pharyngula stage than in the gastrula stage, and as a blastula, the egg is more difficult to penetrate than before spawning. (Kim et al., 2005). Overall, the mechanical resistance of the zebrafish egg declines with development. Artificial softening of the chorion can easily be effected by application of proteolytic pronase (chorionase), an enzyme released immediately before the natural onset of the hatching process (Kim et al., 2005). During normal embryo development, chorionase is secreted by hatching gland cells located in a belt-like arrangement along the anterior side of the yolk sac (Inohaya et al., 1997; Schoots et al., 1982). Since chorionase does not digest the outermost layers of the chorion, premature hatching in consequence of chorionase secretion by adjacent oocytes is prevented.

There is a controversial discussion about the permeability of the zebrafish chorion for organic compounds: Whereas Zhang und Rawson (1996) documented the chorion to be freely permeable for water and "small" molecules, the free permeability has been questioned for later developmental stages (Adams et al., 2005; Harvey et al., 1983). Likewise, the relative contribution to a potential barrier function of the chorion itself and the perivitelline space is under discussion (Harvey et al., 1983; Zhang and Rawson, 1996). Finally, the syncytial layer of the yolk has also been discussed with respect to its contribution to a barrier function (Hagedorn et al., 1997b).

1.3 Structure of the zebrafish chorion

Whether or not the pores are exclusively responsible for the transfer of materials (chemicals) across the chorion is not clear. The spacing of pores on the zebrafish chorion is quite regular (Figs. 3, 4). In the electron microscope, the pores can be visualized as prominent structures across the multi-layered chorion (Fig. 5). In the scanning electron microscope, the surface of the zebrafish chorion displays various depressions and protrusions (Fig. 6). In the scanning electron microscope, broken edges and inner surface views of the chorions document that the pore canals are obstructed close to the outer surface of the chorion (Figs. 6 - 9). For more details, see Fedderwitz (2008).



Preparation of the zebrafish chorion: A) the chorion of a 48 hpf embryo was torn apart along the pores (Alcian blue- Periodic acid-Schiff (PAS) stain, $40\times$); B) outer surface of the chorion with distinctive projections brightly colored (hematoxylin eosin (HE) stain, $60\times$); C) outer chorion surface of a 72 hpf zebrafish embryo (Masson-Goldner, $60\times$); D) inner chorion surface of a 72 hpf zebrafish embryo (Masson-Goldner, $60\times$); E) micropyle (Masson-Goldner, $40\times$); F) outer chorion surface (Alcian blue-PAS-stain, $60\times$). From Böhler (2012).



Figure 4: Surface of the chorion of zebrafish (*Danio rerio*) eggs

Surface views of the chorion of zebrafish (*Danio rerio*) eggs at 24 h after fertilization clearly show the regular dispersion of the chorion pores (\rightarrow). Staining techniques: A: Goldner, B. Azan, C: HE, D: PAS. From Fedderwitz (2008) and Braunbeck et al. (2012).



Figure 5: Layers of the zebrafish (*Danio rerio*) chorion

The multi-layered zebrafish (*Danio rerio*) chorion (Ch) shows multiples pores (P), which are penetrated by microvilli (M) both from the follicular cells (FZ) and the oocyte (O). a: outer apical layer; m: intermediate layer; i: inner layer; d: dense material. Magnification left: 20,000×; right: 6,600×. From Fedderwitz (2008) and Braunbeck et al. (2012).





Outer surface of the zebrafish (*Danio rerio*) chorion in the scanning electron microscope: 12 h after fertilization, the surface displays shallow indentations as well as larger shallow protrusions. 60 h after fertilization, small bubble-like protrusions are visible. A: 12 h; B: 24 h; C: 48 h; D: 60 h. d: electron-dense materials. From Fedderwitz (2008) and Braunbeck et al. (2012).

Figure 7: Inner surface of the zebrafish (*Danio rerio*) chorion



Along broken edges (left) and from the inner surface (right) of the zebrafish (*Danio rerio*) chorion, chorion pores become evident at regular spacing as canals that do not completely penetrate the chorion, but are plugged close to the outer surface. From Fedderwitz (2008).





Left: Field-emission scanning electron microscopy of the outermost granular layer (GL) of the zebrafish (*Danio rerio*) chorion with distinct projections $(2.0 - 3.0 \,\mu\text{m}$ in diameter) at intervals of $2 - 10 \,\mu\text{m}$. Bar: $2 \,\mu\text{m}$. Right: Field-emission scanning electron microscopy view of the outer chorion surface, with the granular layer (GL) and inner surface with openings of the chorion pore canals (CPC). At a few places, remnants of materials from the perivitelline space (PVR) are attached to the inner side of the chorion. Bars: $10 \,\mu\text{m}$. From Rawson et al. (2000).





Field-emission scanning electron microscopy of the openings of the pore canals in the chorion of zebrafish (*Danio rerio*). Bars: 1 µm. From: Rawson et al. (2000).

1.4 Dechorionation as a tool to improve the fish embryo toxicity test (FET) with the zebrafish (*Danio rerio*)

The most obvious approach to investigate the role of the chorion as a barrier for chemical toxicity is to experimentally remove the chorion and expose the "naked" embryo. Several protocols exist for fish embryo dechorionation, using either trypsin/EDTA (Collodi et al., 1992) or pronase (Stuart et al., 1990; Westerfield, 2007) solutions to digest or even dissolve the chorion. Most studies aimed at recovering embryos for genetic engineering or generating embryonic cells for cell cultures, but only few investigations were carried out to reveal the influence of chemicals on dechorionated embryos (Braunbeck et al., 2005; Mizell and Romig, 1997; Ozoh, 1980). According to Westerfield (2007), embryos can also be dechorionated mechanically by use of forceps without any enzymatic digestion (Fig. 10). For the specific purposes of these studies, it seemed sufficient to select undamaged embryos or embryonic cells for further operations. In none of the studies mentioned above, however, the actual survival rates of dechorionated embryos were given special attention and subjected to statistical analysis. For chemical toxicity assessment, however, a reproducibly high survival rate is of fundamental importance. Therefore, the present study was designed to develop and optimize a dechorionation procedure with reproducibly high survival rates. The success of the dechorionation procedure was tested with a cationic polymer (Luviquat HM 552), which has been suspected to be blocked by the chorion (Leonard et al., 2005).



Figure 10: Dechorionation of zebrafish (Danio rerio) embryos

Mechanical dechorionation of zebrafish (*Danio rerio*) embryos using Dumont[™] no. 5 forceps: A: intact embryo 6 h after fertilization; B: egg fixed with one pair of forceps; C: piercing the chorion with a second pair of forceps; D: enlarging the cleft by pulling apart the chorion by means of two forceps; E: chorion with cleft before lifting; and F: dechorionated embryo. Average diameter of the embryo with chorion: 0.7 mm.

After some optimization, zebrafish embryos could be dechorionated from an age of 24 h after fertilization (Henn and Braunbeck, 2011) with satisfactorily high survival rates (Fig. 11). For the model high molecular weight substance Luviquat, it could be demonstrated that access of high molecular weight molecules could be given access to the zebrafish embryo by means of dechorionation (Figs. 12, 13).



Figure 11: Correlation of zebrafish (Danio rerio) age and survival after dechorionation

Correlation between the age of zebrafish embryos at time of dechorionation and the ratio of normal development versus mortality as well as sublethal effect rates on the second day after dechorionation. For comparison, data for non-dechorionated embryos have been added.*Results statistically different compared to the group dechorionated 24 hpf (Dunn's method, p < 0.05). From Henn and Braunbeck (2011) and Braunbeck et al. (2012).

Figure 12: Mortalities of non-dechorionated zebrafish (*Danio rerio*) embryos after exposure to Luviquat



Mortalities of non-dechorionated zebrafish (*Danio rerio*) embryos (—) after 48 (\bullet) and 72h (\circ) and corresponding hatching rates (- -) after 48 h ($\mathbf{\nabla}$) and 72 h (∇) of exposure to Luviquat. From Henn and Braunbeck (2011) and Braunbeck et al. (2012).

Figure 13: Mortalities of dechorionated and non-dechorionated zebrafish (*Danio rerio*) embryos after exposure to Luviquat



Comparison of mortalities after 120 h of exposure to Luviquat in non-dechorionated zebrafish (*Danio rerio*) embryos (\bullet), after 48 h of exposure to Luviquat in eleutheroembryos exposed from the age of 72 h (total age of embryos: 120 h; ∇), and after 48 h of exposure to Luviquat in embryos dechorionated at 24 h post fertilization (total age of embryos: 72 h; \blacksquare). From Henn and Braunbeck (2011) and Braunbeck et al. (2012).

For more details and a detailed protocol for zebrafish dechorionation with reproducibly high survival rates, see Henn and Braunbeck (2011).

1.5 Fluorescent dyes as markers for the permeability of the chorion in zebrafish (*Danio rerio*) embryos

There is limited evidence that the permeability of the chorion changes during the embryonic development and that, after hardening, the chorion is less permeable (Gellert el al. 2001) and might then function as a barrier for even smaller molecules. Ozoh (1980) found dechorionated zebrafish embryos to be more sensitive against copper intoxication than non-dechorionated embryos, which corroborated several other studies with different teleost species showing accumulation of different heavy metals on the outside or within the chorion (Stouthart et al., 1994; Van Leeuwen et al., 1985; Wedemeyer, 1968). Only few studies indicated a weak barrier function of the zebrafish chorion for lipophilic substances such as γ -hexachlorocyclohexane (lindane) (Braunbeck et al., 2005) and small substances such as dimethyl sulfoxide (DMSO) (Harvey et al., 1983). In other teleost species like the medaka (*Oryzias latipes*), the same seems to hold true for substances like cypermethrin and thiobencarb (Gonzalez-Doncel et al., 2004; Villalobos et al., 2000).

An essential aspect of the permeability of the fish egg chorion is the molecular weight. Polymers and high molecular weight surfactants (~ 40,000 – 100,000 g/mol) were suspected to be blocked by the chorion, since comparison between embryo, eleutheroembryo toxicities (Leonard et al., 2005) and dechorionated embryos resulted in considerably higher sensitivity of the latter both (Henn and Braunbeck, 2011). Creton (2004) investigated the inhibition of the endoplasmic reticulum (ER) Ca2+ pump in zebrafish embryos and established that the chorion is permeable to fluorescent dextrans of 3000 Da, but is not permeable to fluorescent dextrans of 10,000 Da. To identify a "critical molecular size" for the passage across the chorion, polymers are ideal test substances: a polymer is a macromolecule composed of repeating identical structural units typically connected by covalent chemical bonds, which are available in various molecular weights. Since the basic unit is always the same, no additional functional groups that might interact with the chorion are introduced. Due to the size evidences of between 3 kDa and 10 kDa (Creton, 2004), Kais (2009) tested fluorescent dextran between 3 and 40 kDa with respect to their ability to pass the chorion.

The use of fluorescence dyes allows the visualization of test substances and, thus, their uptake into the egg as well as its accumulation can be followed. Wavelength and amount of the emitted energy depend on both the fluorophore and the chemical environment of the fluorophore (Tsien et al., (eds) 1995). The two well-known classes of highly fluorescent dyes, rhodamines and fluoresceins, are derivatives of xanthenes and belong to heterocyclic compounds. In addition to their similarity to PAHs, rhodamine and fluorescein are ecologically relevant by themselves. Uranin is the water-soluble sodium salt of fluorescein and has been used in river systems, most notably in the case of the Chicago River, where fluorescein was the first substance used to dye a river green on St. Patrick's Day in 1962. In industrial applications, it is used to color automotive coolants (anti-freeze), for dying wool and silk, as an Air-Sea rescue marker, as ground water tracing dye, soap solutions and for the coloring of agrochemicals and fertilizers. Rhodamines are used as dyes for paper and textiles, in luminous pigments, in dye lasers, as well as in cell biology, where they have served as markers in fluorescence microscopy.

Apart from water solubility and partition coefficient, other parameters have to be considered, e.g. Topological Polar Surface Area (TPSA), which is an estimation of the polar fraction of the molecular surface area. Similar to the physicochemical parameters such as the influence of the particular structure and its complexity, it affects the environmental impact and toxicological potential. The complexity rating of the compounds is a rough estimate of how complicated a structure is in terms of both elements and the displayed structural features including symmetry. Generally, larger compounds are more complex than smaller ones, but highly symmetrical compounds or compounds with few distinct atom types or elements are downgraded.

The diploma thesis by Kais (2009) was designed to investigate the mechanism(s) of uptake and distribution of fluorescent dyes with different chemical characteristics in the zebrafish embryo. Lipophilicity, molecular weight, different substituents, as well as different charging are important issues for ecotoxicological questions. Rhodamine b and sulforhodamine b were representatives for good water solubility and low log POW. Fluorescein is a small molecule and is regarded as a basic molecule. This study investigates how the chemical behavior of fluorescein might change with the insertion of two chlorine atoms or with an additional carboxyl group. Embryos were studied after 24 h and 48 h in the standard fish embryo toxicity test (FET). The time between 26 h and 48 h (called pre-hatching time) is especially important for the question of distribution due the fact that the structure of the chorion profoundly changes during this developmental period. Experimental results provide evidence of massive quantitative changes in the uptake and distribution following the chorion softening process, which is mostly due to the proteolytic activity at the pre-hatching stage (Kim et al., 2005).

In fact, fairly water-soluble molecules of lower molecular weight such as rhodamine rapidly penetrate the zebrafish chorion and accumulate within the embryo (Fig. 14). Although only of slightly higher molecular weight (580 versus 479) and an even lower KOW value (1.44 versus 2.28), sulforhodamine is transferred much more slowly than rhodamine, most likely due to (1) the overall negative charge of sulforhodamine and (2) the bulky nature of the two carboxyl groups (Fig. 14). In addition to molecular size and charge, exposure time is a critical factor for the transfer of substances across the zebrafish chorion (Fig. 15). Most importantly, solvents with detergent-like properties such as dimethyl sulfoxide massively facilitate the transfer of

substances with moderate to higher lipophilicity such as fluorescein (Fig. 16). However, it should be noted that DMSO also supports the uptake of markers into dechorionated embryos, i.e. the solvent not only modifies the properties of the chorion, but also of the zebrafish embryo itself (Fig. 17). For more details, see Kais (2009) and Kais et al. (2013).





Rhodamine and sulforhodamine transport across the zebrafish (*Danio rerio*) chorion after 48 h water-borne exposure: Since the accumulation of rhodamine inside the embryo by far exceeds that of sulforhodamine, there is evidence that positively charged molecules show lower penetration rates across the zebrafish chorion. From Kais (2009).

Figure 15: Time-dependent passage of the fluorescent marker dichlorofluorescein across the zebrafish (*Danio rerio*) chorion



Prolonged exposure to the fluorescent marker dichlorofluorescein increases the transfer across the zebrafish (*Danio rerio*) chorion and the accumulation inside the embryo, thus documenting the time-dependence of the passage of materials across the chorion. From Kais (2009) and Braunbeck et al. (2012).





Increased concentrations of the solvent (detergent) dimethyl sulfoxide (DMSO) facilitate the uptake of fluorescent markers such as fluorescein across the chorion of the zebrafish (*Danio rerio*) embryo. The use of detergent may thus be a tool to facilitate / improve the passage of materials across the fish chorion. From Kais (2009) and Braunbeck et al. (2012).



Figure 17: Impact of DMSO on substance uptake into zebrafish (Danio rerio) embryos

DMSO strongly increases the uptake of the fluorescent marker fluorescein into dechorionated zebrafish (*Danio rerio*) embryos, ie. the solvent not only modifies the properties of the chorion (cf. Fig. 16), but also of the zebrafish embryo itself. From Kais (2009) and Braunbeck et al. (2012).



Figure 18: Epifluorescence images of zebrafish (Danio rerio) embryos

Epifluorescence images of zebrafish (*Danio rerio*) embryos at the age of 24 hpf (A, C, E) and 48 hpf (B, D, F) exposed to 100 mg/L fluorescein dissolved in 1% (A, B), 0.1% (C, D) and 0.01% DMSO (E, F): With 0.1 and 1% DMSO, fluorescein showed a strong signal inside the embryo (A–D); especially in the common cardiac vein (=duct of Cuvier) (\blacktriangleright) including the heart (+) and vessels inside the brain (*). In contrast, with 0.01% DMSO, fluorescein fluorescence inside the embryo was weak, but still quite high (E and F). In the embryo, fluorescence intensity increased with DMSO concentration (A > C > E). Exposure time: 30 ms (24 hpf) and 2 ms (48 hpf) (Kais et al., 2013).



Figure 19: Cross-sections and surface images of zebrafish (Danio rerio) embryos

Cross-sections (A, C, E) and composite three-dimensional surface (B, D, F) confocal laser scanning images of zebrafish (*Danio rerio*) embryos at the age of 24 hpf after exposure to 100 mg/L fluorescein dissolved in 1% (A and B), 0.1% (C and D) and 0.01% DMSO (E and F): With 0.1 and 1% DMSO, fluorescein passed the chorion and accumulated within the embryo (A and C); in contrast, with 0.01% DMSO, fluorescein did not pass the chorion, and no signal in the embryo was detectable (E). All images were taken at 100% laser power and a high voltage setting of 20. From Kais et al. (2013) and Braunbeck et al. (2012).



Figure 20: Transmission, cross-section and 3D images of zebrafish (Danio rerio) embryos

Transmission detector (A, D, G) as well as cross-section (B, E, H) and 3D maximum intensity rendering (C, F, I) confocal laser scanning images of zebrafish (*Danio rerio*) embryos (48 hpf) after exposure to 100 mg/L fluorescein dissolved in 1 % (A - C), 0.1% (D - F) and 0.01 % DMSO (G - I): In 0.1 and 1 % DMSO, showed a strong signal inside in the embryo (A–F), which clearly outshined the signal of the chorion and a small portion of fluorescein in the intestinal lumen (G - I). Fluorescence intensity inside the embryo decreased with DMSO concentrations (C > F > I). Fluorescence images were taken at 100 % laser power and a high voltage setting of 1 (B, C, E, F) and 20 (H and I). From Kais et al. (2013) and Braunbeck et al. (2012).



Figure 21: Composite 3D surface confocal laser images of zebrafish (Danio rerio) embryos

Composite three-dimensional surface (maximum intensity rendering) confocal laser images of zebrafish (*Danio rerio*) embryos, dechorionated after 24 h (A, C, E) and 48 h (B, D, F) exposure to 100 mg/L fluorescein dissolved in 1% (A and B), 0.1 % (C and D) and 0.01 % DMSO (E and F): With 0.1 and 1 % DMSO, fluorescein gave a strong signal inside the embryo (A - D); with 0.01 % DMSO, fluorescein fluorescence inside the embryo (E and F) and in a small section of the intestinal lumen (F) was weak. Fluorescence images of 24 hpf embryos were taken at full laser power and a high voltage setting of 20 (A), 30 (C) and 150 (E), whereas 48 hpf embryo images were taken at laser power of 5 (B), 3 (D) and 100 (F) and a high voltage setting of 15 (B and D) and 100 (F). From Kais et al. (2013) and Braunbeck et al. (2012).
Figure 22: Cross-section and composite 3D surface images of zebrafish (*Danio rerio*) embryos after exposure to 2,7-dichlorofluorescein I



Cross-section (A and C) and composite 3D surface (B, D - F) confocal laser images of zebrafish (Danio rerio) embryos at the age of 48 hpf after exposure to 50 mg/L 2,7-dichlorofluorescein in 1 % (A and B), 0.1 % (C and D) and 0.01 % DMSO (E and F): In 0.1 and 1 % DMSO, there was an increased fluorescence intensity of 2,7-dichlorofluorescein inside the embryo (A - D); 0.01 % DMSO, however, did not show a definite change in the signal strength (F). All images were taken at full laser power and a high voltage setting of 10. From Kais et al. (2013) and Braunbeck et al. (2012).



Figure 23: Cross-section and composite 3D surface images of zebrafish (*Danio rerio*) embryos after exposure to 2,7-dichlorofluorescein II

Composite three-dimensional surface confocal laser images of zebrafish (*Danio rerio*) embryos dechorionated after 24 h (A, C, E) and 48 h (B, D, F) exposure to 50 mg/L 2,7-dichlorofluorescein dissolved in 1 % (A and B), 0.1 % (C and D) and 0.01 % DMSO (E and F): Following exposure in 0.1 and 1 % DMSO, 2,7-dichlorofluorescein gave an increased signal inside the embryo (A - D), especially in the intestinal lumen. In contrast, with 0.01 % DMSO, fluorescence intensity was much lower (E and F). All fluorescence images were taken at full laser power and a high voltage setting of 50 (24 hpf) and 5 (48 hpf) (Kais et al., 2013) and Braunbeck et al. (2012).

1.6 Polyethylene glycols of different molecular size as markers of chorion permeability in the zebrafish (*Danio rerio*) embryo

In order to more closely identify the upper limits of the molecular size of chemicals to cross the chorion of zebrafish, differently sized, non-toxic and chemically inert polyethylene glycols (PEGs; 2000 - 12,000 Da) were applied at concentrations (9.76 mM) high enough to provoke osmotic pressure. Whereas small PEGs were expected to be able to cross the chorion, restricted uptake of large PEGs was hypothesized to result in shrinkage of the chorion (Fig. 24). Due to a slow, but gradual uptake of PEGs over time, molecular size-dependent equilibration in conjunction with a regain of the spherical chorion shape was observed. Thus, the size of molecules able to cross the chorion could be narrowed down precisely to \leq 4000 Da, and the time-dependency of the movement across the chorion could be described. To account for associated alterations in embryonic development, fish embryo toxicity tests (FETs) according to OECD TG 236 (OECD, 2013) were performed with special emphasis to changes in chorion shape.

FETs revealed clear-cut size-effects: the higher the actual molecular weight (= size) of the PEG, the more effects (both acutely toxic and sublethal) were found (Fig. 25). No effects were seen with PEGs of 2000 and 3000 Da. In contrast, PEG 8000 and PEG 12,000 were found to be most toxic with LC_{50} values of 16.05 and 16.40 g/L, respectively. Likewise, the extent of chorion shrinkage due to increased osmotic pressure strictly depended on PEG molecular weight and duration of exposure. A reflux of water and PEG molecules into the chorion and a resulting reshaping of the chorion could only be observed for eggs exposed to PEGs \leq 4000 Da. Results clearly indicate a barrier function of the zebrafish chorion for molecules larger than 3000 to 4000 Da.



Figure 24:Shrinkage of the chorion of zebrafish (Danio rerio) embryos after transfer into
polyethylene glycols of different molecular mass

After transfer into PEG test solutes, the chorion of zebrafish (*Danio rerio*) embryos showed a concentrationdependent shrinkage due to effects by PEG 3000 (A - C) and PEG 6000 (D - F). Only minor depression of the chorion were found at 3.125 g/L PEG 3000 (A) and PEG 6000 (D), whereas at 25 g/L (B, E) and 100 g/L (C, F) stronger depressions were evident, finally leading to a tightly wrapped chorion around yolk and cells (C and F). From Pelka et al. (2017).





Sublethal effects in zebrafish (*Danio rerio*) embryos after 96 h exposure to 12.5 g/L (A, B), 25 g/L (C) and 50 g/L PEG 6000 (D). Note the absence (A) or deformation of the eyes (B - C), sharply bent tails, formation of edemata as well as yolk deformations of PEG-exposed embryos. From Pelka et al. (2017).

In any case, the PEG-size-dependent deformation of the chorion showed a clear-cut doseresponse relationship. In addition, this could be correlated with a concentration-dependent increase of acute lethality and sublethal effects (Fig. 26).

The minor (PEG 2000 and PEG 3000) to strong deformations of the chorions (PEGs \geq 4000 Da) observed immediately after the onset of exposure gradually changed with progressing exposure. Whereas eggs exposed to PEG 2000 quickly regained the spherical shape of their chorions (Fig. 27), the recovery of chorion shape of embryos exposed to PEG 3000 took 48 h. For PEG 4000, minor recovery from chorion deformation could be observed. Since the space between the chorion and the embryo slightly increased within 48 h, the embryos were able to resume their movements, even within the deformed chorions. For PEGs \geq 6000 Da, however, the deformations of the chorion observed at the very start of exposure seemed to even increase after 24 and 48 h, indicating a continuation of the shrinkage process. Finally, embryos exposed to PEGs 8000 and 12,000 did not develop further and quickly coagulated (Fig. 27).

In order to graphically illustrate the PEG size- and time-dependence of the equilibration process, the average difference of the chorion area at different time points in relation to the area at the start of exposure was determined (Fig. 28). Again, two groups of PEG size ranges could be distinguished: For PEGs \leq 4000, an increase of the chorion area could be observed, indicating a recovery from the deformation due to gradual influx of external medium. The biggest difference compared to the start of exposure was found for PEG 3000 with a difference in chorion area of 13.16, 14.44 and 16.52 % after 24, 36 and 48 hpf, respectively. In contrast, for PEGs \geq 6000 a decrease of chorion area was evident. For embryos exposed to PEGs 8000 and 12,000, the chorion area even decreased by 1.54 and 2.18 %, respectively. For more details, see Pelka et al. (2017).



Figure 26: Chorion deformation of zebrafish (Danio rerio) eggs after exposure to PEGs

Exposure of zebrafish (*Danio rerio*) eggs for 24 h to different polyethylene glycols (PEGs) reveals chorion deformation (grey columns) at almost all concentrations of the differently sized PEGs. Likewise, for all tested PEGs, there was a concentration-dependent increase of acute lethality (solid black line) and sublethal effects (dotted blue line). From Pelka et al. (2017).



Figure 27:Chronological sequence of deformation of the zebrafish (Danio rerio) chorion
following exposure to differently sized polyethylene glycols (PEGs)

Chronological sequence of deformation of the zebrafish (*Danio rerio*) chorion following exposure to differently sized polyethylene glycols (PEGs). Each column shows the same egg exposed to dilution water (ctrl) or 9.67 mM of the different PEGs at start of exposure (approx. 1 hpf) as well as at 24 and 48 hpf. Whereas the chorions of embryos exposed to PEG 2000 and 3000 regained a spherical shape, the chorions of PEGs \geq 4000 Da remained deformed. Note coagulation of embryos (darkening, intransparency) exposed to PEGs \geq 8000 Da. From Pelka et al. (2017).

Figure 28: Changes in the area of the chorion of zebrafish (*Danio rerio*) embryos exposed to polyethylene glycols (PEGs) of different molecular weight



Changes in the area of micrographs of the chorion of zebrafish (*Danio rerio*) embryos exposed to polyethylene glycols (PEGs) of different sizes (molecular weight) in relation to the areas at the onset of exposure. The exposure covers the time-period from the onset of exposure (approx. 5 min) to the time point of no further shrinkage of the chorion (approx. 24 h). For PEG 3000, additional measurements were made after 36 and 48 h. Note that maximum shrinkage is already reached after approx. 7 h for most PEGs. Data are given as percent change over negative controls at the beginning of the exposure for three independent runs with 20 embryos (n = 3). Coagulated eggs at PEGs 8000 and 12,000 were excluded from evaluation. From Pelka et al. (2017).

1.7 Conclusions: Barrier function of the zebrafish chorion

From multiple experiments with both fluorescent markers and differently sized polyethylene glycols, the following conclusions can be drawn:

- With appropriate modifications of the FET protocol, embryos can reproducibly be dechorionated at 24 h post-fertilization (hpf) with reliably high survival rates of ≥ 90 %. Dechorionation of younger stages (< 24 hpf) is generally possible, however with lower survival rates.</p>
- By means of fluorescent marker molecules, the molecular size for the passage of noncharged chemical substances across the chorion of the zebrafish could be set at approx. 3000 Da.
- More in-depth studies with differently sized polyethylene glycols allowed narrowing down the limit for free passage across the chorion of the zebrafish between 3000 and 4000 Da. Depending on time, polyethylene glycols with a molecular weight ≤ 3000 Da show a more or less free passage.
- The passage of most uncharged industrial chemicals is not impaired by a systematic size limitation.
- Dechorionation of zebrafish eggs eliminates problems associated with the passage of the zebrafish chorion by high molecular weight substances.
- The passage of charged molecules across the chorion of zebrafish is substantially lower than that of non-charged molecules. Given the complex interaction of three-dimensional

molecular configuration and molecular charge, a precise prediction of the penetration barrier for charged molecules is not possible without a complex model based on a comprehensive database (which are not yet available).

- Dimethyl sulfoxide (DMSO) concentrations of 0.1 % and 1 % decrease the barrier function of the chorion, whereas a concentration of 0.01 % DMSO remains ineffective. As a consequence, DMSO may be used without complications as a solvent, however, only at a maximum concentration of 0.01 % (0.1 ml/L; 100 mg/L), which is in line with the recommendations of the OECD GD 23 on difficult substances (OECD, 2018).
- Since dimethyl sulfoxide (DMSO) does not impact zebrafish embryo survival up to concentrations well above 2 %, DMSO may be used as a detergent to facilitate the passage of substances that might be critical (1) in terms of their molecular size to pass the zebrafish chorion (2) for handling in aqueous solutions due to their high lipophilicity. So far, there is no convincing evidence that after appropriate solution by the aid of solvents highly lipophilic substances are not taken up by zebrafish embryos.
- For highly lipophilic substances, neither the Acute Fish Toxicity test (AFT, OCED TG 203) nor the Fish Embryo Acute Toxicity test (FET, OECD TG 236) appear to be suitable protocols, since highly lipophilic substances might accumulate over extended periods of time and might not have reached a steady-state after an exposure time restricted to 96 h. Rather, for highly lipophilic substances, longer-term tests such as a Fish Early Life-Stage test according to OECD TG 210 (OECD, 2013b) should be conducted (Oris et al., 2012).

2 Biotransformation and bioactivation capacities in early life stages of zebrafish (*Danio rerio*) – a literature review

2.1 Introduction

Major objections to the use of the FET have mainly been based on the assumption of limited biotransformation capacities in early life stages of fish. Repeatedly, doubts have been expressed concerning the comparability of metabolic capacity between early juvenile and adult life stages of zebrafish (Busquet et al., 2008a; Chng, 2013; Verbueken et al., 2017). In fact, a limited biotransformation or bioactivation capacity could lead to underestimations of toxic or teratogenic potency of a xenobiotic, which would be bioactivated in other life stages (Busquet et al., 2008a; Kluver et al., 2014). On the other hand, given the *ex utero* development, early-life stages of (zebra)fish depend on their intrinsic metabolic capacity for elimination or transformation of xenobiotics (Verbueken et al., 2017); therefore, it seems reasonable to assume at least qualitatively appropriate biotransformation capacities.

As a consequence, in order to strengthen the acceptance of the FET, it seemed essential to attempt to better understand the metabolic capacities of zebrafish embryos across different lifestages. However, while literature concerning spatial and temporal expression patterns of xenobiotic metabolizing enzymes has been growing fast at the genome and RNA levels, only few investigations addressed protein function and catalytic activities, although these are ecologically much more relevant parameters than DNA- and RNA-based characteristics. The database concerning zebrafish metabolism, especially in early-life stages, is much less complete than that available for other model organisms (e.g. rainbow trout) and is scattered over more than 100 studies (Spitsbergen and Kent, 2003).

The following chapter has been designed to provide a state-of-the-art summary aimed at facilitating the identification of remaining gaps in our knowledge on biotransformation in zebrafish in general and zebrafish embryos in specific. The literature study has been initiated as a starting-point for prioritization of further research into biotransformation in zebrafish embryos.

2.2 Metabolism of xenobiotics

The majority of xenobiotics (compounds foreign to the organism) undergo metabolic transformation (biotransformation) in order to be excreted (more) effectively (Donato and Castell, 2003). In vertebrates, xenobiotic metabolism is primarily located in the liver; however, extrahepatic tissues such as kidney, intestine and gills may also make major contributions to xenobiotic metabolism (Buhler and Williams, 1988; Chambers and Yarbrough, 1976).

As a rule, metabolism of xenobiotics is a biphasic process subdivided into phase I and II reactions. Phase I reactions introduce (*de novo* synthesis) or unmask polar groups (e.g. -OH, - COOH, -SH, -NH₂) in xenobiotics. These oxidative, reductive, or hydrolytic processes provide functional sites for subsequent conjugation reactions (phase II reactions). The key enzymes for phase I reactions are, among other less important enzyme systems, various isoforms of cytochrome P450-dependent monooxygenases (CYPs). CYP-mediated reactions primarily include hydroxylation and epoxidation of aromatic and aliphatic compounds, ester and ether cleavage, as well as heteroatom dealkylation.

In phase II reactions, functional groups are conjugated with polar endogenous compounds. Phase II reactions include glucuronidation, glutathione and amino acid conjugation as well as acetylation; all of these reactions significantly increase the water solubility of the generated metabolites, allowing and facilitating their excretion *via* the bile (intestinal route) and/or as urine *via* the kidney route (Buhler and Williams, 1988; Donato and Castell, 2003; Parkinson and Ogilvie, 2001; Schlenk et al., 2008).

2.3 The role of metabolism in activation and detoxification of xenobiotics

Metabolism is well recognized as a critical factor influencing the teratogenic and toxic potency of numerous xenobiotics. In the majority of cases, metabolism leads to detoxification (bioinactivation); however, toxification (bioactivation) of xenobiotics has also been reported for multiple substances (Goldstone et al., 2010; Nebert and Dalton, 2006). For instance, it is assumed that only about one quarter of all carcinogenic xenobiotics are tumorigenic in their non-metabolized form, whereas the remaining xenobiotics require CYP-mediated activation before they can develop full toxicity or teratogenicity (Nebert and Dalton, 2006). Proteratogens and protoxicants can thus be activated enzymatically into electrophilic and reactivate intermediates such as electrophiles, epoxides and free radicals, which readily interact with cellular components such as proteins, DNA or lipids (Blaschke et al., 2010; Miller, 1970; Wells et al., 2004). Typical model xenobiotics that are bioactivated via metabolism include benzo[a]pyrene, allyl alcohol, aflatoxin, cyclophosphamide and carbamazepine (Kluver et al., 2014; Weigt et al., 2011).

Most importantly, toxicity and teratogenicity of metabolically activated substances can be underestimated, if biotransformation capacities are limited (Busquet et al., 2008a; Kluver et al., 2014). In case of bioinactivation, limited biotransformation capacities might lead to a falsenegative result. A typical example could be albendazole, which has been reported to be metabolically inactivated in adult zebrafish, but not in embryonic zebrafish (Boix et al., 2015; Carlsson et al., 2013; Mattsson et al., 2012). Supplementation by a mammalian metabolization system resulted in a compensation of the lack of stage-dependent deactivation, i.e. a decrease of developmental toxicity in embryonic zebrafish (Mattsson et al., 2012).

2.4 Cytochrome P450-dependent monooxygenases (CYPs)

It is well established that cytochrome P450-dependent monooxygenases (CYPs) catalyze the oxidative metabolism (phase I metabolism) of many xenobiotics, and approximately 75 % of all xenobiotics are estimated to undergo CYP mediated phase I biotransformation including hydroxylation, dealkylation and/or epoxidation (Guengerich, 2007; Nebert and Dalton, 2006). In general, for zebrafish, much is known about spatial and temporal expression patterns (Otte et al., 2010), but catalytic activities, substrate specificity, inducibility and stereoselectivity of CYPs have only rarely been examined. In general terms, in vertebrates, the number of CYP genes ranges from 40 to > 100, with CYPs distributed among 19 gene families (Nelson et al. 2013).

2.5 Number of CYP isoforms in zebrafish

So far, a total of 86 - 96 CYP genes, distributed among 17 - 18 categories of CYP families, have been described for zebrafish (Goldstone et al., 2010; Saad et al., 2016a; Stegeman et al., 2010). CYP families 1, 2, 3 and – to a smaller extent – CYP family 4 are considered to mainly metabolize xenobiotics, while the CYP families 5 - 51 primarily contribute to the metabolism of endogenous substrates (Goldstone et al., 2010). The full number of CYP genes and their family (> 40 % amino acid identity) and subfamily (> 55 % amino acid identity) memberships have already been well documented (Goldstone et al., 2010; Saad et al., 2016a). The following paragraph will review available data of the CYP families 1 - 4 in zebrafish.

In recent years, the focus has been put on genes in CYP family 1, especially on CYP1A as a bioindicator for detecting dioxin-like compounds, polychlorinated biphenyls and polyaromatic hydrocarbons (Jonsson et al., 2009; Kais et al., 2017; Noury et al., 2006). Far less attention has been given to the CYP families 2 to 4.

2.6 Expression patterns of CYPs during zebrafish development

In recent years, most studies have focused on expression patterns of CYPs. They have determined the temporal and spatial expression of CYPs in unchallenged (control) zebrafish by using quantitative real-time PCR (qPCR), microarray techniques, transgenic zebrafish lines, and whole-mount *in situ* hybridization (WISH) technologies. The diversity of methodological approaches, methodology-dependent detection limits and test intervals makes a direct comparison between studies difficult and sometimes leads to conflicting results; as an example, data for CYP3A65 gene expression trends in early-life stages of zebrafish are given in Fig. 29. Fig. 30 summarizes trends in temporal expression patterns for all CYP450 isoforms that could be located in the literature for embryonic and juvenile life-stages zebrafish. Table 1 gives a comprehensive account of spatial and temporal patterns of enzyme expression in zebrafish.





Comparison of CYP3A65 gene expression trends in early-life stages of zebrafish (*Danio rerio*). Data were generated by (1) whole transcriptome microarray (Goldstone et al. 2010), (2) whole-mount in situ hybridization (Tseng et al. 2005), and (3) the use of transgenic zebrafish lines (Chang et al. 2013). From Lörracher & Braunbeck (2019)

CYP Embryonic zebrafish Juvenile zebrafish Methodology Reference Unfertilized eggs Significant ARNT2-nomalized expression Quantitative real-time PCR Goldstone et al. 201 Significant ARNT2-nomalized expression Single-color Agilent custom whole- new whole- method method read-time PCR Goldstone et al. 201 VPLA Method logy Reference Method logy Color Agilent custom whole- method method read-time PCR Goldstone et al. 201 Method logy Method logy Color Agilent custom whole- method method read-time PCR Goldstone et al. 201 Method logy Method logy Custom target read-time PCR Goldstone et al. 201 Method logy Method logy Custom target read-time PCR Goldstone et al. 2014 Method logy Method logy Custom target read-time PCR Goldstone et al. 2014 Method logy Method logy Custom target read-time PCR Goldstone et al. 2014 Method logy Method logy Custom target read-time PCR Goldstone et al. 2014 Method logy Method logy Method logy Color logy Goldstone et al. 2014 Method logy Method logy Method logy Method logy Goldstone et al. 2014 Method logy Method logy Method logy Method logy Goldstone et al. 2014					
Unterlitized eggs: Duratification Conditione et al. 201 Significant ARN12-normalized expression Single-color Aglient custom whole- are arise ari	сүр	Embryonic zebrafish	Juvenile zebrafish	Methodology	Reference
CYPIA CY		Unfertilized eggs: Significant ARNT2-normalized expression		Quantitative real-time PCR	Goldstone et al. 2010
CYP1A CUANTIATIVE real-time PCR CUANTIATIVE real		There are a here and a here at here	97 4 2	Single-color Agilent custom whole- transcriptome microarray	Goldstone et al. 2010
CYP1A Cumitative real-time PCR Glisic et al. 2014 Cumitative real-time PCR Glisic et al. 2014 Cumitative real-time PCR Glisic et al. 2010 Cumitative real-time PCR Jones et al. 2010 Cumitative real-time PCR Briauning et al. 2015 Cumitative real-time Briauning et al.		ohpt 24 hpt 48 hpt 72 hpt 96 hpt 120 h	pr 144 hpr 192 hpr 73 hpr 67 dpr	Quantitative real-time PCR	Goldstone et al. 2009
Image: Control of the state	СҮР1А	the and and Zhat what to he		Quantitative real-time PCR	Glisic et al. 2014
Object 24 hpt 25 hpt 36 hpt 27 hpt Bräuning et al. 2015 Object 24 hpt 72 hpt 30 hpt 140 hpt 140 hpt 140 hpt 21 dpt 2007		00pt 24 hpt 22 hpt 45 hpt 12 hpt 120 hpt		Quantitative real-time PCR	Jones et al. 2010
Object 24 hpt 48 hpt 120 hpt 21 dpt 21 dpt </td <td></td> <td>0hpr 24.hpr 48.hpr 72.hpr 98.hpr 120.hpr</td> <td></td> <td>Quantitative real-time PCR</td> <td>Bräuning et al. 2015</td>		0hpr 24.hpr 48.hpr 72.hpr 98.hpr 120.hpr		Quantitative real-time PCR	Bräuning et al. 2015
		0hpt 24 hpt 48 hpt 72 hpt 96 hpt 120 hpt	144 hpt 188 hpt // 14 dpt 21 dpt 28 dpt // 27 dpt	Quantitative real-time PCR	Jönsson et al. 2007

Figure 30:	Temporal expression of	CYP450 in embryonic and	juvenile zebrafish ((Danio rerio)
------------	------------------------	-------------------------	----------------------	---------------



СҮР	Embryonic zebrafish	Juvenile zebrafish	Methodology Single-color Agilent custom whole-	Reference Goldstone et al. 2010
	ange one same and and		transcriptome microarray	
	48 hpt 72 hpt 96 hpt 120 hpt	14 hpr 188 hpr // 14 dpr 21 dpr 28 dpr // 57 dpr	Quantitative real-time PCR	Bräuning et al. 2015
CYP1C2	0hpt 24 hpt 48 hpt 72 hpt 96 hpt 120 hpt		Quantitative real-time PCR	Jönsson et al. 2007
	80 hpf Expression tends to be higher than the expression of CYP1A or CYP1B1 (Statistically not significant)		Quantitative real-time PCR	Jõnsson et al. 2007
			Quantitative real-time PCR	Goldstone et al. 2010
	Short 6 hort 12 hort 24 hort 36 hort 48 hor		Single-color Agilent custom whole- transcriptome microarray	
			Quantitative real-time PCR	Goldstone et al. 2009
	Ohpt 24 hpt 48 hpt 72 hpt 96 hpt 120 hpt	144 hpf 168 hpf 1 192 hpf 1 57 dpf		
СУРОЛИ			Whole mount in situ hybridization	Wang et al. 2007
	 + 36 hpf		Quantitative real-time PCR	Wang et al. 2007
	abort 6 hor 12 hor 24 hor 36 hor 48 hor		Single-color Agilent custom whole-transcriptome microarray	Goldstone et al. 2010
CYP2K6	0hpr 24 hpr 48 hpr 22 hpr 86 hpr 120 hpr	14 hpt	Quantitative real-time PCR	Wang-Buhler et al. 2005

СҮР	Embryonic zebrafish	Juvenile zebrafish	Methodology	Reference
CYP2K8	Not that take what what		Single-color Agilent custom whole-transcriptome microarray	Goldstone et al. 2010
CYP2K16	New Cherr 22 hpr 24 hpr 24 hpr		Single-color Agilent custom whole-transcriptome microarray	Goldstone et al. 2010
CYP2K17	aber the 24 ber 24 ber 24 ber		Single-color Agilent custom whole-transcriptome microarray	Goldstone et al. 2010
CYP2K18	Abr 6 hpt 22 hpt 24 hpt 26 hpt 49 hpt		Single-color Agilent custom whole- transcriptome microarray	Goldstone et al. 2010
	0 1 0 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1		Whole mount in situ hybridization Transgenic line Tg(cyp2k18:egfp)	Poon et al. 2017
CYP2K19	Ref that 21 hor 24 hor 44 hor 44 hor		Single-color Agilent custom whole- transcriptome microarray	Goldstone et al. 2010
CYP2K20	abort at hor 24 hor 24 hor 24 hor 24 hor		Single-color Agilent custom whole- transcriptome microarray	Goldstone et al. 2010
CYP2K21	jatu an and a set an set and a set a		Single-color Agilent custom whole- transcriptome microarray	Goldstone et al. 2010
CYP2K22 CYP2K7	adata atalaa aatalaa aatalaa		Single-color Agilent custom whole- transcriptome microarray	Goldstone et al. 2010
CYP2K31	April 6 hpr 23 hpr 24 hpr 24 hpr 24 hpr		Single-color Agilent custom whole- transcriptome microarray	Goldstone et al. 2010

СҮР	Embryonic zebrafish	Juvenile zebrafish	Methodology	Reference
67100A0	April Objet 12 hpc Abort Abort		Single-color Agilent custom whole- transcriptome microarray	Goldstone et al. 2010
CITZNIS	00hpf 24 hpf 48 hpf 72 hpf 98 hpf 120 hpf		Whole mount in situ hybridization Transgenic line Tg(cyp2n13: egfp)	Poon et al. 2017
CYP2P6	Apple 6 hpd 12 hpd 24 hpd 24 hpd 46 hpd		Single-color Agilent custom whole- transcriptome microarray	Goldstone et al. 2010
CYP2P7	Shert ohjet 12 hert 24 hert 36 hert 46 her		Single-color Agilent custom whole- transcriptome microarray	Goldstone et al. 2010
СҮР2Р9	abort 6 hort 22 hort 24 hort 36 hort 41 hort		Single-color Agilent custom whole- transcriptome microarray	Goldstone et al. 2010
CYP2P10	add the log to the log		Single-color Agilent custom whole- transcriptome microarray	Goldstone et al. 2010
CYP2P14	and other and and and		Single-color Agilent custom whole- transcriptome microarray	Goldstone et al. 2010
	Sheet a hear so hear as hear as hear as hear		Single-color Agilent custom whole- transcriptome microarray	Goldstone et al. 2010
CYP2R1	obpf 3 hpt 6 hpt 12 hpt 24 hpt 72 hpt	// // // // // // // // //	Quantitative real-time PCR	Peng et al. 2017
CYP2U1	aber cher 23 her 24 her 26 her 48 her		Single-color Agilent custom whole- transcriptome microarray	Goldstone et al. 2010

СҮР	Embryonic zebrafish	Juvenile zebrafish	Methodology	Reference
	Unfertilized eggs Significant ARNT2-normalized expression		Quantitative real-time PCR	Goldstone et al. 2010
CYP2V1 CYP2J26	Abor 6 hpd 12 hpd 24 hpd 26 hpd 46 hpd		Single-color Agilent custom whole- transcriptome microarray	Goldstone et al. 2010
	there are a set of the		Quantitative real-time PCR	Jones et al. 2010
CYP2X7	ator elar schor ather ather		Single-color Agilent custom whole- transcriptome microarray	Goldstone et al. 2010
CYP2X8	Aper 6 hpt 28 hpt 26 hpt 49 hpt		Single-color Agilent custom whole- transcriptome microarray	Goldstone et al. 2010
CYP2X10	alor short as hort as hort		Single-color Agilent custom whole- transcriptome microarray	Goldstone et al. 2010
СҮР2ҮЗ	idea idea advise advise advise advise		Single-color Agilent custom whole- transcriptome microarray	Goldstone et al. 2010
CYP2Y4	alor that still alor		Single-color Agilent custom whole- transcriptome microarray	Goldstone et al. 2010
CYP2AA1	age that 24 by 26 by 46 by		Single-color Agilent custom whole- transcriptome microarray	Goldstone et al. 2010
CYP2AA2	age of the 24 hpr 24 hpr 24 hpr		Single-color Agilent custom whole- transcriptome microarray	Goldstone et al. 2010
CYP2AA3	alor that 2 hor 2 hor 2 hor 2 hor		Single-color Agilent custom whole- transcriptome microarray	Goldstone et al. 2010
	unfertilized eggsunfertilized eggs significant ARNT2-normalized expression		Quantitative real-time PCR	Goldstone et al. 2010
CYP2AA4	and the second		Single-color Agilent custom whole- transcriptome microarray	Goldstone et al. 2010

СҮР	Embryonic zebrafish	Juvenile zebrafish	Methodology	Reference
CYP2AA7	April 12 hpr 24 hpr 24 hpr 44 hpr		Single-color Agilent custom whole- transcriptome microarray	Goldstone et al. 2010
CYP2AA8	abor the ather ather ather		Single-color Agilent custom whole- transcriptome microarray	Goldstone et al. 2010
CYP2AA9	and the advect advect		Single-color Agilent custom whole- transcriptome microarray	Goldstone et al. 2010
CYP2AA11	about a hor at hor at hor at hor		Single-color Agilent custom whole- transcriptome microarray	Goldstone et al. 2010
CYP2AA12	Apple the athen athen athen athen		Single-color Agilent custom whole- transcriptome microarray	Goldstone et al. 2010
CYP2AD2	ater cher strier artist artist		Single-color Agilent custom whole- transcriptome microarray	Goldstone et al. 2010
CYP2AD3	Sheet a high at high at high at high		Single-color Agilent custom whole- transcriptome microarray	Goldstone et al. 2010
CYP2AD6	and the title allow allow allow		Single-color Agilent custom whole- transcriptome microarray	Goldstone et al. 2010
CYP2AE1	Shert of her 23 her 26 her 26 her 26 her		Single-color Agilent custom whole- transcriptome microarray	Goldstone et al. 2010



СҮР	Embryonic zebrafish	Juvenile zebrafish	Methodology	Reference
CVP3C2	0 0 0 1 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0		Quantitative real-time PCR	Shaya et al. 2014
	abort the athor athor athor		Single-color Agilent custom whole- transcriptome microarray	Goldstone et al. 2010
rvb3r3	the short take a hor		Quantitative real-time PCR	Shaya et al. 2014
	abpr chor 12 hpr 24 hpr 24 hpr		Single-color Agilent custom whole- transcriptome microarray	Goldstone et al. 2010
CYP3C4	chort 1 hpd 5 hgd 12 hgd 28 hgd 48 hgd		Quantitative real-time PCR	Shaya et al. 2014
CYP4F43	aber a her 22 her 24 her 26 her 46 her		Single-color Agilent custom whole- transcriptome microarray	Goldstone et al. 2010
CYP4V7	and the active active active active		Single-color Agilent custom whole- transcriptome microarray	Goldstone et al. 2010
CYP4V8	and the state as her as her as her		Single-color Agilent custom whole- transcriptome microarray	Goldstone et al. 2010
CYP4T8	abort object 24 hor 26 hor 48 hor		Single-color Agilent custom whole- transcriptome microarray	Goldstone et al. 2010

Comparative trends within the examined CYP450 isoforms (Fig. 30):



Trends in temporal expression patterns for all CYP450 isoforms that could be located in the literature for embryonic and juvenile life-stages zebrafish (*Danio rerio*). From Lörracher & Braunbeck (2019)

References		Otto at al 2010	Scomaienchi et al. 2010			Bräuning et al. 2015	Otte et al. 2017				Otte et al. 2010. Scomaienchi et al. 2010		Saad at al. 2018			Pauka et al. 2011	Jönsson et al. 2009
Adult												Much higher EROD activity than	developmental stages.	No gender differences.			Constitutive and significantly inducible EROD activity
												Liver microsomes					Whole gill arch and liver
Juvenile	Intestine, liver, around the anal pore, circulatory system, head kidney, nephric duct, kidney	Same spatial distribution as embryo (104 hpf).									Activity approximately 50 - 30% of the level at 104 hpf					Constitutive and significantly inducible EROD activity in the supernatant of body homogenates	
	128 hpf										128 hpf					2 wpf	
Embryo	Cytoplasm of cells of the envelope layer, yolk syncytial layer, developing germ layers	Different parts of the head, straight tube of the heart, dorsal aorta, myotome, envelope of the yolk, pronephric duct, urogenital pore	Circulatory system, heart, inner party of the eye, mesencephalon, fourth ventriole, liver primordium, primordia of the pronephrio kidney, pronephrio duct, urogenital pore	All vessels, heart, intestine, pancreas, gut, liver, head kidney, pronephrio duct	Intestine, liver, around the anal / urinary pore, circulatory system, head kidney, nephric duct, kidney	Significant increase in activity	Activity above the level of detection, but lower than the limit of quantification.	Decrease in activity. Activity below the limit of detection	Increase in activity. Activity above the limit of quantification	Activity above the limit of quantification	Activity varies over time Peak in expression at 8 hpf and 104 hpf	Highest activity with a large inter-batch variation	Negligible activity (significantly lower than at 48 hpf)	Increase in activity, even further at the end of organogenesis	Negligible activity		
	8 hpf	32 hpf	56 hpf	80hpf	104 hpf	24-120 hpf	2,5 hpf	48 hpf	96 hpf	120hpf	8 - 120 hpf	5 hpf	24 -48 hpf	72 - 96 hpf	120 hpf		
СҮР						Mammals:	CYP1 mediated	Zebrafish: CYP1A > CYP1C2 > CYP1B1 = CYP1C1 >	CYP1D1 (Heterologous expressed CYP1s)								
Activity assay			In vivo- EROD assay Substrate: 7-Ethoxyresorufin								In vitro- EROD assay Substrate: 7-Ethoxyresorufin						

The chorion structure and biotransformation capacities of zebrafish as boundary conditions for OECD Test Guideline 236
(Acute Fish Embryo Toxicity Test)

Table 1:Spatial and temporal patterns of CYP enzyme expression in zebrafish (Danio rerio)

Activity assay	СҮР		Embryo	Juvenile		Adult	References
In vitro- BROD assay	Human: CYP1A, CYP2B, CYP3A	2.5 -86 hpf	No activity above the limit of detection				Otto, 64 41, 2017
Substrate: 7-0-Benzylresonufin	Zebrafish: CYP1A = CYP1B1 > CYP1Cs > CYP1D1	120 hpf	Activity above the limit of detection but under the limit of quantification				
In vitro-PROD assay Substrate: 7-Pentoxyresondin	Zebrafish: CYP1A, CYP1Cs > CYP1D	2.5 - 120 hpf	No activity above the limit of detection				Otte et al. 2017
In vivo-ECOD assay Substrate: 7-Ethoxycoumarin		96 hpf	Constitutive ECOD activity Incubation time dependence in activity				Jones et al. 2010
In vitro- ECOD assay	Mammals: CYP2 mediated				Liver microsomes	Constitutive ECOD activity ECOD activity does not differ significantly from the activities of Cyprinus carpio , Oryzias latipes and Poecilia reticulata	Funari et al. 1997
Substrate: 7-Ethoxycoumarin				Relative constitutive ECOD activity: liver > gill > muscle > brain significantly inducible			Wu et al. 2014
In vivo-OOMR assay Substrate: Octyloxymethylresonufin	Mammals: CYP3 mediated	96 hpf	Incubation time dependence in activity significantly inducible				Jones et al. 2010
In vivo-Iuminescence-based Promega P450-Glo™ CYP3A4 assay Substrate: Luciferin-IPA Substrate: Luciferin-IPA	Human:	120 hpf	Activity well distinguished from background noise				Chng 2013
In vitro-Luminescence-based Promega P450-Glo™ CYP3A4 assay Substrate: Luciferin-IPA	CTPSAA				Liver microsomes	Metabolite concentration below the lower limit of quantification	Verbueken et al. 2017
<i>In vivo</i> -luminescence-based Promega P450-Glo™ CYP3A4 assay Substrate: Luciferin-BE	Human: CYP3A4	48 hpf 72 hpf	Constitutive activity Significant modification by inducers and inhibitors Constitutive activity higher than at 48 hpf				Li et al. 2011

Activ	ity assay	СҮР		Embryo	Juvenile		Adult	References
In vivo-BFCOD	Aesse	Human: CYP3A4 >>CYP2B8 Zebrafish: Minimal contribution	96 hpf	Activity consistently measured Significant modification by inducers and inhibitors				Creusot et al. 2014, Rerwick et al. 2008
Substrate: 7-Bei trifluoromethylox	nzyloxy-4- sumarin	of CYP1A to the BFCOD activity CYP1A> CYP1C2>CYP1C1, CYP181 (Heterologous	120 hpf	Constitutive and significantly inducible BFCOD activity				Oziolor et al. 2017
In vitro-BOMR Substrate: Benz methylresorufin	assay yloxy-	Human: CYP3A	5 - 120 hpf Microsomes of whole embryo homogenates	BOMR activity only observed at 72 hpf and 96 hpf Activity close to the limit of quantification		Liver microsomes	Activity significantly higher than in microsomes of whole embryo homogenates	Verbueken et al. 2017
Abbreviatio	ns:				*			
ECOD		Ethoxycoumarin-O	h-deethylase					
EROD		Ethoxyresorufin-O.	-deethylase					
BFCOD		7-Benzyloxy-4-trifl	uoromethylcou	imarin-O-debenzyloxylase				
BOMR BROD		Benzyloxy-methyl- 7-benzylresorufin-(resorufin O- debenzylast	<i>0</i> 1				
00MR PROD		Octyloxymethylres 7-pentoxvresorufin	torufin O-depentvlas	ũ				
		and an and a second a	the second se					

Spatial and temporal patterns of CYP enzyme expression in zebrafish (*Danio rerio*). From Lörracher & Braunbeck (2019)

Most CYP genes (51 out of 52 CYP genes investigated for zebrafish) are expressed as mRNA at least once during the embryonic development until < 120 hours post-fertilization (hpf) (Braunig et al., 2015; Chang et al., 2013; Corley-Smith et al., 2006; Glisic et al., 2016; Goldstone et al., 2010; Jones et al., 2010; Peng et al., 2017; Poon et al., 2017a; Shaya et al., 2014; Tseng et al., 2005; Wang-Buhler et al., 2005; Wang et al., 2007). The only exception, so far, is CYP3C4, which, however, has only been investigated from 0 to 48 hpf (Shaya et al., 2014). For most CYPs, expression throughout the early development fluctuates in a wave-like fashion (Fig. 29) peaking at different time points (Goldstone et al., 2010).

For CYP1A, CYP2V1, and CYP2AA4, maternal contribution to mRNA transcript abundance was detected in unfertilized oocytes (Goldstone et al., 2010). CYP1B1 (24 hpf) and CYP1D1 (9 hpf) show their highest expression in early stages of embryonic development (Goldstone et al., 2009; Jonsson et al., 2007a). Other CYPs such as CYP1A, CYP2J26, CYP2K6, and CYP2R1 show an increase in expression after 48 - 72 hpf, which is the period, when zebrafish embryos hatch (Jones et al., 2010; Peng et al., 2017; Wang-Buhler et al., 2005). In the case of CYP1C1, CYP1C2, and CYP1D1, there is a decrease in mRNA transcript abundance towards the end of embryonic development (96 - 120 hpf) (Jonsson et al., 2007a).

Information concerning temporal CYP expression patterns in juvenile life-stages (eleuthero¬embryos and larvae) of zebrafish (between 120 hpf and 3 months post-fertilization; mpf) are available for the following eight isoforms: CYP1A, CYP1B1, CYP1C1, CYP1C2, CYP1D1, CYP2K6, CYP2R1, and CYP3C1. However, as holds for embryos, the abundance of CYP mRNA transcripts in juvenile life-stages is still fluctuating. CYP1A, CYP1C1, CYP2K6, and CYP2R1 have higher expression levels in juvenile zebrafish than in embryos. For the isoforms CYP1B1, CYP1D1, and CYP3C1, the levels of expression in embryonic life-stages is higher than or equal to the level in juvenile life-stages (Braunig et al., 2015; Corley-Smith et al., 2006; Goldstone et al., 2009; Jonsson et al., 2007a; Peng et al., 2017; Wang-Buhler et al., 2005).

Adult		Juvenile	Embryonic
CYP1A	CYP2R1	CYP1A	СҮР2К18
CYP1B1	CYP2Y3	СҮР2К6	СҮР2К7
CYP1C1	CYP2AA1	СҮР2К7	CYP2N13
CYP1C2	CYP2AA2	CYP3A65	СҮРЗА65
CYP1D1	CYP3A65		
CYP2J1	CYP3C1		
СҮР2К6	CYP3C2		
СҮР2К7	СҮРЗСЗ		
CYP2K18	CYP3C4		

 Table 2:
 CYP isoforms in adult, juvenile, and embryonic life-stages of zebrafish (Danio rerio)

CYP isoforms investigated concerning their organ-specific spatial expression patterns in adult, juvenile, and embryonic life-stages of zebrafish (*Danio rerio*). From Lörracher & Braunbeck (2019)

2.7 Spatial expression patterns of CYPs in zebrafish

The overall metabolic capacity is determined by the isoforms and abundance of CYPs expressed in the respective organs. In fish, the liver is considered as the main organ responsible for biotransformation of xenobiotics (Pesonen and Andersson, 1991). CYPs have been described as mainly liver-localized enzymes, but are also expressed in extrahepatic tissues and organs including kidneys, intestine, brain, eyes, heart, muscles, and gills. Using qPCR, antibody staining, transgenic zebrafish lines, and whole mount in situ hybridization, spatial expression patterns of 18 CYPs have been described (Table 2) (Corley-Smith et al., 2006; Fetter et al., 2015; Goldstone et al., 2009; Jonsson et al., 2007b; Kubota et al., 2013; Peng et al., 2017; Poon et al., 2017a; Shaya et al., 2014; Siegenthaler et al., 2017; Taylor, 2005; Tseng et al., 2005; Wang-Buhler et al., 2005; Wang et al., 2007; Yang et al., 2017).

Only for CYP3C1 and CYP3A65, data are available on spatial expression patterns covering all developmental stages. CYP3C1 shows pronounced organ and developmental differences in spatial expression (Table 3). At 12 hpf, CYP3C1 mRNA transcripts are widely distributed through the whole embryo. In later embryonic development (48 and 120 hpf), CYP3C1 mRNA is restricted to the brain and the gastrointestinal tract (Corley-Smith et al., 2006). In juvenile life-stages CYP3C1 is expressed in many organs and tissues including intestine, liver, kidneys, gills, and eyes (Taylor, 2005). In adult zebrafish, expression of CYP3C1 is distributed over multiple organs, with clear sex-differences in the spatial distribution profile (Corley-Smith et al., 2006; Shaya et al., 2014). For a detailed account of spatial expression patterns in embryonic, juvenile and adult zebrafish, see Tables 4 and 5.

Knowledge of spatial and temporal CYP expression is important for a deeper understanding of metabolism, but does not provide a direct insight into functional metabolic activity itself (Sadler et al., 2016), since there is no direct correlation between CYP expression levels, amounts of corresponding proteins, and measurable CYP activities (Bluhm et al., 2014; Goldstone et al., 2010). The CYP activity, for example, can be influenced by changes in protein stability or the efficiency of translation (Bluhm et al., 2014).

Age	and sex	Brain	Eye	Gills	Intestine	Liver	Kidney	Heart	Muscle	Testis	Ovary
	12 hpf				++ Widely d	listributed ac	ross the entir	e embryo +	+		
Embryo	48 hpf	‡									
	120 hpf	ŧ			ŧ						
Juvenile	21 dpf		+ + +	++++	ŧ	‡	++				
Adult	Female		+	+	ŧ	ŧ		+			ŧ
	Male	+	++++	+	+	+++	‡	+		‡	

Table 3: Spatial expression of CYP3C1 in developmental stages of zebrafish (Danio rerio)

The spatial expression profile of CYP3C1 in different developmental stages of zebrafish. White blocks denote non-existence of data, grey blocks denote non-detectable expression, light green low expression, intermediate green blocks denote moderate expression, and dark green blocks high expression (in comparison within the given CYP isoform and life-stage; data from Corley-Smith et al., 2006; Shaya et al., 2014; Taylor, 2005) . From Lörracher & Braunbeck (2019)

	References	Taylor 2005	Taylor 2005	Poon et al. 2017		Fetter et al. 2015	Taylor 2005		Poon et al. 2017	
	Applied method	Antibody staining	Antibody staining	Transgenic line TG (CYP2K18: egfp)	Whole mount in situ hybridization	Whole mount in situ hybridization	Antibody staining	Transgenic line (CYP2N13:egfp)	Transgenic line TgFOSMID (CYP2N13:egfp)	Whole mount in situ hybridization
	Others	Pseugobranch Epitrelium, Skin of the head Oropharymx Taste bud Esophagus	Skin Oropharymx Esophagus Muscle tunic of the intestine				Oropharymx Esophagus Skin of the head Taste bud Cartilage of the head Skin of the trunk and tail		Olfactory bulb Cloaca Skin	
		• •	o •				•		0	
	Muscle	×	×	×	×	×	s. L	×	×	×
	Heart	O Ventricle	0/0	×	×	×	л.s.	×	0	×
istribution	Kidney Pronephros	Mesonephric duct	 Renal tubules 	×	×	0	0	×	×	×
	Liver	n.s.	×	×	0	×	×	×	0	0
	Intestinal tract	•	×	0	0	×	•	O Intestinal bulb region	0	0
	Gill	n.s.	×	×	×	×	•	×	×	×
	Eye	×	Lens O	×	×	×	0	×	×	×
	Brain	×	×	×	×	×	n.s.	×	×	×
	Age	21 dpf	21 dpf	120 hpf	120 hpf	96 hpf	21 dpf	120 hpf	120hpf	120 hpf
	сүр	CYP1A	CYP2K6	CYP2K18			CYP2K22 CYP2K7		CYP2N13	

Table 4: Spatial expression patterns of CYP 450 in zebrafish (Danio rerio)

							Distribution						
СҮР	Age	Brain	Eye	Gill	Intestinal tract	Liver	Kidney Pronephros	Heart	Muscle		Others	Applied method	References
	72 hpf	×	×	×	×	0	×	×	×			Whole mount in situ hybridization	
	84 hpf	×	×	×	 Especially in the foregut 	0	×	×	×			Whole mount in situ hybridization	
CYP3A65	96 hpf	×	×	×	 Especially in the foregut 	0	×	×	×			Whole mount in situ hybridization	Tseng et al. 2005
	120 hpf	×	×	×	•	•	×	×	×			Whole mount in situ hybridization	
	21 dpf	×	Comea	×	0	×	O Renal tubules	0	0	0	Skin of the trunk Cartilage	Antibody staining	Taylor 2005
	12 hpf				Wi	dely distribute	ed thorugh the who	ole embryo				Whole mount in situ hybridization	
	48 hpf	•	×	×	×	×	×	×	×			Whole mount in situ hybridization	Corley-Smith et al. 2006
CYP3C1	120 hpf	•	×	×	0	×	×	×	×	•	Pharynx	Whole mount in situ hybridization	
							•			0	Skin of the tail and the trunk		
	21 dpf	n.s.	•	•	0	0/0	Mesonephric duct renal tubes	n.s.	n.s.	0	Taste bud, skin of the head	Antibody staining	Taylor 2005
										•	Oropharynx		
Abbreviation	:51		5	not studied									

65

studied with negative results

×

moderate expression high expression

00.

< 120 hpf < 3 mpf

embryo juvenile

low expression

Comparison within the given CYP isoform:

From Lörracher & Braunbeck (2019)

The chorion structure and biotransformation capacities of zebrafish as boundary conditions for OECD Test Guideline 236 (Acute Fish Embryo Toxicity Test)

Table 5: Spatial expression patterns of cytochrome P450 in adult zebrafish (Danio rerio)

							Distrib	ution						
CYP	Sex	Brain	Eye	Gill	Intestine	Liver	Kidney	Heart	Muscle	Testis	Ovary	Others	Applied method	References
	\$1\$	•	0	•	•	•	0	•	n.s.	n.s.	n.s.		Quantitative real-time PCR	Goldstone et al. 2008 Cytochrome P450 1D1
	No significant sex differences	0	•	•	•	•	•	•	n.s.	0	0		Quantitative real-time PCR	Jönsson et al. 2007
CYP1B1	No significant sex differences	•	•	•	0	0	•	•	n.s.	0	0		Quantitative real-time PCR	Jönsson et al. 2007
CYP1C1	No significant sex differences	•	•	•	0	0	0	•	n.s.	0	0		Quantitative real-time PCR	Jönsson et al. 2007
CYP1C2	No significant sex differences	•	•	0	0	0	•	•	n.s.	0	0		Quantitative real-time PCR	Jönsson et al. 2007
CYP1D1	\$15	•	0	•	•	•	0	•	n.s.	n.s.	n.s.		Quantitative real-time PCR	Goldstone et al. 2008
	\$15	•	n.s.	n.s.	×	•	•	•	0	0	۰		Quantitative real-time PCR	Wang et al. 2007
011201	\$15	•	n.s.	n.s.	×	×	×	×	n.s.	•	•		Whole mount in situ hybridization	Wang et al. 2007
CYP2K6	\$1\$	×	×	×	×	•	n.s.	×	×	×	•		Quantitative real-time PCR	Wang-Buhler et al. 2005
CYP2K7 CYP2K22	\$1\$	n.s.	n.s.	n.s.	n.s.	•	n.s.	n.s.	n.s.	0	0		Quantitative real-time PCR	Siegentahler et al. 2017
CYP2K18	64	n.s.	n.s.	n.s.	n.s.	•	n.s.	n.s.	n.s.	n.s.	n.s.		Quantitative real-time PCR	Yang et al. 2017
CYP2R1	\$1\$	×	×	×	×	•	×	×	•	×	0	 Visceral adipose tissue 	Quantitative real-time PCR	Peng et al. 2017
CYP2Y3	64	n.s.	n.s.	n.s.	n.s.	•	n.s.	n.s.	n.s.	n.s.	n.s.		Quantitative real-time PCR	Yang et al. 2017
	4	0	0	n.s.	•	0	0	0	n.s.		0		Quantitative real-time PCR	2002 la ta atorio
	۴0	0	ο	n.s.	•	•	•	•	n.s.	•			Quantitative real-time PCR	

Spatial expression patterns of cytochrome P450 in adult zebrafish (Danio rerio)

							Distrib	ution						
сүр	Sex	Brain	Eye	Gill	Intestine	Liver	Kidney	Heart	Muscle	Testis	Ovary	Others	Applied method	References
	0+	•	•	n.s.	•	•	•	0	п.s.		0		Quantitative real-time PCR	
CIPZAAZ	50	•	•	n.s.	•	•	•	0	n.s.	0			Quantitative real-time PCR	- Kubota et al. 2013
CYP3A65	\$15	0	0	•	•	•	n.s.	•	n.s.	n.s.	n.s.		Quantitative real-time PCR	Tseng et al. 2005
	0+	×	0	0	•	•	n.s.	0	n.s.		•	O Skin	Quantitative real-time PCR	Corley-Smith et al 2005
CYP3C1	0+	•	0	0	•	0	0	•	n.s.		•	O Offactory rosette Spieen		Shaya et al. 2014
-	50	0	•	0	0	•	•	0	n.s.	•		Olfactory rosette Spleen	Quantitative real-time PCR	Shaya et al. 2014
CVD3C3	0+	•	•	•	•	•	0	•	n.s.		•	 Olfactory rosette Spleen 		Cherry of Mild
CTP3CZ	50	0	0	0	•	•	0	0	n.s.	•		 Officatory rosette Spleen 	quantrative real-time PCK	ଧୀଙ୍ଗର ୧୩.୧୬.୮୬
CYP3C3	0+	•	0	0	•	•	•	•			•	 Olfactory rosette Spleen 	Quantitative real-time PCR	Shaya et al. 2014
	Q	0	0	0	•	•	0	•		•		O Olfactory rosette Spleen		
- Jenno	¢+	•	•	•	0	0	•	•			•	O Offactory rosette Spleen		
CTP3C4	Ş	•	•	•	•	•	•	•			•	O Offactory rosette Spieen		oraya et al. 2014
Abbreviation	ŝ													
			n.s. ×	not studied studied wit	t h negative rest	Ħ								
Comparison with	hin the given CYP450 is	soform:	00.	weak expre moderate e strong expr	ession sypression ession									

Spatial expression patterns of CYP 450 in embryonic and juvenile zebrafish (*Danio rerio*). From Lörracher & Braunbeck (2019)

2.8 CYP-dependent activities

For the understanding of metabolic capacity, it is important to measure CYP-dependent activity. Various preparations have been used to investigate CYP-dependent activities of zebrafish, including whole body, liver microsomes, S9 samples, heterologously expressed CYPs, as well as primary hepatocytes. Most studies have determined CYP-dependent activities using fluorescent-based catalytic activity assays (Braunig et al., 2015; Chng, 2013; Creusot et al., 2015; Jones et al., 2010; Jonsson et al., 2009; Otte et al., 2017; Otte et al., 2010; Pauka et al., 2011; Saad et al., 2016b; Scornaienchi et al., 2010; Verbueken et al., 2017; Wu et al., 2014). The problem with fluorescent-based assays is that their interpretation is complicated due to overlapping and still unknown substrate specificities of CYP isoforms in zebrafish (Scornaienchi et al., 2010). CYP1s have been most extensively studied of all the CYPs. By heterologous expression of CYP1A, CYP1B1, CYP1C1, CYP1C2, and CYP1D1 (e.g. over-expression of zebrafish CYP genes in E. coli or yeast), catalytic activities and specificities towards 11 fluorometric substrates (e.g. 7-ethoxyresorufin, 7-methoxyresorufin, 7-benzyloxyresorufin) have been demonstrated in zebrafish (Scornaienchi et al., 2010).



Figure 31: In vivo localization of EROD activity in 8 h old zebrafish (Danio rerio) embryos

In vivo localization of EROD activity at 8 hpf zebrafish (*Danio rerio*) embryos exposed to β-naphthoflavone (A, B) or artificial water only (C, D) imaged by CLSM at 8 hpf. Lateral views on the embryos (A, C) showed EROD activity in the cytoplasm of the envelope layer and of the layers below (a). EROD activity also appears in the yolk syncytial layer (b). Nuclei of the envelope layer are marked with arrowheads (A, C) and yolk granules are indicated (c). Sections at the median plane (B, D) indicated EROD activity in the developing germ layers (marked by arrowheads). Scale bar: 200µm (insets: 100µm); animal pole top. From Otte et al. (2010)

Commonly, ethoxyresorufin-O-deethylase (EROD) activity is used as marker for CYP1 activity. All CYP1s, especially CYP1A, metabolize 7-ethoxyresorufin into the highly fluorescent metabolite 7-hydroxyresorufin (Scornaienchi et al., 2010; Stegeman et al., 2015). In zebrafish, basal CYP1A activity has been detected from 2.5 hpf onwards (Otte et al., 2017). Because of the zebrafish embryo's transparency, CYP1A activity can be measured in vivo. By using confocal laser scanning microscopy, Otte et al. (2010) could demonstrate CYP1 activity as early as 8 hpf (Fig. 31).

Only recently, Kais et al. (2017, 2018) developed the technique by Otte et al. (2010) further and optimized it to a rapid screening test (Fig. 32). Kais et al. (2017, 2018) documented the applicability of in vivo EROD-imaging for a selection of model compounds (Fig. 33 - 37). For more details, see Kais et al. (2017).

Deethylation of 7-ethoxycoumarin (ECOD) is an established marker for CYP2 activities in mammals. Activity towards 7-ethoxycoumarin was detected at 96 hpf (Jones et al., 2010). Earlier life-stages of zebrafish have not been investigated, yet.

For human CYP3-dependent activities, several substrates are available (e.g. luciferin-derivatives, 7-benzyloxy-4-trifluoromethylcoumarin, and benzyloxy-methylresorufin). CYP3-dependent activities in zebrafish have been detected from 48 hpf onwards, but in most cases starting at 72 hpf (Chng, 2013; Creusot et al., 2015; Jones et al., 2010; Li et al., 2011b; Oziolor et al., 2017; Renwick et al., 2008; Verbueken et al., 2017).





EROD induction patterns in zebrafish (*Danio rerio*) embryos after 72 h (A, D), 96 (B, E) and 120 h (C, F) of exposure to 10 μ g/l β -naphthoflavone (maximum EROD induction) following 3 h (A - C) and continuous (D - F) exposure. There is no difference in signal intensity between 3 h and continuous exposure, except for the 120 h exposure, which gives a slightly lower signal following continuous exposure (F). The areas of fluorescence in the embryos exposed for 3 h appear smaller than those seen after continuous exposure. Number of experiments: n = 12 with 4 - 6 embryos each. Epifluorescence microscopy (exposure time: 40 ms). From Kais et al. (2017)





In vivo localization of EROD activities in zebrafish (*Danio rerio*) embryos after 72 (A, D, G, J), 96 (B, E, H, K) and 120 h (C, F, I, L) exposure to 10 μ g/L β -naphthoflavone (A - C), solvent (0.1 % DMSO; D-F), 0.6 mg/L (EC₁₀) chlorpyrifos (G - I) and 0.6 mg/L chlorpyrifos + 10 μ g/L β -naphthoflavone (J - L). In all developmental stages, embryos exposed to β -naphthoflavone show a bright signal in the liver, whereas embryos of the solvent control show a weaker signal reflecting constitutive activity. The signal intensities after single chlorpyrifos exposure as well as after co-exposure to chlorpyrifos and β -naphthoflavone show lower signal intensities than in the solvent control after both 96 and 120 h. The signal in the co-exposure approach after 72 h is as strong as that of the solvent control. Epifluorescence microscopy, exposure times: 60 ms (72 h), 30 ms (96 h), 40 ms (120 h). From Kais et al. (2018)





In vivo localization of EROD activities in zebrafish (*Danio rerio*) embryos after 72 (A, D, G, J,M), 96 (B, E, H, K, N) and 120 h (C, F, I, L, O) exposure to 10 μ g/L β -naphthoflavone (A - C), solvent (0.1 % DMSO; D - F), 0.25 (G - I), 0.5 (J - L) and 1.0 mg/L chlorpyrifos (M - O)+ 10 μ g/L β -naphthoflavone each. Embryos exposed to β -naphthoflavone show a bright signal in the liver, whereas the embryos of the solvent control show a weaker signal reflecting constitutive activity. Particularly after 96 and 120 h, signal intensities after co-exposure to 0.5 and 1.0mg/L chlorpyrifos and β -naphthoflavone were lower than in the solvent, whereas 0.25mg/L chlorpyrifos did not inhibit EROD activity completely (G - I). Epifluorescence microscopy, exposure times: 40 ms (72 h), 100 ms (96 h), 100 ms (120 h). From Kais et al. (2018).





In vivo localization of EROD activities in zebrafish (*Danio rerio*) embryos after 72 (A, D, G, J), 96 (B, E, H, K) and 120 h (C, F, I, L) exposure to 10 μ g/L β -naphthoflavone (A - C), solvent (0.1 % DMSO; D - F) and 0.6 mg/L chlorpyrifos following 3 h (G - I) and continuous (cont.) exposure (J - L). In all developmental stages, embryos of short-term exposure to β -naphthoflavone show bright signals in the liver (A - C), whereas embryos of the solvent control show weaker signals (D - F) and demonstrate no differences to continuous exposure. Signals after short-term exposure to chlorpyrifos also show no differences, if compared to continuous exposure, except for the short-term exposure of 72 h embryos (G), which show stronger fluorescence than after continuous exposure (J) and are similar to the signal of the solvent control. Epifluorescence microscopy, exposure times: 60 ms (72 h), 30 ms (96 h), 40 ms (120 h). From: Kais et al. (2018)
2.9 Inducibility of cytochrome P450s in zebrafish

Various CYPs such as CYP1A, CYP1B1, CYP1C1/C2 and CYP3A65 can be induced or inhibited by xenobiotics (Braunig et al., 2015; Cunha et al., 2016; Glisic et al., 2016; Kubota et al., 2013; Tseng et al., 2005). In many cases, the induced CYP is at the same time capable of metabolizing the inducing agent (or other xenobiotics) (Zhu, 2010). Induction of CYP activities, especially in the liver, can therefore lead to accelerated xenobiotic metabolism (Celander, 2011).

In zebrafish, xenobiotic-mediated induction of CYP1s is well documented as, e.g., expression, protein abundance, and 7-ethoxyresorufin-*O*-deethylase activity. In zebrafish, CYP1A, CYP1B1, CYP1C1, and CYP1C2 are induced by aryl hydrocarbon receptor (AhR) agonists such as polycyclic aromatic hydrocarbons (Braunig et al., 2015; Jonsson et al., 2007b). In contrast, CYP1D1 was found not to be inducible by AhR agonists such as PCB126 and TCDD (Goldstone et al., 2009). In contrast, exposure to caffeine, ibuprofen, or carbamazepine (0.05 - 5 μ M) leads to a significant down-regulation of CYP1A (Aguirre-Martinez et al., 2017). For the CYP families 2, 3, and 4, mechanism of down- and upregulation, inductors, and inhibitors, are less well understood.

Starting from 8 hpf onwards, induction of CYP1s has been shown to occur in all developmental stages of zebrafish embryos. However, tissue distribution and maximum inducibility vary substantially between developmental stages (Aguirre-Martinez et al., 2017; Cunha et al., 2016; Glisic et al., 2016; Jones et al., 2010; Kubota et al., 2015; Liu et al., 2016; Otte et al., 2010; Zhang et al., 2015).

When assessing metabolization capacities based on CYP expression and activity, respectively, potentially adverse effects by the xenobiotic itself should be considered whenever possible.

2.10 Bioactivation capacity of early-life stages: functional confirmation of CYP activity

It is assumed that only about one quarter of all carcinogenic xenobiotics are tumorigenic in their non-metabolized form; the remaining 75 % xenobiotics require CYP-mediated metabolic activation before elucidating their full carcinogenicity, toxicity, or teratogenicity (Fantel, 1982; Nebert and Dalton, 2006). In fact, a limited biotransformation capacity can lead to an underestimation of protoxicants or proteratogens (Busquet et al., 2008b; Kluver et al., 2014).

So far, the only protoxicant known to be less toxic to zebrafish early-life stages than to the adult and juvenile zebrafish is allyl alcohol (Kluver et al., 2014). In mammals, allyl alcohol is metabolized *via* oxidation into the highly reactive metabolite acrolein. In zebrafish embryos, the reduced toxicity of allyl alcohol is caused by a lack of alcohol dehydrogenase 8a (adh8a), which is responsible for the oxidative formation of acrolein (Kluver et al., 2014).

Concerning proteratogens, there is prominent evidence that zebrafish embryos possess significant biotransformation capacities. Weigt et al. (2011) demonstrated that ten well-known proteratogens were teratogenic in zebrafish embryos exposed for 3 days (Table 6). All ten substances tested (2-acetylaminoflurene, benzo[a]pyrene, aflatoxin B1, carbamazepine, phenytoin trimethadione, cyclophosphamide, ifosfamide, tegafur, and thio-TEPA), are known to undergo CYP-mediated bioactivation in mammals. The result suggests that zebrafish early-life stages (< 72 hpf) are capable of testing proteratogenic substances, even without addition of an exogenous metabolic activation system (e.g. S9 mix, microsomes; Weigt et al., 2011).

Substance	LC ₅₀	EC ₅₀	TI (LC ₅₀ /EC ₅₀)
2-Acetylaminofluorene	6.9 μM	-	< 1
Benzo[a]pyrene	5.1 μΜ	0.52 μΜ	9.81
Aflatoxin B1	2.3 μΜ	2.2 μΜ	1.05
Carbamazepine	> 500 µM	222 μΜ	> 1
Phenytoin	> 250 μM	386 μM	> 1
Trimethadione	45.7 mM	23.5 mM	1.95
Cyclophosphamide	8.4 mM	4.7 mM	1.79
Ifosfamide	3.2 mM	3.1 mM	1.03
Tegafur	30.3 mM	3.4 mM	8.91
Thio-TEPA	306 µM	53.2 μM	5.76

Table 6:LC50, EC50 and teratogenicity index (TI) of selected proteratogens in 3 d old zebrafish
(Danio rerio) embryos

LC₅₀, EC₅₀ and teratogenicity index (TI) of 2-acetylaminoflurene, benzo[a]pyrene, aflatoxin B1, carbamazepine, phenytoin trimethadione, cyclophosphamide, ifosfamide, tegafur, and thio-TEPA d old zebrafish (*Danio rerio*) embryos. From Weigt et al. (2011)

2.11 Metabolism capacity – chemical analysis

Advances in chemical analytical methods greatly contributed to the understanding of metabolic activities (Guengerich, 2000). Highly sensitive methods (e.g. liquid chromatography tandemmass spectrometry) allow the detection and measurement of low abundance metabolites in body homogenates and water. Depletion of substances or formation of metabolites allow conclusions to be drawn about metabolic processes and activities of enzymes (Li et al., 2011b).

In zebrafish, studies have been primarily focusing on metabolism of human drugs such as paracetamol, benzocaine, bupropion, verapamil, and phenacetin (Alderton et al., 2010; Brox et al., 2016b; Chng et al., 2012; Jones et al., 2010; Kantae et al., 2016; Saad et al., 2017). So far, several phase-I- and phase-II-related biotransformation products have been described. Based on mass analysis and comparison to standards, the molecular structure of occurring transformation products can be estimated (Alderton et al., 2010).

Studies comparing xenobiotic metabolism of embryonic, juvenile, and adult life-stages are rare. Since there are different analytical approaches using different methods as well as targets (e.g. liver microsomes, primary hepatocytes, whole body homogenates) comparability is not given among the available studies. No investigation yet has systematically studied xenobiotic metabolism within all developmental life-stages of zebrafish. Especially for juvenile zebrafish life-stages, there is hardly any information available (Table 7).

Occurrence of seven transformation products after three hours exposure to clofibric acid, depletion of benzocaine, and increase of its phase II metabolites (4-aminobenzoic acid, 4-acetaminobenzoic acid) indicate early (< 24 hpf) presence of metabolic activity in zebrafish (Brox et al., 2016a; Brox et al., 2016b). In most cases, such as dextromethorphan, diclofenac, and phenacetin, the maximum number and highest level of metabolites appear post hatch (Alderton et al., 2010; Brox et al., 2016b; Chng et al., 2012; Saad et al., 2017). For testosterone, e.g., metabolism capacity of zebrafish early-life stages (< 120 hpf) appears rather limited (Table 8).

	References		Weigt et al. 2011		Chng et al. 2012, Weigt et al. 2010, Kantae et al. 2016, Jones 2010		Weigt et al. 2011, Troxel et al. 1997	Carlsson et al. 2011.	Carisson et al. 2013. Mattsson et al. 2012	Poon et al. 2017	Weigt et al. 2011,	Hsu et al. 1996 Amanuma et al. 2002,
	s identified	Phase II					*					
¢,	Metabolite	Phase I			>		*			*		
Adult (> 3.5 m	Biotransformation		n.s.	Liver microsomes - N-acety-p-tenzoquinone imine (NAPQI)			- Aflatoxicol Aflatoxico-glucuronide - Unidentified polar metabolite - Aflatoxin-epoxide		n.s.	 Hydroxy-amiodarone Mono-N-desethylamiodarone Hydroxy-mono-N- desethylamiodarone 	Bioactivation of proteratogen: - Formation B[a]P-DNA adducts	
	s identified	Phase II							>			
(Juli	Metabolites	Phase I										
Juvenile (> 6.	Biotransformation		n.s.		જુર		n.s.	144 hpf - Albendazole suffoxide - Albendazole suffone - Albendazole-2-aminosulfone	Albendazole was reduced to a high extend at lower concetrations. At higher concertration the reduce in concentration was not observed or much less evident.	n.s.		n.s.
	s identified	Phase II			>				:			
(Jul	Metabolite	Phase I	:				:					:
Embryo (< 5 c	Biotransformation		72 hpf Bioactivation of proteratogen: - teratogenic effects	25 MgC Quantification of: - Paraetamol-sulfate - Paraetamol-glucuronide Level of metabolites below or dose to the limit of quantification	72 hpf Bouchristion of proteratiogen: - No teratogenio effecta after exposure (2 - 3 hpf)) (MAS) leads to activation system (MAS) leads to significant increase of affected embryos	<u>86 hpf</u> - Acetaminophen-sultate - Acetaminophen-olutathione	<u>12 hpf.</u> Bioactivation of proteratogen: - teratogenic effects	2-34 hpf High spontanous transformation to albendazole sulfoxid after 10 h (even in control).	it is not shure if emoryos contribute to comersation	n.s.	24-40 hpf Bioactivation of proteratogen: - increased mutant frequency in the rpsL tarrost gene	22 hef 22 hef Bioactivation of proteratiogen: teratogenic effects
	Substance		-Acetylaminofluorene		Letaminophen Baracetamol		Aflatoxin B,		libendazole	Amiodarone		3enzo[a]pyrene

Embr	yo (< 5 c	(Juli		Juvenile (> 6	đpf)		Adult (> 3.5 mp	6		
Biotransformation	Metabolites identifie	s identified		Biotransformation	Metabolites	s identified	Biotransformation	Metabolite	es identified	References
Phase I Phase	Phase I Phase	Phase	=		Phase I	Phase II		Phase I	Phase II	
2.34 hpf 4-aminobentacio acid 4-acetaminobentacio acid 4-acetaminobentacio: acid	>	>		n.s.			ъ.s.			Brox et al. 2016
88 hpt - Two mono-gluouronidated metabolites - Mono-suffaked metabolite - Di-suffaked metabolite - Gluruonidated and sulftaked metabolite	*	*		n.s.			Two mono-glucuronidated metabolite Nono-suidated metabolite On-suidated metabolite - Di-suidated + suiffated metabolite - Di-glucuronidated e metabolite		×	Fol et al. 2017
n.s.				n.s.			 Demethylated metabolite Reduced metabolite Glucuronidated metabolite Sulfated metabolite 	*	*	Li et al. 2014
<u>08 hpf</u> - Mono-glucuronidated metabolite - Mono-suffated metabolite	*	>		n.s.			 Mono-glucuronidated metabolite Mono-sultated metabolite 		>	Fol et al. 2017
n.s.				<u>168 hpf</u> - Hydraxybupropian	*		n.s.			Alderton et al. 2010
72-90 hpf 7 ometabolities could be measured continously 7 metabolities name. All of them showed staady increase during the experimental period. - Mono-subtaned metabolite - Mono-subtaned metabolite - Mono-subtaned metabolite - Di-conjugated: glucorsylated and - Di-conjugated: glucorsylated and - Di-conjugated: glucorsylated and - Di-conjugated: glucorsylated and suftaed metabolite - Di-conjugated: glucorsylated and suftaed metabolite - Mono-hydroxylated metabolite	*	*		u.s.			S.			Hu et al. 2012
12 h <u>of</u> Bioactivation of proteratogen: - teratogenio effects	:			n.s.			Π.S.			Weigt et al. 2011
72.hpf No metabolite detected				<u>168 hpf</u> - Cisapride N-sulfate		*	n.s.			Alderton et al. 2010
7 <u>hef</u> - 7 Transformation products 18 hef - 14 Transformation products - 18 Transformation products	*	×		n.s.			y y			Brox et al. 2016
n.s.				n.s.			 Demethylated metabolite Reduced metabolite Methylated matbolite 	*		Li et al. 2014

	References		Weigt et al. 2011, Busquet et al. 2008	Saad et al. 2017, Alderton et al. 2010	Alderton et al. 2010, Saad et al. 2017	Li et al. 2014	Carlsson et al. 2013	Carlsson et al. 2013	Jones et al. 2009, Jones et al. 2012	Weigt et al. 2011	Li et al. 2014	Alderton et al. 2010,
	s identified	Phase II									*	
ĥ	Metabolite	Phase I		×	×	>					*	
Adult (> 3.5 m)	Biotransformation		n.s.	Liver microsomes - 3-Methorymorphinan - Dectrophan	Liver microsomes: - 4 hydroxy-diclofenac - 5 hydroxy-diclofenac (Distinction not possible)	 Demethylated metabolite Methylated matbolite 	n.s.	n.s.	n.s.	n.s	 Demethylated metabolite Methylated matbolite Methylated matbolite Glucuronidated metabolite Sulfated metabolite 	n.s.
	s identified	Phase II										:
(jdp	Metabolites	Phase I		×			>	>	>			:
Juvenile (> 6	Biotransformation		Ъ.S.	108.bpf - O-desmethyl dextromethorphan - N-desmethorybhan - On the methoroyphan hydroxylated metabolite - Dextrorphan	103 hpr - Hydroxy-diclofenac	n.s.	144 h <u>pf</u> - Fenbendazole - Oxfendazole	144 hpf - Oxfendazole		S L	n.s.	168 hpf - More polar, earlier eluting metabolites (HPLC)
	s identified	Phase II										
(Jd	Metabolite	Phase I	:	×	>				>	-		
Embryo (< 5 c	Biotransformation		48 her Bioactivation of proteratogen: no teratogenic effects after incubation (2 - 3 hpf) 22 her Bioactivation of innitizationen:	 - teralogenic effects - Mincle embryor microsomess. 5 hpt. 48 hpt. 5 hpt. 48 hpt. - 3 Methoxymorphinan destrophan :below the limit of quantification - Destrophinan Destrophinan - Destrophinan: unquantifiable - Destrophan - Destrophan 	Whole embryon prima: unquantance Whole embryon microsomes. - No metabolite detected <u>26 hof</u> - Hydroxylated metabolite (close to the lower finant of detection) - Level of hydroxylated metabolites 10 finanes of	adults. n.s.	n.s.	n.s.	<u>90 hpf</u> - Hydroxyibuprofen	72 hpf Bioactivation of proteratogen: - teratogenic effects	n.s.	n.s.
	Substance		Cyclophosphamide	Dextromethorphan	Dictofenac	Epiberberine	Febantel	Fenbendazole	lbuprofen	lfosfamide	Jatrorrhizine	Lauric acid

	References		derton et al. 2010 ing et al. 2012.	aad et al. 2017, oon et al. 2017	xon et al. 2017	et al. 2014		ox et al. 2016	eigt et al. 2011	derton et al. 2010	eigt et al. 2011		denton et al. 2010, hng et al. 2012, kad et al. 2017	eigt et al. 2011	eigt et al. 2011	ang et al. 2016
	s identified	Phase II	<u> </u>	>	ă	>		ä	M	R.	M		₹0 <i>8</i>	×	M	×
Ģ	Metabolite	Phase I		>	>	8							>			*
Adult (> 3.5 mp	Biotransformation		low level of - 1'-Hydroxymidazołam - 4- Hydroxymidazołam - N-Glucuronidation metabolites	Liver microsomes - No depletion of mid.azolam - Low, non-quantifiable oncentrations of 1-hydroxy- mid.azolam	- Hydroxy-nefazodone	 Demethylated metabolites Hydroxylated metabolites Glucuronidated metabolite Sulfated metabolite 		u.s.	n.s.	n.s	n.s.	Liver microsomes In total 8 hydroxylated metabolites with several icromers	 - By Hydroxytestosterone - 108Hydroxytestosterone - 20-Hydroxytestosterone - Other metabolites not identified. 	n.s.	n.s.	 Diphenyiphosphate Mondynorcylated diphenyiphosphate Mondynorcylated triphenyiphosphate Dihydroxylated triphenyiphosphate Dihydroxylated triphenyiphosphate Glouromot odd conjugated metabolites Sludy of 10 additional metabolites were detectable or below
	s identified	Phase II											>			
(Jab	Metabolite	Phase I		>						>			>			
Juvenile (> 6	Biotransformation		168 hpf A total of 5 metabolites - Mono-oxydated metabolite - Oxydated metabolite	- retenting are metabolite	n.s.	n.s		n.s.	n.s.	168 hpf - Hydroxy-tacrine	s n	<u>168 hpf</u> - Testosterone glunuronide conjugate - One hydroxylated metabolite		n.s.	n.s.	ns.
	s identified	Phase II						>								
(Juli	Metabolite	Phase I		>				*	:		-		>			
Embryo (< 5	Biotransformation		5-120 hpf (microsomes) - No depletion of midazolam 72 hpf Eiro meterbriter detected		n.s.	ų š	<24 hpt - Paracetamol	24-96 hpf - Paracetamol - Paracetamol glucuronide - Paracetamol suffate	<u>72 hpf</u> Bioactivation of proteratogen: - teratogenic effects	.s.п	22 hpt. Bioactivation of proteratogen: - teratogenic effects	Whole embryo microsomes. 5 hpt - 120hpf - No substrate consumption	12.0 hof 11.0 hof In total 2 hydroxytestosterone metabolites - Main metabolite, was unique, and not 66-Hydroxytestosterone - testosetrone glucuronide	72 hpf. Bioactivation of proteratogen: - teratogenic effects	72 hpf. Bioactivation of proteratogen: - teratogenic effects	ц.
	Substance			Midazolam (MDZ)	Nefazodone	Palmatine		Phenacetin	Phenytoin	Tacrine	Tegafur		Testosterone	Thio-TEPA	Trimethadione	Triphenyl phosphate (TPHP)



Biotransformation metabolites identified in embryonic, juvenile, and adult stage zebrafish. From Lörracher and Braunbeck (20199

	0		
Metabolites	Embryo (microsomes)	Juvenile	Adult (microsomes)
Phase I	5-120 hpf: no testosterone consumption 120 hpf: two hydroxylated metabolites: (1) main metabolite, unique, and not found in microsomes of adult zebrafish, (2) 6β-hydroxytestosterone	168 hpf: hydroxylated metabolite	Six hydroxylated metabolites with several isomers: 6β-hydroxytestosterone 16β-hydroxytestosterone 2α-hydroxytestosterone Three not identified metabolites
Phase II	Testosterone glucuronide conjugate	Testosterone glucuronide conjugate	Not studied

Table 8:Testosterone metabolite profiles zebrafish (Danio rerio) at various developmental
stages

Testosterone metabolite profiles of embryo, juvenile, and adult zebrafish (*Danio rerio*) reveal differences in the metabolism capacity. Data from Alderton et al. (2010) and Saad et al. (2017), compiled by Lörracher and Braunbeck (2019)

2.12 Conclusions and recommendations for future research into the biotransformation capacities of zebrafish (*Danio rerio*)

From the considerations above, the following conclusions can be drawn:

- So far, zebrafish embryos, juveniles and adults have been studied to a very different extent with respect to their biotransformation capacities. Since especially juveniles have been neglected largely, transfer of biotransformation data from juvenile fish to adult fish does not appear justified. Overall, our knowledge about xenobiotic transformation capacities in zebrafish can – at best – be called fragmentary.
- However, whenever studied in more detail, biotransformation in zebrafish embryos could be documented. So far, only for rare exceptions such as allyl alcohol and albendazol, a lack of biotransformation (allyl alcohol to acrolein) could be demonstrated due to a lack of the enzyme required (alcohol dehydrogenase in the case of allyl alcohol).
- Studies on biotransformation gene expression (induction) is not as conclusive as studies in enzyme biosynthesis (transcriptomics, proteomics) or – even better – biochemical activity of enzymes. Since our knowledge on the spectrum of substances accepted as substrates by various cytochrome P450 isoforms is little, there is no way to extrapolate gene activation to biochemical functionality.
- ► There is no adequate database to extrapolate observations from mammalian studies to fish and fish embryos. For many P450 isoforms, we do not have conclusive evidence of congruence in cytochrome P450 terminology between mammalian studies and studies in

lower vertebrates. Therefore, transfer of a link between nomenclature and metabolic function from mammals to fish or vice versa ("same nomenclature means same function") is not necessarily justified.

- With respect to biotransformation phase II reactions, our knowledge is even more fragmentary than for phase I metabolization.
- In case there is suspicion from studies with mammalian models that metabolic activation or inactivation by biotransformation might play a role for the toxicity profile of a given substance, additional experiments with external biotransformation via, e.g., mammalian S9 preparations may help to compensate for a potential lack of biotransformation in zebrafish.
- As a gold standard for studies into biotransformation capacities in (zebra)fish embryos, juveniles and adults, gene expression studies need to be linked to transcriptome / proteome analysis data as well as chemical analytical analyses into the metabolites formed by cytochrome P450 enzymatic activity. Such studies, however, are most challenging in terms of expertise, resources, time, number of animals used and funding.
- Given the limitations of resources and the multitude of potential substrates for cytochrome P450 conversion, studies providing indirect evidence of the existence of active biotransformation seem more appropriate for screening purposes. Such indirect evidence may come from
 - studies designed to visualize the formation of fluorescent or colored metabolites (at best, in living embryos: in situ live imaging); unfortunately, such studies so far suffer from an apparent lack of appropriate fluorescent metabolites;
 - studies based on chemical analyses of emerging metabolites; however, in many instances, protocols for the chemical identification of metabolites still need to be developed;
 - studies based on the detection of biological effects by substances activated by cytochrome P450 action.

3 Analysis of the relevance and adequateness of the Fish Acute Toxicity Test (AFT) according to OECD TG 203 to fulfil the information requirements and addressing concerns under REACH

3.1 Summary

The aim of this study was a re-evaluation of historical AFT data produced according to OECD TG 203. The selection of compounds used for the present study was identical to that used by (Scholz et al., 2016) for the analysis of the relevance and adequateness of using the Fish Embryo Acute Toxicity Test (FET; OECD TG 236) to fulfill the information requirements and to address concerns under REACH. The data were retrieved from the ECHA database in July 2017 and were analyzed for the relevance and adequateness of using the AFT to fulfill the information requirements and addressing concerns under REACH. The selection criteria used in the present study to obtain the AFT database are identical to those that were used for the ECHA FET study (Scholz et al., 2016; Sobanska et al., 2018).

Starting with a dataset of 2936 studies of a total of 1842 substances, in a first step (I), all studies with LC_{50} entries not precisely defined (e.g. LC_{50} data given as range only) were removed. In the second step (II), duplicate studies were eliminated. These two initial steps reduced the number of studies to 56.4 % of all studies and 54.3 % of the test substances originally contained in the dataset. Further filtering steps included: (III) studies with LC_{50} values above half the water solubility; (IV) studies into substances with log $P_{0W} > 4$, when test concentrations were not verified by chemical analytics; and (V) inorganic substances. Application of filtering steps 3 to 5 resulted in a remainder of 38 % of studies and of 38 % of the substances of the original dataset.

An intra- and interspecies comparison of the remaining data revealed robust values for AFT data: The analysis illustrated that almost all studies of intra- and interspecies comparison (93.45 % and 96.22 %) lie within a deviation range between 1 and 100, which is currently accepted within the REACH process. A comparison of AFT data from the present study to the data from the corresponding FET study (Scholz et al., 2016; Sobanska et al., 2018) showed some comparability, although not all filtering steps were identical in both studies. Namely, the filtering steps "Keep studies with predicted pH at saturation >5 or <9" and "Remove studies with no toxicity and maximum test concentration < 10 fold above baseline toxicity" were not performed in the analysis of AFT studies due to the tools being unavailable. In the end, when only zebrafish (Danio rerio) studies are considered (as was done in the ECHA FET study), the potential for elimination of data from the database was similar for both FET and AFT data. However, the rationale for only using zebrafish data is debatable, because data from other fish species is used for regulatory purposes and also correlates well with FET data (Belanger et al., 2013). Since these findings indicate that the robustness of FET and AFT data are similar, the suitability of the FET for REACH purposes should be reconsidered, and political action(s) should be taken to make FET data acceptable for application in REACH.

With regard to the use of OECD 203 for regulatory purposes, more attention should be paid to the physicochemical properties of the test substances and to the correct preparation and use of test substance concentrations. Additionally, chemical verification of test concentrations should be mandatory for substances with log $P_{OW} > 4$ or tested above water solubility for a better reliability of the data.

3.2 Introduction

The ECHA database represents a valuable tool to quickly obtain information on chemical components. Apart from substance description and physical and chemical properties, it provides information on the environmental fate and pathways as well as ecotoxicological and toxicological information. Given the rapid pace of scientific developments and the significant increase of information within the REACH process, the ECHA database undergoes constant modification of data and is updated frequently.

The aim of the present study was a retrospective re-evaluation of a dataset for the short-term Acute Fish Toxicity Test (AFT) according to OECD TG 203, which had been retrieved from the ECHA database (https://www.echemportal.org/echemportal/propertysearch/) in July 2017, with respect to its suitability for the information requirements and addressing concerns under REACH. Furthermore, the study has been designed as an analysis in parallel to the already existing analysis of FET data conducted by Scholz et al. (2016).

3.3 Treatment of datasets

The dataset was extracted in July 2017 and has been worked with since then, being aware that some data especially on significant ecotoxicological studies in the REACH process may have been changing in the meantime. Therefore, this analysis should be regarded as a snapshot of the AFT data available in the ECHA database from July 2017.

In collaboration with Dr. Stefan Scholz and Dr. Nils Klüver (UfZ Leipzig), the ECHA database was screened for historical AFT data. All studies performed with the major species used for ecotoxicological testing, rainbow trout (*Onchorhynchus mykiss*), carp (*Cyprino carpio*), fathead minnow (*Pimephales promelas*), zebrafish (*Danio rerio*), Japanese medaka (*Oryzias latipes*), bluegill sunfish (*Lepomis macrochirus*) and guppy (*Poecilia reticulata*) were incorporated.

Search criterion	Keyword
Study result type	Experimental result
Reliability (Klimisch score)	1, 2
Test guideline	OECD Guideline 203
Test guideline, qualifier	According to, equivalent or similar to
GLP compliance	yes
Test type	Static, semi-static, flow-through
Test type; Water media type	all
Total exposure	96 h
Effect concentrations, Endpoint	LC ₅₀
Effect concentrations, Effect conc.	overlapping 0 – 10,000 g/L
Effect concentrations, Basis for effect	all

 Table 9:
 Overview of the search criteria in the ECHA database for historical AFT data

Overview of the search criteria in the ECHA database for historical Acute Fish Toxicity data

In order to include preferably all AFT data, studies containing old species names such as *Salmo gairdneri* for rainbow trout and *Brachydanio rerio* for zebrafish were also included. Subsequently, the search criteria listed in Tab. 9 were used.

Primary search results were organized in an Excel spreadsheet. Subsequent analyses were performed on the Excel sheet with the acute toxicity data in fish, as obtained from the ECHA database and modified by means of the KNIME software (<u>www.knime.org</u>) by Dr. Stefan Scholz. The actual analyses were conducted using modified versions of the Excel data sheet as well as a KNIME workflow and appropriate pivot tables and filters. The final search dataset after reformatting with KNIME consisted of a collection of 2936 studies covering 1842 substances. The single consecutive steps of the analysis are summarized in Fig. 36.



Figure 36: Overview on the filtering steps in the ECHA dataset

Flow-chart representation of the filtering steps in the ECHA dataset

Step I: Filtering of unsuitable LC50 values

In order to obtain suitable LC_{50} concentrations, the dataset was first filtered by removing any data from plates with unsuitable entries (e.g. " $LC_{50} = x$ to y mg/L"; " $LC_{50} > ...$ " or " $LC_{50} < ...$ " or acute tests without concentration units (e.g. g/ha or ml/ha).

Step II: Filtering of duplicate studies

By revising the dataset, it became obvious that numerous studies on one substance showed the same (sequence of) LC_{50} values as those of other substances. Thus, it was assumed that data for either single studies or structurally related substances were duplicated in the dataset. These studies were removed, if either no studies for analogues of the corresponding substances could be found in the ecotoxicological information in the ECHA database or in some cases it became

obvious that mixtures of different components were used that were enlisted separately in the dataset (e.g."A mixture composed primarily of resin acids and modified resin acids such as dimers and decarboxylated resin acids" was used to cover all substances with overall 10 studies).

Steps III - V: Identification of physicochemical properties

The ECHA database was checked for the most important physicochemical properties of the substances including water solubility and log P_{0W} .

In the third step, all studies tested with an LC_{50} above the water solubility were excluded. In the fourth step, the criteria were exacerbated by eliminating all studies from the dataset, if the LC_{50} was higher than half the level of water solubility. Substances without any specifications on water solubility were also excluded.

In the fourth step, a similar procedure was used for substances with a log Pow > 4 with the exception, that substances with log Pow > 4 were kept, if the studies were performed with a chemical analysis.

In the fifth step, all remaining studies performed with inorganic substances were removed.

Studies with zebrafish (Danio rerio)

Since the purpose of this study was also to compare the AFT dataset to the FET dataset from a recent study, in a final step the remaining dataset was filtered for Acute Fish Toxicity Tests, performed with the zebrafish (*Danio rerio*).

Intra- and interspecies comparison

For a better appreciation of the reliability and consistency of LC_{50} data, inter- and intraspecies comparisons were performed. For this purpose, all substances with only one single entry per substance in the database were excluded, since for these neither intra- nor interspecies comparisons could be made.

For the interspecies comparison, the dataset was filtered to include all substances, where studies for more than one fish species were available. For the intraspecies comparison, all substances analyzed in more than one study and fish species were used.

3.4 Results of the data filtering procedure

The study was carried out with a dataset generated in July 2017, containing 2936 studies for AFT tests retrieved from the ECHA database representing a total of 1842 substances (for an overview of absolute numbers and ratios of the different steps, cf. Fig. 42 and Table 10).

3.4.1 Filtering steps I and II: Unsuitable LC₅₀ values and duplicate studies

After identification and deletion of all database entries containing no clearly defined values (LC₅₀ value ranges only given: "x to y", " \geq ", " \leq ", "ca."), 1594 studies out of original 2936 studies, representing 929 out of 1842 substances remained within the database (Fig. 37). The first filtering step ("unclear entries"; 1342 studies, 9113 substances) thus resulted in a reduction to 54.3 % and 50.5 % remaining studies and substances for further evaluation, respectively. In filtering step II, 52 studies (1.77 % of the original dataset) dealing with 20 substances (1.09 % of the original dataset) could clearly be identified as either duplicate studies of the same substance or substances tested with the same mixture of components; these were therefore removed from the dataset. Filtering steps I and II thus resulted in an elimination of 47.5 % of all studies (n = 1394) and of 50.7 % of all substances (n = 933) of the original dataset.



Figure 37: Elimination of unclear entries and duplicate studies in the ECHA dataset

Absolute numbers of studies and substances in the database during the elimination of unclear entries and duplicate studies. The elimination of unclear entries and duplicates reduced the number of studies and substances in the dataset by around 50 %.

3.4.2 Filtering steps III and IV: LC₅₀ and water solubility

When comparing the LC_{50} values with the water solubility of the substances given in the ECHA database, it became obvious that the LC_{50} of 177 studies out of the 1542 studies after filtering step II (11.49 % of studies after filtering step II) lies above water solubility. This correlates to 78 (8.58 % of studies after filtering step II) from overall 909 remaining substance (Fig. 38).



Figure 38: Entries in the ECHA database with LC₅₀ > water solubility, no data on water solubility and water solubility given as range with the LC₅₀ lying within this range

Overview on the ratio of studies and substances with LC_{50} > water solubility, no data on water solubility and water solubility stated as range with the LC50 lying within this range. The data refers to the remaining 909 substances and 1542 studies after step I and II. Studies and substances with LC50 above water solubility (light green) or without any information on water solubility (grey) were excluded from the dataset.

For 78 studies (5.06 % of the remaining dataset after step I and II) and 59 substances (6.49 % of the remaining dataset after step I and II), no data on water solubility was available in the ECHA database. Both, studies with LC_{50} > water solubility (177 studies with 78 substances) as well as substances, where no data on water solubility exists, were therefore excluded from the dataset. Additionally, for 40 studies (2.60 %, referring to 1594 studies after step I and II) and 22 substances (2.42 %, referring to 909 substances after step I and II), the water solubility was given as range with the LC_{50} lying within that range. These studies and substances remained in the datasheet for the next step, resulting in an overall of 1287 studies and 772 substances.

Figure 39: Ratio of the studies in the ECHA database with use of analytics and vehicle in studies with LC₅₀ tested above water solubility



Ratio of the use of analytics and vehicle in studies with LC₅₀ tested above water solubility. In almost the same ratio of studies, no data are available on analytics or vehicle use

Figure 40: Solvent use and analytical verification in substances from the ECHA database tested with LC₅₀ over water solubility



Overview on solvent use and analytical verification in substances tested with LC₅₀ over water solubility

In only 38.5 % of the studies tested with an LC_{50} over water solubility, a chemical analysis was performed (Fig. 39a). In 42.3 % of the cases, no information on analytics was available, and 19.2 % of the studies were conducted without any analysis. Although the selected concentrations of the test substances were above water solubility, surprisingly 41 % of the studies were not performed with a solvent. For the same ratio (41 %), no information on the use of a vehicle was available. Only in 18 % of the studies, a vehicle was used (Fig. 39b).

A detailed analysis revealed a differentiated picture (Fig. 40): For the majority (38 %) of studies, information on neither solvent use nor analytical verification was available. In 24 % of the cases, analytics were performed although the LC_{50} lies over the value of water solubility. In these studies, the measured concentrations often were available; however, the nominal concentration was used for the LC_{50} . 15 % of the studies were conducted without solvent or analytical verification. Only in 12 % of studies both, a solvent was used and analytics were performed.

The elimination of all studies with an LC_{50} higher than half the level of water solubility resulted in a further elimination of 40 studies (1.36 % of the original dataset) and 27 substances (1.47 % of the original dataset). Thus, 1247 studies corresponding to 745 substances remained for the next step, meaning 42.47 % of the studies and 40.45 % of the substances from the original dataset, respectively.

3.4.3 Elimination step V: Analysis of log Pow data

Since log Pow values refer to a specific substance, the elimination took place on the substance level, eliminating all studies from one substance, when it was not meeting the criteria. On the other hand, since the ecotoxicological information in the ECHA database does not normally include all study results, the information on analytical results was adopted for the substance. Criteria for elimination were as follows: log Pow > 4, when the studies were not supported by analytics and no data available for log Pow within the ECHA database. Additionally, another cohort was formed, if the log Pow was specified as a range from < 4 to > 4, with the LC₅₀ lying within this range. Analogously to the data on water solubility, these data remained in the dataset for further calculation. Results from the analysis of log Pow values are summarized in Fig. 41.



Figure 41:Analytical verification of the test concentrations of substances tested within the
ECHA database, when log Pow > 4

Analysis of log Pow values of substances and analytical verification of the test results when log Pow > 4

Reduction steps			Ste	pl	Ste	li q	Ste	III d	Step	N	Step	>	Step	N	Step	VII
	Origina datal	al ECHA base	Unc	lear ies	Dupli	cates	LC ₅₀ > 1 solut	Water bility	LC ₅₀ > 0.5 solub	x Water ility	Log Po	w > 4	Inorge substa	anic nces	Speciselect	ies tion
Total dataset	Subst.	Studies	Subst.	Studies	Subst.	Studies	Subst.	Studies	Subst.	Studies	Subst.	Studies	Subst.	Studies	Subst.	Studies
	1842	2936														
Without unclear entries			929	1594												
Erased unclear entries			913	1342												
Without duplicate studies					606	1542										
Erased duplicate studies					20	52										
LC ₅₀ < water solubility (ws)							750	1247								
LC _{s0} > water solubility (ws)							78	177								
No data (n.d.) on water solubility							59	78								
Water solubility stated as range							22	40								
Dataset without LC ₅₀ >ws & n.d.							772	1287								
LC ₅₀ < 0.5x water solubility									745	1247						
LC _{s0} > 0.5x water solubility									27	40						
Log Pow < 4											533	857				
Log Pow > 4											81	145				
No data Log Pow											35	85				
Log Pow < 4, with analytics											59	104				
Log Pow < 4, without analytics												10				
Log Pow < 4, n. d. on analytics											17	32				
Log Pow <4 to >4 total											96	160				
Log Pow <4 to >4 with analytics											65	103				
Log Pow <4 to >4 without analytics											×	46				
Log Pow <4 to >4 with n. d. on analytics											23	10				
Remaining Studies/substances after											689	1120				
Inorganic substances													∞	12		
Organic substances													681	1108		
All fish species except zebrafish															416	744
Zebrafish (Danio rerio) studies															265	364

Table 10:

Absolute number of studies/substances removed or considered for final analysis

Processing of the ECHA dataset for AFT data: number of studies/substances removed or considered for the final dataset. Green fields: values removed from the dataset



Figure 42: Summary of study elimination per filtering step relative to original data (%)

(Acute Fish Embryo Toxicity Test)

The chorion structure and biotransformation capacities of zebrafish as boundary conditions for OECD Test Guideline 236

Overview on studies/substances eliminated per filtering step relative to original data (%)

Based on the remaining number of substances (745), the majority (71.54 %) of substances (n = 533) tested had a log Pow below 4. Overall, 10.9 % of the substances (n=81) had log Pow values > 4; yet, the majority (8.1 %) of these substances were verified by analytics and, thus, kept in the dataset. For 96 of the substances (12.9 %), log Pow was given as range. Referring to the original dataset, only substances with log Pow < 4 were kept as well as log Pow stated as range and Log Pow > 4 when chemical analysis were performed. Thus, 37 % of substances and 38% of studies from the original dataset remained after the elimination of log Pow data, representing 689 substances and 1120 studies (Table 10; Fig. 42).

3.4.4 Further filtering: Inorganic substances

The ECHA dataset also contained a number of inorganic substances for which the log Pow is not significant. Most of them had already been deleted in previous steps. The remaining dataset still contained 8 inorganic substances with altogether 12 studies (Table 10). Following this step, 36.97 % of substances and 37.74 % of the studies relative to the original dataset remained.

3.4.5 Further filtering: AFT studies with zebrafish (Danio rerio) only

For comparison with the results from the FET study, in this step only studies with zebrafish (*Danio rerio*) were included (Table 10). 364 studies and 266 substances of the whole dataset were performed with zebrafish (*Danio rerio*), representing 12.40 % of studies and 14.44 % of substances of the total dataset.

3.4.6 Further filtering: Interspecies and intraspecies comparisons

In a further step and as preparation for subsequent inter- and intraspecies comparisons, all studies for substances with only one single entry in the database were excluded (i.e. no comparison possible). When compared to the original number of studies and substances in the ECHA database (100 %), this step resulted in a reduction of the number of substances from 682 to 453 (remaining 12.4 % of original number of substances) and a reduction of the number of studies from 1108 to 655 (remaining 22.3 % of original number of studies).

For intraspecies comparisons, 493 of the remaining studies (72.3 %) covering 181 substances (16.34 %) were available. For interspecies comparison, only 178 studies (26.0 %) representing 53 substances (4.8 %) were used.

The minimal and maximal LC_{50} values of a substance were identified, and the maximum deviation (maximum deviation = maximum value/minimum value) for each substance was calculated. For a better overview, deviations were categorized as follows (Fig. 43):

= 1	identical value
≤ 10	deviation by a factor between 1 and 10
≤ 100	deviation by a factor between 10 and 100
≤ 1000	deviation by a factor between 100 and 1000

Since in the intraspecies comparison two substances were tested with several studies in two species, two substances for the calculation of the percentile of ranges were added. The analysis illustrated that almost all studies of intra- and interspecies comparison (93.45 % and 96.22 %) are within the deviation ranges from 1-100, which is currently accepted within the REACH process.

The majority of studies per substance (85.3 % intraspecies and 86.8 % interspecies, respectively) varied by a factor ≤ 10 . In the next category with a factor ≤ 100 , interspecies comparison showed with 9.4 % of the substances nearly the same value as the intraspecies comparison with 8.2 %. Only few studies showed a deviation factor ≤ 1000 in intra- as well as in interspecies comparison. The value of the interspecies comparison (3.8 %) was about a sevenfold higher when compared to the intraspecies value (0.6 %). In the intraspecies comparison in 6.0 % of the substances, two or more studies per substance were found with exactly the same LC₅₀ values, resulting in a deviation factor of 1. On the other hand, no substance with a deviation factor of 1 could be detected in the interspecies comparison.



Figure 43: Range of deviation factors in intra- and interspecies comparisons within the ECHA database

Range of deviation factors in intra- and interspecies comparisons: values were categorized into 4 ranges: 1 for identical same values; \leq 10 for deviations by a factor between 1 and 10; \leq 100 for deviation by a factor between 10 and 100; \leq 1000 deviation by a factor between 100 and 1,000

There is a difference to the distribution in the first inter- and intraspecies comparison in the AFT status report presented to the UBA in November 2017 prior to most of the processing steps. In the November 2017 status report, 22.2 % of interspecies studies met the deviation factor \leq 100. In this dataset, nearly twice as many studies in the interspecies (5.6 %) and 4 times as many studies in the intraspecies comparison (1.9 %) showed a deviation factor \leq 1000, indicating that the processing of the initial dataset, especially the exclusion of studies with LC₅₀ > water solubility erased most of the outlier studies.

3.5 Conclusions drawn from the analysis of AFT data from the ECHA database and recommendations for future AFT studies

3.5.1 Comparison of data handling in the present AFT study and the FET by (Scholz et al., 2016)

One purpose of the present study was to give an overview on the reliability and possible elimination capacities of AFT data by setting similar criteria as in the ECHA FET study by (Scholz et al., 2016) and (Sobanska et al., 2018). However, the procedure and order of the processing steps varied between the present study and the study by (Scholz et al., 2016), and not all steps were equally performed. In case of the ECHA FET study, unclear values were not explicitly removed as were duplicate studies; in the present AFT study, studies with "predicted pH at saturation >5 or <9" and studies with "no toxicity and the maximum test concentration <10 fold above baseline toxicity" were not taken into account due to the tools not being available.

Nevertheless, parts of the findings allow a direct comparison (Table 11). The processing of studies with an LC_{50} above 0.5 times of water solubility in the ECHA FET study resulted in elimination of about 22 % for both tests and substances. In the present AFT study, this step (separated in two steps: LC_{50} > water solubility and LC_{50} > 0.5 times water solubility) reduced the number of tests by only 10.1 and % and the number substances by only 5.6 %. This indicates that – in this respect – the AFT data are obviously more robust than the FET data. On the other hand, the elimination potential of the analysis of log Pow data is almost identical in both the FET and the present AFT study.

Although the last step of the present study, the reduction of studies to those performed with the zebrafish (*Danio rerio*), is not necessarily a valid step, since historically for the AFT more fish species are used. However, it could be shown that by keeping only zebrafish data, the percentage of remaining studies is 7.6 % for the FET and 12.5 % for the AFT, i.e. very similar for both test systems. Thus, the selection of only zebrafish data does not lead to a massive loss of data for neither the FET nor the AFT analysis.

A closer look at the ECHA FET study reveals the biggest potential for elimination of data in the two steps: "Keep studies with predicted pH at saturation >5 or <9" and "Remove studies with no toxicity and maximum test concentration <10 fold above baseline toxicity", which were not performed in the analysis of AFT studies due to the tools being unavailable. This indicates that these two steps might be important for a more significant result and should be analyzed in additional studies.

	Studie	ss (n)	Substar	ices (n)	Studi	es (%)	Substan	ces (%)
	EE	AFT	FET	AFT	EE	AFT	EE	AFT
Initial Fet and AFT dataset	2065	2936	1415	1842	2065	2936	1415	1842
Remove studies with unclear LC ₅₀ - values		1594		929		54.3		50.4
Remove duplicate studies		1542		606		52.5		49.4
Remove studies when LC50 > water solubility		1287		772		43.81		41.91
Keep only studies conducted with zebrafish	2036	*	1415	*	98.60	*	100	÷
Keep only studies with organic substances	1996	*	1391	*	96.66	*	98.30	*
Keep tests with exposure duration of 96 to 120	1610	*	1191	*	76.77	*	84.17	*
Keep studies with LC50 or maximum test concentration below 0.5 times of water solubility	1155	1247	882	745	55.93	42.40	62.33	40.45
Keep studies with predicted pH at saturation >5 or <9	1131	n.p.	873	n.p.	54.77	n.p.	61.70	n.p.
Remove studies with no toxicity at maximum test concentration < 10 fold above baseline toxicity	411	ч.	343	n.p.	19.90	n.p.	24.24	ч.
Keep compounds with corresponding AFT data for fathead minnow, rainbow trout, bluegill and zebrafish	238	n.r.	185	u.r.	11.53	u.r.	13.07	n.r.
Keep only substances with Kow <4 and/or Log Kaw <4, unless the test concentrations were confirmed by analytical chemistry	156	1120	123	689	7.55	38.15	8.69	37.40
Keep only studies with organic substances		1108		681		37.74		36.97
Keep only studies conducted with zebrafish		364		266		12.40		14.44
*conducted in AFT analysis at a later timepoint; **already selected	l within the	filtering proc	bedure of orig	ginal dataset;	: n.p not pe	erformed; n.r.	- not relevan	t

The chorion structure and biotransformation capacities of zebrafish as boundary conditions for OECD Test Guideline 236 (Acute Fish Embryo Toxicity Test)

Table 11:

FET and AFT sets and numbers of substances/studies considered in final dataset

Overview of processing the FET and AFT datasets and absolute numbers of substances and studies considered for the final dataset. Green fields represent the last filtering step for both, FET and AFT analysis, the elimination of all studies except those with zebrafish.

3.5.2 Conclusions and recommendations for future AFT testing

Analyzing the AFT data from the ECHA database clearly showed that the number of studies removed from the database after applying the filtering criteria is nearly as the same as in the ECHA FET study. The largest filtering of the AFT data took place when the entries with unclear LC_{50} values were excluded. Another very critical filtering step was the analysis of LC_{50} tested above water solubility, indicating that such studies might not be fully reliable.

With regard to conducting future AFT studies for regulatory purposes, the following recommendations can be given:

- ► Only studies with clearly defined toxicity values should be accepted. Existing studies with readings like e.g. "LC₅₀ = x to y mg/L"; "LC₅₀ >..., %w/t" or "LC₅₀ <..."should be revised or rejected.</p>
- The recently revised OECD Guidance Document 23 should be consulted when testing difficult-to-test test substances (OECD, 2018).
- ► In the process of test design and the selection of test concentrations, the limits of water solubility as well as other physicochemical properties of a substance must be considered.
- For tests above water solubility, analytical validation of test concentrations should be mandatory.
- If the test concentration needs to be above water solubility, the thorough use of a vehicle should be considered.
- Measured concentrations should be given preference for ecotoxicological results. Given the importance of e.g. LC₅₀ values for risk assessment, nominal concentrations should only be accepted, if there is evidence that degradation or other losses from the system (e.g. metabolism, volatilization, adsorption to the surface of the tank, etc.) do not occur.
- Within the REACH process, elevated attention should also be paid to the preparation of test solutions. Procedures such as "ultrasonic treatment followed by filtration" or "test media with visible unsolved matter" should be discarded. In any case, as mentioned above, such procedures need to be accompanied by thorough sampling and analytical verification.
- In particular, studies with a log Pow > 4 should generally be accompanied by analytical verification.
- Given that approx. 60 % of the existing AFT data would not fulfil the requirements set in the ECHA FET study (Scholz et al., 2016; Sobanska et al., 2018), massive re-testing of acute fish toxicity would be required.
- Study results for substances tested at concentrations close or above their water solubility, but without analytical verification should be interpreted with care or considered for revision.

4 Contributions to OECD project no. 2.54: "Guidance Document on an Integrated Approach on Testing and Assessment for Fish Acute Toxicity Testing" – Integration of the Fish Embryo Test into the Threshold Approach (OECD Guidance Document 126)

In 2010, the threshold approach was adopted by the OECD (OECD, 2010) based on the observation that fish are not always the most sensitive species (Hutchinson et al., 2003; Weyers et al., 2000). In this approach, an initial Acute Fish toxicity Test (AFT) according to OECD TG 203 (OECD, 1992) is conducted at one concentration, i.e. the lowest EC₅₀/LC₅₀ derived from test responses in Daphnia and algae, and continued testing is triggered only if death is observed at this threshold concentration. Since the threshold approach was developed, the Fish Embryo Toxicity Test (OECD TG 236) has been validated and accepted by the OECD (OECD, 2013a). It seemed, therefore, appropriate to update the threshold approach to incorporate OECD TG 236, because this test offers a significant 3Rs (refinement) benefit compared to the AFT as embryos are used instead of juvenile or adult fish. As a consequence, a project proposal to update the threshold approach to incorporate the Fish Embryo Acute Toxicity Test (in particular OECD TG 236) was submitted to the OECD by Austria as the lead country and the International Council on Animal Protection in OECD Programs (ICAPO) in 2015 (OECD project no. 2.54: "Guidance Document on an Integrated Approach on Testing and Assessment for Fish Acute Toxicity Testing").

Incorporation of OECD TG 236 into the threshold approach was deemed appropriate, because LC_{50} values from the FET correlate very well with LC_{50} values from OECD TG 203 (Belanger et al., 2013; Klüver et al., 2015; Knobel et al., 2012; Lammer et al., 2009; Ratte and Hammers-Wirtz, 2003). However, some exceptions to this good correlation have been described and a review was performed by the European Chemicals Agency (ECHA) suggesting some limitations of TG 236 for regulatory use that require further consideration (Scholz et al., 2016).

Within the present UBA project, the authors of this report made continuous contributions during telephone conferences and face-to-face meetings as well as in written especially to the background document, which is designed to accompany the central updated guidance document. This background document provides relevant background information and discussions considered when updating OECD GD 126 with the aim of integrating OECD TG 236. It focuses on the characterization of the performance of the OECD TG 236 in terms of its reliability within the threshold approach and the applicability domain of OECD TG 236.

In this context, chapter 3 of this report ("Analysis of the relevance and adequateness of the Fish Acute Toxicity Test (AFT) according to OECD TG 203 to fulfil the information requirements and addressing concerns under REACH") will attempt to provide a counterpart to the corresponding ECHA FET study (Scholz et al., 2016). Likewise, the literature review in chapter 2 of the present report ("Biotransformation and bioactivation capacities in early life stages of zebrafish (*Danio rerio*)" will provide valuable input to the discussion of the applicability of the Fish Embryo Test. Since the background document has not been finalized yet, further input will be given by the authors of this report in the near future.

5 References

Adams, S.L., Zhang, T., Rawson, D.M., 2005. The effect of external medium composition on membrane water permeability of zebrafish (*Danio rerio*) embryos. Theriogen. 64, 1591-1602.

Aguirre-Martinez, G.V., Reinardy, H.C., Martin-Diaz, M.L., Henry, T.B., 2017. Response of gene expression in zebrafish exposed to pharmaceutical mixtures: Implications for environmental risk. Ecotox. Environ. Safety 142, 471-479.

Alderton, W., Berghmans, S., Butler, P., Chassaing, H., Fleming, A., Golder, Z., Richards, F., Gardner, I., 2010. Accumulation and metabolism of drugs and CYP probe substrates in zebrafish larvae. Xenobiotica 40, 547-557.

Amanuma, K., Tone, S., Saito, H., Shigeoka, T., Aoki, Y., 2002. Mutational spectra of benzo[a]pyrene and MelQx in rpsL transgenic zebrafish embryos. Mutat. Res. 513, 83-92.

Arukwe, A., Goksoyer, A., 2003. Eggshell and egg yolk proteins in fish: hepatic proteins for the next generation: oogenetic, population, and evolutionary implication of endocrine disruption. Comp. Hepatol. 2, 1-21.

Belanger, S.E., Rawlings, J.M., Carr, G., 2013. Use of fish embryo toxicity tests for the prediction of acute fish toxicity to chemicals. Environ. Toxicol. Chem. 32, 1768-1783.

Blaschke, U., Paschke, A., Rensch, I., Schuurmann, G., 2010. Acute and chronic toxicity toward the bacteria *Vibrio fischeri* of organic narcotics and epoxides structural alerts for epoxide excess toxicity. Chem. Res. in Toxicol. 23, 1936-1946.

Böhler, S., 2012. The fathead minnow embryo as a model for the development of alternative testing methods in ecotoxicology. Diploma thesis, Dept. of Zoology, Heidelberg University, 136 pp.

Boix, N., Teixido, E., Vila-Cejudo, M., Ortiz, P., Ibáñez, E., Llobet, J.M., Barenys, M., 2015. Triclaben-dazole sulfoxide causes stage-dependent embryo lethality in zebrafish and mouse *in vitro*. PLoS One 10(3), e0121308.

Braunbeck, T., Kais, B., Lammer, E., Otte, J., Schneider, K., Stengel, D., Strecker, R., 2015. The fish embryo test (FET): origin, applications, and future. Environ. Sci. Pollut. Res. 22, 16247–16261.

Braunbeck, T., Lammer, E., Leist, E., Rudolf, M., 2005. Towards an alternative for the acute fish LC_{50} test in chemical assessment: the zebrafish (*Danio rerio*) embryo toxicity test – an update. ALTEX 22, 87-102.

Braunbeck, T., Völkl, A., 1991. Induction of biotransformation in the liver of eel (*Anguilla anguilla* L.) by sublethal exposure to dinitro-*o*-cresol: an ultrastructural and biochemical study. Ecotoxicol. Environ. Safety 21, 109-127.

Braunbeck, T., Böhler, S., Strecker, R., Fedderwitz, F., Henn, K., Kais, B., Lammer, E., Schneider K., Weigt, S., 2012. Der Fischembryotest als Alternativmethode für den akuten Fischtest – abschließend notwendige Laboruntersuchungen und Datenanalysen zur Validierung des Fischembryotests für das OECD Prüfrichtlinienprogramm. Bericht an das Umweltbundesamt (FKZ 3708 65 400).

Braunig, J., Schiwy, S., Broedel, O., Muller, Y., Frohme, M., Hollert, H., Keiter, S.H., 2015. Time-dependent expression and activity of cytochrome P450 1s in early life-stages of the zebrafish (*Danio rerio*). Environ. Sci. Pollut. Res. 22, 16319-16328.

Brivio, M., Bassi, R., Cotelli, F., 1991. Identification and characterization of the major components of the *Oncorhynchus mykiss* egg chorion. Mol. Reprod. Dev. 28, 85-93.

Brox, S., Seiwert, B., Haase, N., Kuster, E., Reemtsma, T., 2016a. Metabolism of clofibric acid in zebrafish embryos (*Danio rerio*) as determined by liquid chromatography-high resolution-mass spectrometry. Comp. Biochem. Physiol. 185 C, 20-28.

Brox, S., Seiwert, B., Kuster, E., Reemtsma, T., 2016b. Toxicokinetics of polar chemicals in zebrafish embryo (*Danio rerio*): Influence of physicochemical properties and of biological processes. Environ. Sci. Technol. 50, 10264-10272.

Buhler, D.R., Williams, D.E., 1988. The role of biotransformation in the toxicity of chemicals. Aquat. Toxicol. 11, 19-28.

Busquet, F., Nagel, R., von Landenberg, F., Mueller, S.O., Huebler, N., Broschard, T.H., 2008a. Development of a new screening assay to identify proteratogenic substances using zebrafish *Danio rerio* embryo combined with an exogenous mammalian metabolic activation system (mDarT). Toxicol. Sci. 104, 177-188.

Carlsson, G., Patring, J., Kreuger, J., Norrgren, L., Oskarsson, A., 2013. Toxicity of 15 veterinary pharmaceuticals in zebrafish (*Danio rerio*) embryos. Aquat. Toxicol. 126, 30-41.

Carlsson, G., Patring, J., Ulleras, E., Oskarsson, A., 2011. Developmental toxicity of albendazole and its three main metabolites in zebrafish embryos. Reprod. Toxicol. 32, 129-137.

Celander, M.C., 2011. Cocktail effects on biomarker responses in fish. Aquat. Toxicol. 105, 72-77.

Chambers, J.E., Yarbrough, J.D., 1976. Xenobiotic Biotransformation Systems in Fishes. Comp. Biochem. Physiol. 55C, 77-84.

Chang, C.T., Chung, H.Y., Su, H.T., Tseng, H.P., Tzou, W.S., Hu, C.H., 2013. Regulation of zebrafish CYP3A65 transcription by AHR2. Toxicol. Appl. Pharmacol. 270, 174-184.

Cheng, J., Flahaut, E., Cheng, S.H., 2007. Effect of carbon nanotubes on developing zebrafish (Danio rerio) embryos. Environ. Toxicol. Chem. 26, 708-716.

Chng, H.T., 2013. Evaluation of alternative *in vivo* and *in vitro* models for drug metabolism testing in drug discovery, Pharmacy. Nat. Univ. of Singapore.

Chng, H.T., Ho, H.K., Yap, C.W., Lam, S.H., Chan, E.C.Y., 2012. An investigation of the bioactivation potential and metabolism profile of zebrafish versus human. J. Biomol. Screen. 17, 974-986.

Collodi, P., Kamei, Y., Ernst, T., Miranda, C., Buhler, D.R., Barnes, D.W., 1992. Culture of cells from zebrafish (*Brachydanio rerio*) embryo and adult tissues. Cell. Biol. Toxicol. 8, 43-61.

Corley-Smith, G.E., Su, H.T., Wang-Buhler, J.L., Tseng, H.P., Hu, C.H., Hoang, T., Chung, W.G., Buhler, D.R., 2006. CYP3C1, the first member of a new cytochrome P450 subfamily found in zebrafish (*Danio rerio*). Biochem. Biophys. Res. Comm. 340, 1039-1046.

Coward, K., Bromage, N.R., Hibbitt, O., Parrington, J., 2002. Gamete physiology, fertilization and egg activation in teleost fish. Rev. Fish Biol. Fisheries 12, 33-58.

Creton, R., 2004. The calcium pump of the endoplasmic reticulum plays a role in midline signaling during early zebrafish development. Brain. Res. Dev. Brain Res. 151, 33-41.

Creusot, N., Brion, F., Piccini, B., Budzinski, H., Porcher, J.M., Ait-Aissa, S., 2015. BFCOD activity in fish cell lines and zebrafish embryos and its modulation by chemical ligands of human aryl hydrocarbon and nuclear receptors. Environ. Sci. Poll. Res. 22, 16393-16404.

Cunha, V., Rodrigues, P., Santos, M.M., Moradas-Ferreira, P., Ferreira, M., 2016. *Danio rerio* embryos on Prozac – Effects on the detoxification mechanism and embryo development. Aquat. Toxicol. 178, 182-189.

Donato, M.T., Castell, J.V., 2003. Strategies and molecular probes to investigate the role of cytochrome P450 in drug metabolism – Focus on *in vitro* studies. Clin. Pharmacokinet. 42, 153-178.

Donovan, M., Hart, N.H., 1983. Fine structure of the chorion and site of sperm entry in the egg of *Brachydanio*. J. Exp. Zool. 227, 277-296.

EU, 2010. Directive 2010/63/EU of the European parliament and of the council of 22 September 2010 on the protection of animals used for scientific purposes. Off. J. EU L 276, 33-79.

Fantel, A.G., 1982. Culture of whole rodent embryos in teratogen screening. Teratogen. Carcinogen. Mut. 2, 231-242.

Fedderwitz, F., 2008. Morphologie und Ultrastruktur des Chorions vom Zebrabärbling (*Danio rerio*), Dept. of Zoology. Univ. of Heidelberg, Heidelberg, 46 pp.

Fetter, E., Smetanova, S., Baldauf, L., Lidzba, A., Altenburger, R., Schuttler, A., Scholz, S., 2015. Identification and characterization of androgen-responsive genes in zebrafish embryos. Environ. Sci. Technol. 49, 11789-11798.

Funari, E., Zoppini, A., Verdina, A., Deangelis, G., Vittozzi, L., 1987. Xenobiotic-metabolizing enzyme-systems in test fish. 1. Comparative studies of liver microsomal monooxygenases. Ecotox. Environ. Safety 13, 24-31.

Gellert, G., Heinrichsdorff, J., 2001. Effect of age on the susceptibility of zebrafish eggs to industrial wastewater. Water Res. 35, 3754-3757.

Gilkey, J.C., Jaffe, L.F., Ridgway, E.B., Reynolds, G.T., 1978. A free calcium wave traverses the activating egg of the medaka, *Oryzias latipes*. J. Cell. Biol. 76, 448-466.

Glisic, B., Hrubik, J., Fa, S., Dopudj, N., Kovacevic, R., Andric, N., 2016. Transcriptional profiles of glutathione-*S*-Transferase isoforms, Cyp, and AOE genes in atrazine-exposed zebrafish embryos. Environ. Toxicol. 31, 233-244.

Goldstone, J.V., Goldstone, H.M.H., Morrison, A.M., Tarrant, A., Kern, S.E., Woodin, B.R., Stegeman, J.J., 2007. Cytochrome p450 1 genes in early deuterostomes (tunicates and sea urchins) and vertebrates (chicken and frog): Origin and diversification of the CYP1 gene family. Mol. Biol. Evol. 24, 2619-2631.

Goldstone, J.V., Jonsson, M.E., Behrendt, L., Woodin, B.R., Jenny, M.J., Nelson, D.R., Stegeman, J.J., 2009. Cytochrome P450 1D1: a novel CYP1A-related gene that is not transcriptionally activated by PCB126 or TCDD. Arch. Biochem. Biophys. 482, 7-16.

Goldstone, J.V., McArthur, A.G., Kubota, A., Zanette, J., Parente, T., Jonsson, M.E., Nelson, D.R., Stegeman, J.J., 2010. Identification and developmental expression of the full complement of cytochrome P450 genes in zebrafish. BMC Genomics 11, 643.

Gonzalez-Doncel, M., Fernandez-Torija, C., Hinton, D.E., Tarazona, J.V., 2004. Stage-specific toxicity of cypermethrin to medaka (*Oryzias latipes*) eggs and embryos using a refined methodology for an *in vitro* fertilization bioassay. Arch. Environ. Contam. Toxicol. 48, 87-98.

Guengerich, F.P., 2000. Metabolism of chemical carcinogens. Carcinogenesis 21, 345-351.

Guengerich, F.P., 2007. Mechanisms of cytochrome P450 substrate oxidation: mini review. J. Biochem. Mol. Toxicol. 21, 163-168.

Hagedorn, M., Hsu, E., Kleinhans, F.W., Wildt, D.E., 1997a. New approaches for studying the permeability of fish embryos: toward successful cryopreservation. Cryobiology 34, 335-347.

Hagedorn, M., Kleinhans, F.W., Wildt, D.E., Rall, W.F., 1997b. Chill sensitivity and cryoprotectant permeability of dechorionated zebrafish embryos, *Brachydanio rerio*. Cryobiology 34, 251-263.

Hagedorn, M., Kleinhans, F.W., Artemov, D., Pilatus, U., 1998. Characterization of a major permeability barrier in the zebrafish embryo. Biol. Reprod. 59, 1240-1250.

Hart, N.H., Collins, G.C., 1991. An electron microscope and freeze-fracture study of the egg cortex of *Brachydanio rerio*. Cell Tissue Res. 265, 317-328.

Hart, N.H., Pietri, R., Donovan, M., 1984. The structure of the chorion and associated surface filaments in *Oryzias* – evidence for the presence of extracellular tubules. J. Exp. Zool. 230, 273-296.

Harvey, B., Kelley, N.R., Ashwood-Smith, M.J., 1983. Permeability of intact and dechorionated zebrafish embryos to glycerol and dimethyl sulfoxide. Cryobiology 20, 432-439.

Henn, K., Braunbeck, T., 2011. Dechorionation as a tool to improve the fish embryo toxicity test (FET) with the zebrafish (*Danio rerio*). Comp. Biochem. Physiol. 153C, 91-98.

Hisaoka, K.K., 1958. Microscopic studies of the teleost chorion. Trans. Amer. Microsc. Soc. 77, 240 - 243.

Hisaoka, K.K., Firlit, C.F., 1960. Further studies on the embryonic development of the zebrafish, *Brachydanio rerio* (Hamilton-Buchanan). J. Morph. 107, 205-225.

Hsu, T., Deng, F.Y., 1996. Studies on the susceptibility of various organs of zebrafish (*Brachydanio rerio*) to benzo(a)pyrene-induced DNA adduct formation. Chemosphere 33, 1975-1980.

Hutchinson, T.H., Barrett, S., Busby, M., Constables, D., Hartmann, A., Hayes, E., Huggett, D., Länge, R., Lillicrap, A.D., Straub, J.O., Thompson, R.S., 2003. A strategy to reduce the numbers of fish used in acute ecotoxicity testing of pharmaceuticals. Environ. Tocicol. Chem. 22, 60-65.

Inohaya, K., Yasumasu, S., Araki, K., Naruse, K., Yamazaki, K., Yasumasu, I., Iuchi, I., Yamagami, K., 1997. Species-dependent migration of the fish hatching gland cells that commonly express astacin-like proteases in common. Dev. Growth Diff. 39, 191-197.

Jones, H.S., Panter, G.H., Hutchinson, T.H., Chipman, J.K., 2010. Oxidative and conjugative xenobiotic metabolism in zebrafish larvae *in vivo*. Zebrafish 7, 23-30.

Jones, H.S., Trollope, H.T., Hutchinson, T.H., Panter, G.H., Chipman, J.K., 2009. Assessment of an ibuprofen metabolism by zebrafish larvae, using liquid chromatography-mass spectrometry (LC-MS). Toxicology 262, 14-15.

Jones, H.S., Trollope, H.T., Hutchinson, T.H., Panter, G.H., Chipman, J.K., 2012. Metabolism of ibuprofen in zebrafish larvae. Xenobiotica 42, 1069-1075.

Jonsson, M.E., Brunstrom, B., Brandt, I., 2009. The zebrafish gill model: Induction of CYP1A, EROD and PAH adduct formation. Aquat. Toxicol. 91, 62-70.

Jonsson, M.E., Jenny, M.J., Woodin, B.R., Hahn, M.E., Stegeman, J.J., 2007a. Role of AHR2 in the expression of novel cytochrome p450 1 family genes, cell cycle genes, and morphological defects in developing zebra fish exposed to 3,3 ',4,4 ',5-pentachlorobiphenyl or 2,3,7,8-tetrachlorodibenzo-p-dioxin. Toxicol. Sci. 100, 180-193.

Jonsson, M.E., Orrego, R., Woodin, B.R., Goldstone, J.V., Stegeman, J.J., 2007b. Basal and 3,3',4,4',5pentachlorobiphenyl-induced expression of cyto-chrome P450 1A, 1B and 1C genes in zebrafish. Toxi-col. Appl. Pharmacol. 221, 29-41.

Kais, B., 2009. The distribution of different fluorescence dyes in the egg and embryo of the zebrafish (*Danio rerio*). Diploma thesis, Dept. of Zoology, Heidelberg University.

Kais, B., Schneider, K.E., Keiter, S., Henn, K., Ackermann, C., Braunbeck, T., 2013. DMSO modifies the permeability of the zebrafish (*Danio rerio*) chorion-Implications for the fish embryo test (FET). Aquat. Toxicol. 140, 229-238.

Kais, B., Schiwy, S., Hollert, H., Keiter, S.H., Braunbeck, T., 2017. *In vivo* EROD assays with the zebrafish (*Danio rerio*) as rapid screening tools for the detection of dioxin-like activity. Sci Total Environ. 590-591: 269-280

Kais, B., Ottermanns, R., Scheller, F., Braunbeck, T., 2018. Modification and quantification of *in vivo* EROD liveimaging with zebrafish (*Danio rerio*) embryos to detect both induction and inhibition of CYP1A. Sci. Total Environ. 615, 330-347. Kantae, V., Krekels, E.H.J., Ordas, A., Gonzalez, O., van Wijk, R.C., Harms, A.C., Racz, P.I., van der Graaf, P.H., Spaink, H.P., Hankemeier, T., 2016. Pharmacokinetic modeling of paracetamol uptake and clearance in zebrafish larvae: expanding the allometric scale in vertebrates with five orders of magnitude. Zebrafish 13, 504-510.

Kim, D.-H., Sun, Y., Seok, Y.H., Lee, S., Kim, B., 2005. Investigating chorion softening of zebrafish embryos with a microrobotic force sensing system. J. Biomech. 38, 1359-1363.

Klüver, N., König, M., Ortmann, J., Massei, R., Paschke, A., Kuehne, R., Scholz, S., 2015. The fish embryo toxicity test (FET) – identification of com-pounds with weak toxicity and analysis of behavioral effects to improve prediction of acute toxicity for neurotoxic compounds. Environ. Sci. Technol. 49, 7002-7011.

Kluver, N., Ortmann, J., Paschke, H., Renner, P., Ritter, A.P., Scholz, S., 2014. Transient overexpression of adh8a increases allyl alcohol toxicity in zebrafish embryos. Plos One 9, e90619.

Knöbel, M., Busser, F.J.M., Rico-Rico, A., Kramer, N.I., Hermens, J.L.M., Hafner, C., Tanneberger, K., Schirmer, K., Scholz, S., 2012. Predicting adult fish acute lethality with the zebrafish embryo: relevance of test duration, endpoints, compound properties, and exposure concentration analysis. Environ. Sci. Technol. 46, 9690-9700.

Kubota, A., Bainy, A.C., Woodin, B.R., Goldstone, J.V., Stegeman, J.J., 2013. The cytochrome P450 2AA gene cluster in zebrafish (*Danio rerio*): expression of CYP2AA1 and CYP2AA2 and response to phenobarbital-type inducers. Toxicol. Appl. Pharmacol. 272, 172-179.

Kubota, A., Goldstone, J.V., Lemaire, B., Takata, M., Woodin, B.R., Stegeman, J.J., 2015. Role of pregnane X receptor and aryl hydrocarbon receptor in transcriptional regulation of pxr, CYP2, and CYP3 genes in developing zebrafish. Toxicol. Sci. 143, 398-407.

Kuhnert, A., Vogs, C., Seiwert, B., Aulhorn, S., Altenburger, R., Hollert, H., Kuster, E., Busch, W., 2017. Biotransformation in the zebrafish embryo – temporal gene transcription changes of cytochrome P450 enzymes and internal exposure dynamics of the AhR binding xenobiotic benz[a]anthracene. Environ. Pollut. 230, 1-11.

Lammer, E., Carr, G.J., Wendler, K., Rawlings, J.M., Belanger, S.E., Braunbeck, T., 2009. Is the fish embryo toxicity test (FET) with the zebrafish (*Danio rerio*) a potential alternative for the fish acute toxicity test? Comp. Biochem. Physiol. 149C, 196-209.

Le Fol, V., Brion, F., Hillenweck, A., Perdu, E., Bruel, S., Ait-Aissa, S., Cravedi, J.P., Zalko, D., 2017a. Comparison of the *in vivo* biotransformation of two emerging estrogenic contaminants, BP2 and BPS, in zebrafish embryos and adults. Int. J. Mol. Sci. 18, 704.

Leonard, M., Vanpoucke, M., Petit-Poulsen, V., Porcher, J.M., 2005. Evaluation of the fish embryo test as a potential alternative to the standard acute fish toxicity test OECD 203, International Symposium on Toxicology Assessment, Skathios, Greece.

Li, C., Luo, L., Awerman, J., McGrath, P., 2011a. Whole zebrafish cytochrome P450 assay for assessing drug metabolism and safety. Zebrafish: methods for assessing drug safety and toxicity, 103-115.

Li, F., Ji, C., Qin, N., 2011b. Mass spectrometry in drug metabolism and disposition: basic principles and applications.

Li, Y., Wang, H., Si, N., Ren, W., Han, L., Xin, S., Zuo, R., Wei, X., Yang, J., Zhao, H., Bian, B., 2015. Metabolic profiling analysis of berberine, palmatine, jatrorrhizine, coptisine and epiberberine in zebrafish by ultra-high performance liquid chromatography coupled with LTQ Orbitrap mass spectrometer. Xenobiotica 45, 302-311.

Liu, H., Nie, F.H., Lin, H.Y., Ma, Y., Ju, X.H., Chen, J.J., Gooneratne, R., 2016. Developmental toxicity, EROD, and CYP1A mRNA expression in zebrafish embryos exposed to dioxin-like PCB126. Environ. Toxicol. 31, 201-210.

Lörracher, A.-K., Braunbeck, T. 2019. Biotransformation capacities in embryonic, larval and adult zebrafish (*Danio rerio*). Submitted for publication (under review).

Mattsson, A., Ulleras, E., Patring, J., Oskarsson, A., 2012. Albendazole causes stage-dependent develop-mental toxicity and is deactivated by a mammalian metabolization system in a modified zebrafish embryo toxicity test. Reprod. Toxicol. 34, 31-42.

Miller, J.A., 1970. Carcinogenesis by chemicals – an overview. Cancer Res. 30, 559-576.

Mizell, M., Romig, E.S., 1997. The aquatic vertebrate embryo as a sentinel for toxins: zebrafish embryo dechorionation and perivitelline space microinjection. Int. J. Dev. Biol. 41, 411-423.

Mold, D.E., Kim, I.F., Tsai, C.-M., Lee, D., Chang, C.-Y., Huang, R.C.C., 2001. Cluster of genes encoding the major egg envelope protein of zebrafish. Mol. Reprod. Dev. 58, 4-14.

Monod, G., Boudry, M.-A., Gillet, C., 1996. Biotransformation enzymes and their induction by b-naphthoflavone during embryo-larval development in salmonid species. Comp. Biochem. Physiol. 114C, 45-50.

Nebert, D.W., Dalton, T.P., 2006. The role of cytochrome P450 enzymes in endogenous signaling pathways and environmental carcinogenesis. Nat. Rev. Cancer 6, 947-960.

Noury, P., Geffard, O., Tutundjian, R., Garric, J., 2006. Non-destructive *in vivo* measurement of ethoxyresorufin biotransformation by zebrafish pro-larvae: development and application. Environ. Toxicol. 21, 324-331.

OECD, 1992. OECD guideline for the testing of chemicals. Section 2: Effects on biotic systems. OECD Test Guideline 203: Fish, Acute toxicity test. Paris, France: Organization for Economic Cooperation and Development.

OECD, 2000. Guidance document on aquatic toxicity testing of difficult substances and mixtures. OECD Series on Testing and Assessment 23, 53 p.

OECD, 2010. Short Guidance on the Threshold approach for Acute Fish Toxicity, Series on Testing and Assessment No 126, ENV/JM/TG(2010)/7, OECD, Paris.

OECD, 2013a. OECD Guidelines for the Testing of Chemicals. Section 2: Effects on Biotic Systems Test No. 236: Fish Embryo Acute Toxicity (FET) Test. Paris, France: Organization for Economic Cooperation and Development.

OECD, 2013b. OECD guidelines for the testing of chemicals: 210 - Fish, early-life stage toxicity tests. OECD, Paris.

Oris, J.T., Belanger, S.E., Bailer, A.J., 2012. Baseline characteristics and statistical implications for the OECD 210 fish early life-stage chronic toxicity test. Environ. Toxciol. Chem. 31, 370-376.

Otte, J.C., Schmidt, A.D., Hollert, H., Braunbeck, T., 2010. Spatio-temporal development of CYP1 activity in early life-stages of zebrafish (*Danio rerio*). Aquat. Toxicol. 100, 38-50.

Otte, J., Schultz, B., Fruth, D., Fabian, E., Van Ravenzwaay, B., Hidding, B., Salinas, E., 2017. Intrinsic xenobiotic metabolizing enzyme activities in early life stages of zebrafish (*Danio rerio*). Toxicol. Sci. 159, 86-93.

Oziolor, E.M., Carey, A.N., Matson, C.W., 2017. A non-destructive BFCOD assay for *in vivo* measurement of cytochrome P450 3A (CYP3A) enzyme activity in fish embryos and larvae. Ecotoxicology 6, 809-819.

Ozoh, P.T., 1980. Effects of reversible incubations of zebrafish eggs in copper and lead ions with or without shell membranes. Bull. Enviro.n Contam. Toxicol. 24, 270-275.

Parkinson, A., Ogilvie, B.W., 2001. Biotransformation of xenobiotics. McGraw-Hill New York.

Pauka, L.M., Maceno, M., Rossi, S.C., Silva de Assis, H.C., 2011. Embryo toxicity and biotransformation responses in zebrafish exposed to water-soluble fraction of crude oil. Bull. Environ. Contam. Toxicol. 86, 389-393.

Pelka, K.E., Henn, K., Keck, A., Sapel, B., Braunbeck, T., 2017. Size does matter – Determination of the critical molecular size for the uptake of chemicals across the chorion of zebrafish (*Danio rerio*) embryos. Aquat. Toxicol. 185, 1-10.

Peng, X., Shang, G., Wang, W., Chen, X., Lou, Q., Zhai, G., Li, D., Du, Z., Ye, Y., Jin, X., He, J., Zhang, Y., Yin, Z., 2017. Fatty acid oxidation in zebrafish adipose tissue is promoted by 1alpha,25(OH)2D3. Cell. Rep. 19, 1444-1455.

Pesonen, M., Andersson, T., 1991. Characterization and Induction of Xenobiotic Metabolizing enzyme-activities in a primary culture of rainbow trout hepatocytes. Xenobiotica 21, 461-471.

Poon, K.L., Wang, X.G., Lee, S.G.P., Ng, A.S., Goh, W.H., Zhao, Z.H., Al-Haddawi, M., Wang, H.S., Mathavan, S., Ingham, P.W., McGinnis, C., Carney, T.J., 2017a. Transgenic zebrafish reporter lines as alternative *in vivo* organ toxicity models. Toxicol. Sci. 156, 133-148.

Poon, K.L., Wang, X.G., Ng, A.S., Goh, W.H., McGinnis, C., Fowler, S., Carney, T.J., Wang, H.S., Ingham, P.W., 2017b. Humanizing the zebrafish liver shifts drug metabolic profiles and improves pharmacokinetics of CYP3A4 substrates. Arch. Toxicol. 91, 1187-1197.

Ratte, H.T., Hammers-Wirtz, M., 2003. Evaluation of the existing database from the fish embryo test. UBA report no. 363 01 062. 27 pp.

Rawson, D.W., Zhang, T., Kalicharan, D., Jongebloed, W.L., 2000. Field emission scanning electron microscopy and transmission electron microscopy studies of the chorion, plasma membrane and syncytial layers of the gastrula-stage embryo of the zebrafish *Brachydanio rerio*: a consideration of the structural and functional relationships with respect to cryoprotectant penetration. Aquacult. Res. 31, 325 - 336.

Renwick, A.B., Lewis, D.F.V., Fulford, S., Surry, D., Williams, B., Worboys, P.D., Cai, X., Wang, R.W., Price, R.J., Lake, B.G., Evans, D.C., 2008. Metabolism of 2,5-bis(trifluoromethyl)-7-benzyloxy-4-trifluorome-thylcoumarin by human hepatic CYP isoforms: evidence for selectivity towards CYP3A4. Xenobiotica 31, 187-204.

Saad, M., Cavanaugh, K., Verbueken, E., Pype, C., Casteleyn, C., Van Ginneken, C., Van Cruchten, S., 2016a. Xenobiotic metabolism in the zebrafish: a review of the spatiotemporal distribution, modulation and activity of Cytochrome P450 families 1 to 3. J. Toxicol. Sci. 41, 1-11.

Saad, M., Verbueken, E., Pype, C., Casteleyn, C., Van Ginneken, C., Maes, L., Cos, P., Van Cruchten, S., 2016b. *In vitro* CYP1A activity in the zebrafish: temporal but low metabolite levels during organogenesis and lack of gender differences in the adult stage. Reprod. Toxicol. 64, 50-56.

Saad, M., Matheeussen, A., Bijttebier, S., verbueken, E., Pype, C., Casteleyn, C., Van Ginneken, C., Maes, L., Cos, P., Van Cruchten, S., 2017. *In vitro* CYP-mediated drug metabolism in the zebrafish (embryo) using human reference compounds. Toxicol. in vitro 42, 329-336

Schlenk, D., Celander, M., Gallagher, E.P., George, S., James, M., Kullman, S.W., van den Hurk, P., Willett, K., 2008. Biotransformation in fishes. The toxicology of fishes, 153-234.

Scholz, S., Fischer, S., Gündel, U., Küster, E., Luckenbach, T., Voelker, D., 2008. The zebrafish embryo model in environmental risk assessment – applications beyond acute toxicity testing. Environ. Sci. Pollut. Res. 15, 394-404.

Scholz, S., Klüver, N., Kühne, R., 2016. Analysis of the relevance and adequateness of using Fish Embryo Acute Toxicity (FET) Test Guidance (OECD 236) to fulfil the information requirements and addressing concerns under REACH. Report ECHA-UFZ contract ECHA/2014/341.

Scholz, S., Ortmann, J., Klüver, N., Leonard, M., 2014. Extensive review of fish embryo acute toxicities for the prediction of GHS acute systemic toxicity categories. Regul. Toxicol. Pharmacol. 69, 572-579.

Scholz, S., Sela, E., Blaha, L., Braunbeck, T., Galay-Burgos, M., García-Franco, M., Guinea, J., Klüver, N., Schirmer, K., Tanneberger, K., Tobor-Kapłon, M., Witters, H., Belanger, S., Benfenati, E., Creton, S., Cronin, M.T.D., Eggen, R.I.L., Embry, M., Ekman, D., Gourmelon, A., Halder, M., Hardy, B., Hartung, T., Hubesch, B., Jungmann, D., Lampi, M.A., Lee, L., Léonard, M., Küster, E., Lillicrap, A., Luckenbach, T., Murk, A.J., Navas, J.M., Peijnenburg, W., Repetto, G., Salinas, E., Schüürmann, G., Spielmann, H., Tollefsen, K.E., Walter-Rohde, S., Whale, G., Wheeler, J.R., Winter, M.J., 2013. A European perspective on alter-natives to animal testing for environmental hazard identification and risk assessment. Regul. Toxicol. Parmacol. 67, 506-530.

Schoots, A.F.M., Stikkelbroeck, J.J.M., Bekhuis, J.F., Denucé, J.M., 1982. Hatching in teleostean fishes: Fine structural changes in the egg envelope during enzymatic breakdown *in vivo* and *in vitro*. J. Ultrastr. Res. 80, 185-196.

Scornaienchi, M.L., Thornton, C., Willett, K.L., Wilson, J.Y., 2010. Functional differences in the cytochrome P450 1 family enzymes from zebrafish (*Danio rerio*) using heterologously expressed proteins. Arch. Biochem. Biophys. 502, 17-22.

Selman, K., Wallace, R.A., Sarka, A., Qi, X., 1993. Stages of oocyte development in the zebrafish, *Brachydanio rerio*. J. Morphol. 218, 203 - 224.

Shaya, L., Dejong, C., Wilson, J.Y., 2014. Expression patterns of cytochrome P450 3B and 3C genes in model fish species. Comp. Biochem. Physiol. 166C, 115-125.

Siegenthaler, P.F., Zhao, Y., Zhang, K., Fent, K., 2017. Reproductive and transcriptional effects of the antiandrogenic progestin chlormadinone acetate in zebrafish (*Danio rerio*). Environ. Poll. 223, 346-356.

Sobanska, M., Scholz, S., Nyman, A.M., Cesnaitis, R., Gutierrez Alonso, S., Klüver, N., Kühne, R., Tyle, H., de Knecht, J., Dang, Z., Lundbergh, I., Carlon, C., De Coen, W., 2018. Applicability of the fish embryo acute toxicity (FET) test (OECD 236) in the regulatory context of Registration, Evaluation, Authorisation, and Restriction of Chemicals (REACH). Environ. Toxicol. Chem. 37, 657-670.

Spitsbergen, J., Kent, M., 2003. The state of the art of the zebrafish model for toxicology and toxicologic pathology research – advantages and current limitations. Toxicol. Pathol. 31 Suppl., 62-87.

Stegeman, J.J., Behrendt, L., Woodin, B.R., Kubota, A., Lemaire, B., Pompon, D., Goldstone, J.V., Urban, P., 2015. Functional characterization of zebrafish cytochrome P450 1 family proteins expressed in yeast. Biochim. Biophys. Acta 1850, 2340-2352.

Stegeman, J.J., Goldstone, J.V., Hahn, M.E., 2010. 10 - Perspectives on zebrafish as a model in environmental toxicology. In: Steve F. Perry, M.E.A.P.F., Colin, J.B. (Eds.), Fish Physiology. Academic Press, pp. 367-439.

Stouthart, A.J.H.X., Spanings, F.A.T., Lock, R.A.C., Wendelaar Bonga, S.E., 1994. Effects of low water pH on lead toxicity to early life stages of the common carp (*Cyprinus carpio*). Aquat. Toxicol. 30, 137-151.

Strähle, U., Scholz, S., Geisler, R., Greiner, P., Hollert, H., Rastegar, S., Schumacher, A., Selderslaghs, I., Weiss, C., Witters, H., Braunbeck, T., 2012. Zebrafish embryos as an alternative to animal experiments - a commentary on the definition of the onset of protected life stages in animal welfare regulations. Reprod. Toxicol. 33, 245-253.

Stuart, G.W., Vielkind, J.R., McMurray, J.V., Westerfield, M., 1990. Stable lines of transgenic zebrafish exhibit reproducible patterns of transgene expression. Development 109, 577-584.

Taylor, A.E., 2005. Immunohistochemical localization of cytochrome P450s 1A, 2K6, 2K7, 3A65 and 3C1 and expression of P4501A in tumor sensitive and resistant lines of juvenile zebrafish, BioRes. Res. Oregon State University.

Troxel, C.M., Reddy, A.P., ONeal, P.E., Hendricks, J.D., Bailey, G.S., 1997. *In vivo* aflatoxin B-1 metabolism and hepatic DNA adduction in zebrafish (*Danio rerio*). Toxicol. Appl. Pharmacol. 143, 213-220.

Tseng, H.P., Hseu, T.H., Buhler, D.R., Wang, W.D., Hu, C.H., 2005. Constitutive and xenobiotics-induced expression of a novel CYP3A gene from zebrafish larva. Toxicol. Appl. Pharmacol. 205, 247-258.

Tsien, R.Y., Waggoner, A., Pawley, J.B., (eds) 1995. Fluorophores for confocal microscopy. Handbook of biological confocal microscopy. Springer Netherlands, pp. 267-274.

Van Leeuwen, C.J., Griffioen, P.S., Vergouw, W.H.A., Maas-Diepeveen, J.L., 1985. Differences in susceptibility of early life stages of rainbow trout (*Salmo gairdneri*) to environmental pollutants. Aquat. Toxicol. 7, 59-78.

Verbueken, E., Alsop, D., Saad, M.A., Pype, C., Van Peer, E.M., Casteleyn, C.R., Van Ginneken, C.J., Wilson, J., Van Cruchten, S.J., 2017. *In vitro* biotransformation of two human CYP3A. Probe substrates and their inhibition during early zebrafish development. Int. J. Mol. Sci. 18(1), pii: E217.

Villalobos, S.A., Papoulias, D.M., Meadows, J., Blankenship, A.L., Pastva, S.D., Kannan, K., Hinton, D.E., Tillit, D.E., Giesy, J.P., 2000. Toxic responses of medaka, d-rR strain, to polychlorinated naphthalene mixtures after embryonic exposure by in ovo nanoinjection: a partial life-cycle assessment. Environ. Toxicol. Chem. 19, 432-440.

Wang-Buhler, J.L., Lee, S.J., Chung, W.G., Stevens, J.F., Tseng, H.P., Hseu, T.H., Hu, C.H., Westerfield, M., Yang, Y.H., Miranda, C.L., Buhler, D.R., 2005. CYP2K6 from zebrafish (*Danio rerio*): Cloning, map-ping, developmental/tissue expression, and aflatoxin B1 activation by baculovirus expressed enzyme. Elsevier Inc, New York.

Wang, L., Yao, J.H., Chen, L., Chen, J.Z., Xue, J.L., Jia, W., 2007. Expression and possible functional roles of cytochromes P450 2J1 (zfCyp 2J1) in zebrafish. Biochemi. Biophys. Res. Comm. 352, 850-855.

Wedemeyer, G., 1968. Uptake and distribution of ZN 65 in the coho salmon egg (*Oncorhynchus kisutch*). Comp. Biochem. Physiol. 26, 271-279.

Weigt, S., Huebler, N., Braunbeck, T., von Landenberg, F., Broschard, T.H., 2010. Zebrafish teratogenicity test with metabolic activation (mDarT): Effects of phase I activation of acetaminophen on zebrafish *Danio rerio* embryos. Toxicology 275, 36-49.

Weigt, S., Huebler, N., Strecker, R., Braunbeck, T., Broschard, T.H., 2011. Zebrafish (*Danio rerio*) embryos as a model for testing proteratogens. Toxicology 281, 25-36.

Wells, P.G., Bhuller, Y., Chen, C.S., Jeng, W., Kasapinovic, S., Kennedy, J.C., Laposa, R.R., McCallum, G.P., Nicol, C.J., Parman, T., Wiley, M.J., Wong, A.W., 2004. Molecular and biochemical mechanisms in teratogenesis. Toxicol. Appl. Pharmacol. 197, 162-163.

Westerfield, M., 2007. The zebrafish book: a guide for the laboratory use of zebrafish (*Danio rerio*). Eugene: University of Oregon Press.

Weyers, A., Sokull-Klüttgen, B., Baraibar-Fentanes, J., Vollmer, G., 2000. Acute toxicity data: a comprehensive comparison of results of fish, *Daphnia* and algae tests with new substances notified in the EU. Environ. Toxicol. Chem. 19, 1931-1933.

Wu, H.H., Gao, C., Guo, Y.P., Zhang, Y.P., Zhang, J.Z., Ma, E.B., 2014. Acute toxicity and sublethal effects of fipronil on detoxification enzymes in juvenile zebrafish (*Danio rerio*). Pest. Biochem. Physiol. 115, 9-14.

Yamagami, K., Hamazaki, T.S., Yasumasu, S., Masuda, K., Iuchi, I., 1992. Molecular and cellular basis of formation, hardening, and breakdown of the egg envelope in fish. Int. Rev. Cytol. 136, 51-92.

Yang, Y., Ji, D., Huang, X., Zhang, J., Liu, J., 2017. Effects of metals on enantioselective toxicity and biotransformation of cis-bifenthrin in zebrafish. Environ. Toxicol. Chem. 36, 2139-2146.

Zhang, Q., Cheng, J.P., Xin, Q., 2015. Effects of tetracycline on developmental toxicity and molecular responses in zebrafish (*Danio rerio*) embryos. Ecotoxicology 24, 707-719.

Zhang, X.S., Rawson, D.M., 1996. Permeability of the vitelline membrane of zebrafish (*Brachydanio rerio*) embryos to methanol and propane-1,2-diol. Cryo-Letters 17, 273-280.

Zhu, B.T., 2010. On the general mechanism of selective induction of cytochrome p450 enzymes by chemicals: some theoretical considerations. Expert Opin. Drug Metab. Toxicol. 6, 483-494.