TEXTE

64/2014

From theory to reality – Evaluation of suitable organisms and test systems for the biomitoring of pharmaceuticals

Part I: Literature review



TEXTE 64/2014

Environmental Research of the Federal Ministry for the Environment, Nature Conservation, Building and Nuclear Safety

Project No. (FKZ) 3713 634 101 Report No. (UBA-FB) 001941/E

From theory to reality – Evaluation of suitable organisms and test systems for the biomonitoring of pharmaceuticals

Part I: Literature review

by

Rita Triebskorn Animal Physiological Ecology, University of Tübingen; Steinbeis Transfer Center for Ecotoxicology and Ecophysiology, Rottenburg

Simon Schwarz, Heinz-R. Köhler Animal Physiological Ecology, University of Tübingen

Kristin Berg, Dirk Jungmann GWT, Technical University Dresden

Manfred Frey Steinbeis Transfer Center for Applied Biological Chemistry, Mannheim

Jörg Oehlmann, Matthias Oetken Aquatic Ecotoxicology, University of Frankfurt

On behalf of the Federal Environment Agency (Germany)

Imprint

Publisher:

Umweltbundesamt Wörlitzer Platz 1 06844 Dessau-Roßlau Tel: +49 340-2103-0 Fax: +49 340-2103-2285 info@umweltbundesamt.de Internet: www.umweltbundesamt.de

f /umweltbundesamt.de
/umweltbundesamt

Study performed by:

Animal Physiological Ecology, University of Tübingen Konrad-Adenauer-Str. 20 72072 Tübingen, Germany

Study completed in:

November 2013

Edited by:

Section IV 2.2 Pharmaceuticals, Washing and Cleaning Agents Ina Ebert

Publication as pdf:

http://www.umweltbundesamt.de/publikationen/from-theory-to-reality-evaluation-of-suitable

ISSN 1862-4804

Dessau-Roßlau, August 2014

The Project underlying this report was supported with funding from the Federal Ministry for the Environment, Nature Conservation, Building and Nuclear safety under project number FKZ 3713 673 410 1. The responsibility for the content of this publication lies with the author(s).

Abstract

In order to identify sensitive organisms, biological endpoints *in vivo* and suitable *in vitro* test systems for the biomonitoring of pharmaceuticals, a two-part literature search was conducted. In the first part, the database *OEKOTOX* established by Bergmann et al. (2011) [1] was upgraded with effect data published between 2011 and 2013 for 90 pharmaceuticals of high priority. From all available data the lowest effect concentrations were identified and the most sensitive organisms and effect endpoints were determined. In addition, effect data were assessed with respect to their relation to measured environmental concentrations (MECs) in German surface waters by calculations of risk quotients MEC_{max}/LOEC_{min}. Publications providing the lowest effect data were evaluated with respect to their reliability according to the criteria of Wright-Walters et al. (2011) [2]. Out of 72 publications investigated for their reliability, 9 were "reliable", 49 "conditionally reliable" and 14 "not reliable" (of which 4 were, in part, conditionally reliable). Prior to the reliability check, the analgesics paracetamol, diclofenac and ibuprofen, the ßblocker propranolol, the antibiotics sulfamethoxazole and erythromycin and the lipid regulator gemfibrozil were identified as environmentally most relevant pharmaceuticals. The relevance was defined by MEC_{max}/LOEC_{min}-values >0.1 for a minimum of 3 different biota classes. After assessment of publication reliability, however, only for diclofenac MEC_{max}/LOEC_{min}-values >0.1 were found in more than 3 biota classes. As further important pharmaceuticals propranolol, sulfamethoxazole, bezafibrate, 17α -ethinlyestradiol, 17β -estradiol and oxytetracycline were identified with MEC_{max}/LOEC_{min}-values >0.1 for 2 biota classes.

Prior to the reliability assessment, the ciliate *Tetrahymena pyriformis*, zebrafish *(Danio rerio),* rainbow trout *(Oncorhynchus mykiss),* medaka *(Oryzias latipes),* the mussel *Elliptio complanata,* and the mudsnail *Potamopyrgus antipodarum* were identified as most sensitive organisms for pharmaceuticals. The most sensitive effect endpoints were chemotaxis, behavior, vitellogenin synthesis, growth rate, reproduction, histopathological alterations, molecular stress biomarkers, oxidative stress markers, receptor binding, and gene expression. All extremely low LOECs (in the range of pg/L) for chemotaxis alterations in *Tetrahymena pyriformis*, however, derive from only a single publication by Lang & Köhidai (2012) [3], the reliability of which, however, could not be proven.

In the second part, a literature search was conducted with the aim to evaluate the state of knowledge in the field of *in vitro* testing of pharmaceutical effects, to discover advantages and disadvantages of biochemical and cell-based assays and to suggest possibilities for the realisation of cell-based tools for a mode of action-based biomonitoring of pharmaceuticals. Promising approaches for β-adrenoreceptor blockers and cyclooxygenase-inhibitors (analgesics) are presented.

As a future perspective for a biomonitoring of pharmaceuticals, the development of mode of action-based *in vitro* test systems for β-blockers and NSAIDs are recommended. Their suitability to reflect *in vivo* responses of organisms which have been identified to be sensitive in part 1 of this study, or which are relevant for German aquatic ecosystems should be assessed, and they should be validated by *in vivo* studies to be conducted in parallel to their establishment.

Kurzbeschreibung

Mit dem Ziel, sensitive Organismen und organismische Endpunkte sowie geeignete *in vitro*-Testsysteme für ein Biomonitoring von Arzneimitteln zu identifizieren, wurde eine zweiteilige Literaturstudie durchgeführt.

Im ersten Teil wurde die von Bergmann et al. (2011) [1] erstellte Datenbank OEKOTOX um Wirkdaten für 90 Arzneimittel von hoher Priorität bis 2013 erweitert. Von allen verfügbaren Daten wurden die niedrigsten Effektwerte identifiziert und die sensitivsten Organismen bzw. Effektendpunkte bestimmt. Darüber hinaus wurden die Effektkonzentrationen zu gemessenen Umweltkonzentrationen in Deutschen Oberflächengewässern (MECs) in Beziehung gestellt und Risikoquotienten MEC_{max}/LOEC_{min} berechnet. Diejenigen Publikationen, in denen die empfindlichsten Endpunkte bzw. Organismen enthalten waren, wurden auf der Basis der von Wright-Walters et al. (2011) [2] genannten Kriterien auf Reliabilität geprüft. Von 72 auf Reliabilität überprüften Publikationen wurden 9 als "reliabel", 49 als "bedingt reliabel" und 14 als "nicht reliabel" (von denen 4 in Teilen bedingt reliabel waren) klassifiziert. Vor der Reliabilitätsprüfung wurden in der vorliegenden Literaturstudie als Pharmazeutika mit höchster Umweltrelevanz die Schmerzmittel Paracetamol, Diclofenac und Ibuprofen, der ß-Blocker Propranolol, die Antibiotika Sulfamethoxazol und Erythromycin und der Lipidsenker Gemfibrozil identifiziert. Die Umweltrelevanz wurde auf der Basis eines Risikoquotienten (MEC_{max}/LOEC_{min}>0.1 für mindestens 3 Organismengruppen) definiert. Nach erfolgter Reliabilitätsprüfung zeigten sich MEC_{max}/LOEC_{min}-Werte >0.1 bei mehr als 3 Organismengruppen jedoch nur noch für Diclofenac. Als weitere wichtige Pharmazeutika wurden Propranolol, Sulfamethoxazol, Bezafibrat, 17α-Ethinlyestradiol, 17β-Estradiol und Oxytetrazyclin mit MEC_{max}/LOEC_{min}-Werten >0.1 bei 2 Organismengruppen identifiziert. Als sensitivste Organismen für Arzneimittel hatten sich vor der Reliabilitätsprüfung der Ciliat Tetrahymena pyriformis, der Zebrabärbling (Danio rerio), die Regenbogenforelle (Oncorhynchus mykiss), Medaka (Oryzias latipes), die Muschel Elliptio complanata und die Zwergdeckelschnecke Potamopyrgus antipodarum erwiesen. Die sensitivsten Effektendpunkte waren Chemotaxis, Verhalten, Vitellogeninsynthese, Wachstum, Reproduktion, histopathologische Veränderungen, molekulare und oxidative Stressmarker, Genexpression und Rezeptorbindung. Alle extrem niedrigen LOECs (im Bereich von pg/L) für die Beeinflussung der Chemotaxis bei *Tetrahymena pyriformis* stammen allerdings von einer einzigen Publikation von Lang & Köhidai (2012) [3], die sich in der durchgeführten Reliabilitätsprüfung als "nicht reliabel" erwies, so dass der Zebrabärbling als bedeutendster Testorganismus und das Verhalten als sensitivster Wirkendpunkt nachrückt.

Im zweiten Teil der Studie wurde eine Literaturrecherche durchgeführt, die zum Ziel hatte, den Wissensstand im Bereich von *in vitro*-Verfahren zum Nachweis von Arzneimittel-Effekten zu beleuchten, Vor- und Nachteile von biochemischen und zellbasierten Assays herauszuarbeiten und Möglichkeiten für die Realisierung von *in vitro* Testsystemen für ein Wirkmechanismusbasiertes Biomonitoring von Arzneimitteln vorzuschlagen. Am Ende dieses Teils werden vielversprechende Wege zur Etablierung solcher Testsysteme für den Nachweis der Wirkungen von ß-Blockern und Cyclooxygenase-Hemmern (Schmerzmitteln) dargestellt.

Für ein künftiges Biomonitoring von Arzneimitteln wird empfohlen, Wirkstoffklassenspezifische *in vitro*-Testsysteme für ß-Blocker und Schmerzmittel zu entwickeln. Durch parallel durchzuführende *in vivo* Untersuchungen soll überprüft werden, inwieweit diese Testsysteme geeignet sind, *in vivo*-Reaktionen von Organismen abzubilden, die in Teil 1 der vorliegenden Studie als sensitiv für Arzneimittel identifiziert wurden, oder die als relevant für deutsche Fließgewässer bekannt sind.

List of Contents

1 Introduction

2 Materials and Methods

- 2.1 Literature search and data collection
- 2.2 Created library and data files
 - 2.2.1 Endnote library
 - 2.2.2 OEKOTOX_{upgraded}
 - 2.3.3 "Evaluation database" for assessment of data

3 Results

3.1 Effects of pharmaceuticals in vivo

- 3.1.1 Number of publications and database entries
- 3.1.2 Number of publications per organism group
- 3.1.3 Evaluation of data
- 3.1.3.1 Most sensitive biota classes for pharmaceuticals
- 3.1.3.2 Most sensitive effect endpoints per substance classes
- 3.1.3.3 Identification of pharmaceutical of highest relevance
- 3.1.3.4 Data for mixtures
- 3.1.3.5 Data with population relevance
- 3.1.4 Reliability of publications
- 3.1.5 Consequences of reliability assessment for the described results
- 3.1.6 Summary of part 1

3.2 In vitro test systems

- 3.2.1 Introduction
- 3.2.2 Fluorescence Spectroscopy
 - 3.2.2.1 Steady-State and Time-Resolved Fluorescence
 - 3.2.2.3 Fluorescence Polarization (FP)
 - 3.2.2.3 Fluorescence Resonance Energy Transfer (FRET)
- 3.2.3 Fluorescence Sensing
 - 3.2.3.1 Low molecular weight Sensors

3.2.3.2 Protein Sensors

- 3.2.3.3 Green fluorescent protein (GFP) variants
 - 3.2.3.3.1 Single Fluorophore Sensors
 - 3.2.3.3.1.1 Halide and pH Sensors
 - 3.2.3.3.1.2 Redox Potential Sensors
 - 3.2.3.3.1.3 Calcium Sensors
 - 3.2.3.3.1.4 cGMP Sensors
 - 3.2.3.3.1.5 Reactive Oxygen Species Sensor
 - 3.2.3.3.1.6 ATP/ADP ratio sensing
 - 3.2.3.3.1.7 Membrane Potential sensing
 - 3.2.3.3.2 FRET / BRET based Sensors
 - 3.2.3.3.2.1 FRET based cAMP Sensor
 - 3.2.3.3.2.2 BRET based cAMP Sensor
 - 3.2.3.3.2.3 FRET based GPCR activation Sensor
 - 3.2.3.3.2.4 FRET based protein kinase activity Sensor
 - 3.2.3.3.2.5 Coupled FRET based NO Sensor
 - 3.2.3.3.3 Protein Fragment Complementation Sensor
- 3.2.4 Immunoassays
 - 3.2.4.1 cAMP AlphaScreen® (PerkinElmer)
 - 3.2.4.2 cAMP LANCE [®] (PerkinElmer)
 - 3.2.4.3 cAMP HTRF (CisBio International)
- 3.2.4.4 cAMP Screen (Life Technologies)
- 3.2.5 Reporter Assays
- 3.2.6 β -adrenoceptor assays
 - 3.2.6.1 Tagged ligand binding assays
 - 3.2.6.2 cAMP luminescent biosensor assay
- 3.2.6.3 cAMP CRE reporter assays
- 3.2.6.4 Receptor conformation based FRET assay
- 3.2.6.5 cAMP gated ion channel
- 3.2.6.6 HCN2 cAMP FRET assay
- 3.2.7 Cyclooxygenase assays
 - 3.2.7.1 Oxygen consumption assay
 - 3.2.7.2 Peroxidase co-substrate oxidation assay
 - 3.2.7.3 COX immunoassay
 - 3.2.7.4 COX fluorescence assay

3.2.8 Sensitivity of in vitro and in vivo assays

3.2.8.1 Comparative in vivo | in vitro analysis of endocrine disrupting chemicals

3.2.9 Evaluation

3.2.9.1 Selection of substances with high priority

3.2.9.2 Evaluation of published in vitro assays

3.2.9.2.1 β-adrenoceptor compatible *in vitro* assays

3.2.9.2.2 Cyclooxygenase compatible *in vitro* assays

3.2.9.3 Suggestion for in vitro assay development

3.2.9.3.1 β-adrenoceptor *in vitro* assay suggestion

3.2.9.3.2 Cyclooxygenase in vitro assays suggestion

4 Conclusions

5 Identification of shortcomings

6 Future perspectives in biomonitoring of pharmaceuticals

7 References

Part 1 Part 2 Reviewed literature (only relevant articles)

List of Figures

Part 1

Figure 1:	Number of publications per substance class
Figure 2:	Number of database entries per substance class
Figure 3:	Number of studies per biota group
Figure 4:	Number of database entries per biota group
Figure 5:	Number of lowest plus second lowest effect values for 13 pharmaceutical classes
Figure 6:	Number of pharmaceuticals with defined LOECs for investigated species
Figure 7:	Number of pharmaceuticals with defined LOECs for investigated effect endpoints
Figure 8:	Number of pharmaceuticals per pharmaceutical class with MECmax/LOECmin values ${\succ}0.1$
Figure 9:	Number of pharmaceuticals with defined $\ensuremath{\text{MEC}_{\text{max}}}\xspace/\ensuremath{\text{LOEC}_{\text{min}}}$ values per organism group
Figure 10:	Number of database entries with population-relevant endpoints

Part 2

Figure 1:	Fluorescence Resonance Energy Transfer (FRET)
Figure 2:	Spectral observables for fluorescence sensing
Figure 3:	Surface bound maltose sensor
Figure 4:	Molecular switch
Figure 5:	AlphaScreen cAMP assay
Figure 6:	LANCE cAMP assay
Figure 7:	HTRF cAMP assay
Figure 8:	Receptor binding and G-protein-dependent reporter gene assays
Figure 9:	cAMP luciferase assay
Figure 10:	Development of a β -1 adrenoceptor FRET sensor
Figure 11:	COX fluorescence assay
Figure 12:	Modulation of redox regulation
Figure 13:	Arachidonic acid metabolism through the COX pathway

List of Tables

Part 1

Table 1:	List of pharmaceuticals with defined priority classes.
Table 2:	Pharmaceuticals for which $MEC_{max}/LOEC_{min}$ values were >0.1 in 1, 2, 3 or 4 biota groups
Table 3:	Comparison of pharmaceutical effect concentrations applied either isolated or mixture of pharmaceuticals
Table 4:	Publications proved for reliability, reliability criteria met, and decision on reliability
Table 5:	Further information on pharmaceuticals tested, test organisms, effect endpoints and reasons for lacking reliability for not reliable studies
Table 6:	Assessment of data for most sensitive species with respect to reliability of publications
Table 7:	Priority pharmaceuticals before reliability evaluation, only based on data from $OEKOTOX_{upgrade}$
Table 8:	Priority pharmaceuticals after reliability evaluation, only based on data from $OEKOTOX_{upgrade}$

Part 2

Table 1:	β1-adrenoceptor biosensors
Table 2:	Cyclooxygenase activity assays

List of Abbreviations

Part 1

CAS	Chemical Abstracts Service
DOI	Digital Object Identifier
LOEC	Lowest observed effect concentration
LOEC min	Minimum of lowest observed effect concentrations reported
LogK _{ow}	Octanol/water partitioning coefficient (logarithmic form)
MEC	Measured environmental concentration
MEC_{max}	Maximum of measured environmental concentrations reported
MOA	Mode of action
NOEC	No observed effect concentration
NSAID	Non-steroidal anti-inflammatory drug
PBT	persistent, bioaccumulating, and toxic
PEC	Predicted environmental concentration
PNEC	Predicted no effect concentration
TER	Toxicity exposure ratio
UBA	Umweltbundesamt der Bundesrepublik Deutschland

Part 2

A. victoria	Aequorea victoria
ADP	Adenosinediphosphate
ALPHA	Amplified Luminescent Proximity Homogeneous Assay
AMP	Adenosine monophosphate
AP	Alkaline phospatase
ATP	Adenosinetriphosphate
b-AR	b-adrenoceptor
BRET	Bioluminescence resonance energy transfer
C.elegans	Caenorhabditis elegans
CaM	Calmodulin
cAMP	Cyclic adenosinemonophosphate
CDCFDA	5', 6' Dicarboxy-2',7'-dichlorodihydrofluorescein
CDCFDA	2′,7′-dichlorodihydrofluorescein
CFP	Cyan fluorescent protein
cGMP	Cyclic guanosinemonophosphate

CNG	Cyclic nucleotide-gated
COX	Cyclooxygenase
CRE	Cyclic adenosinemonophosphate response element
CREB	cAMP response element binding protein
E.coli	Escherichia coli
EC ₅₀	half-maximal effective concentration
EGFP	Enhanced green fluorescent protein
Ерас	Exchange protein directly activated by cyclic adenosinemonophosphate
EYFP	Enhanced yellow fluorescent protein
FlincGs	fluorescent indicators of cyclic guanosinemonophosphate
FP	Fluorescence polarization
FRET	Fluorescence Resonance Energy Transfer or Förster Resonance Energy Transfer
G12/13	RhoA signalling G proteins
GCa-MPs	M13 domain of a myosin light chain kinase and calmodulin based genetically encoded calcium indicator
GFP	Green fluorescent protein
Gi	inhibitory G proteins
GlnK1	nitrogen regulatory protein P-II
GPCR	G protein-coupled receptor
Gq	Phospholipase C signalling G proteins
Gs	cAMP signalling G proteins
HCN2	Cyclic nucleotide-gated channel 2
HEK293	Human Embryonic Kidney 293 cells
HTRF	Homogeneous time resolved fluorescence
ICUE	Indicator of cAMP using Epac
LANCE	Lanthanide chelate exite
MBP	Maltose binding protein
NAD	Nicotinamide adenine dinucleotide
NFAT-RE	Nuclear factor of activated T-cells response element
NF-ĸB	Nuclear Factor kappaB
NO	Nitrogen oxide
РКА	Phospho kinase A
РКС	Phospho kinase C
RhoA	Ras homolog gene family member A
roGFPs	Redoxsensitive fluorescence protein indicators

rxYFPs	Redox yellow fluorescence protein sensors
SPA	Scintillation proximity assay
SRE	Serum response element
SRF-RE	serum response factor response element
t1/2	Half-life
UBA	Umweltbundesamt der Bundesrepublik Deutschland
YFP	Yellow fluorescent protein

1 Introduction

Chemical analysis has regularly revealed the presence of human and veterinary pharmaceuticals in wastewater and surface water in nanogram to microgram per liter concentrations (e.g. Fent et al., 2006 [4], Brauch, 2011 [5]). Decisions on the environmental relevance of these substances usually rely on data recorded for current and future consumption rates, environmental concentrations, environmental fate and pathways (persistence), and ecotoxicological effects recorded in laboratory studies. The problem with this approach is the fact that ecotoxicological routine testing mainly focuses on acute or chronic unspecific toxicity, whereas pharmaceuticals, according to their envisaged specific action in man (human medicine) or companion animals (veterinary drugs) more likely exert specific effects based on their mode of action (MOA) (Brausch et al. 2012 [6]). Information on the effects of active pharmaceutical ingredient classes, either acting on targets or exerting unwanted adverse side effects are, therefore, mandatory prerequisites for an effect-directed monitoring of pharmaceuticals in aquatic ecosystems.

One focus of the present literature review study thus was on the question which organisms sensitively respond to pharmaceuticals and which sublethal parameters can be used as warning sentinels (biomarkers) to monitor action of pharmaceuticals in a sensitive and specific way. In a comprehensive report of Bergmann et al. (2011) [1] a prioritization of human and veterinary pharmaceuticals was undertaken on the basis of data on the presence of compounds in the aquatic environment, their ecotoxicological effects, and their consumption rates up to the year 2011. This report lists analytical data for 274 ingredients and ecotoxicological effect data for 251 ingredients, all of them incorporated in both the MEC (Measured Environmental Concentrations) and ÖKOTOX (effect data) databases. Bergmann et al. (2011) [1] have outlined 24 substances with high priority for environmental monitoring programs. However, the databases also provide evidence that 31 pharmaceuticals with partially high and steadily increasing consumption rates lacked ecotoxicological data until 2009 resulting in a very high uncertainty regarding the assessment of their environmental relevance. Supplementary to the substances regarded as relevant for further research by Bergmann et al. (2011) [1], the German Federal Ministry for the Environment, Nature Conservation and Nuclear Safety has identified further substances as to be of high priority for an evaluation with respect to possible environmental effects. In the present study, we therefore focused on a set of 90 substances, prioritized by Bergmann et al. (2011) [1] and UBA, for which literature was analyzed.

Bergmann et al. (2011) [1] concluded from their research that already the number of substances with high priority (24) would exceed the capacity of routine chemical monitoring. They therefore suggested having leading substances defined for particular ingredient classes that should be included in monitoring programs. The apparent but crucial drawback of such an approach, however, lies in the risk of overlooking effective concentrations of non-leading substances. In contrast to it, the use of tests that could visualize specific molecular interactions of chemicals exhibiting the same mode of action may enable a monitoring of an entire class of compounds. Particularly in view of the vast number of pharmaceuticals with increasing consumption rates but lacking ecotoxicological effect data, a mode of-action-specific *in vitro* assay is a matter of paramount interest; also because new and future compounds that act in the same way on the same molecular target can easily be traced, once such an assay has been established.

In vitro systems using highly sensitive fluorescence detection technologies are already used by the pharmaceutical industry to identify compound classes as promising candidates in the

development of new pharmaceuticals (Eggeling et al., 2003) [7]. A biomonitoring on the basis of such specific mechanisms of action could be possible for e.g. non-steroidal anti-inflammatory drugs (NSAIDs) or β -blockers. For these, the inhibition of the cyclooxygenase and the blocking of β -receptors could be used as mode of actions to be targeted. In contrast to Escher et al. (2005) [8] who use "*in vitro* Assessment of Modes of Toxic Action" in that sense that specific mode of actions defined by the test system itself are investigated (e.g. photosynthesis inhibition of pharmaceuticals by a photosynthesis inhibition assay), we thus favor an approach to use the specific target molecules for pharmaceutical classes (responsible for the pharmacological effect of the respective substance class) as monitoring tools. In the second part of this study, we therefore evaluated different *in vitro* test systems with respect to their suitability as future test systems in routine monitoring of pharmaceutical classes.

In summary, this study reviews (1) the current state of knowledge in effect-directed research and analyzes present data with focus on suitable compounds, organisms, and endpoints which may be combined in a monitoring approach, and (2) in addition, the state of knowledge for *in vitro* testing of pharmaceuticals as a prerequisite for the development of mode of-action-based monitoring tools.

2 Materials and Methods

2.1 Literature search and data collection

The literature search is divided into two parts.

In the first part, effect data for 90 pharmaceuticals defined as relevant according to either Bergmann et al. (2011) [1] or UBA were collected for several biota classes (bacteria, protists, plants/algae, mollusks, crustaceans, insects, fish, tetrapods, others incl. communities). The list of pharmaceuticals and the priority classes are shown in table 1. The prioritization of UBA is based on MECs obtained from the German counties, consumption rates (tendencies between 2002 and 2012), effect values of UBA-internal studies or literature, suspicion to be a PBT compound, degradability and metabolism. The prioritization of Bergmann et al. (2011) [1] is only based on literature data which resulted in partial differences between the two lists of priority substances available.

The search was restricted to literature published between 2011 and 2013, since all earlier published data have been analyzed by Bergmann et al. (2011) [1]. For the search, Web of Knowledge (including the databases "Web of Science[®]", "BIOSIS Citation IndexSM", "BIOSIS Previews[®]", "MEDLINE[®]" and "Journal Citation Reports[®]") were used. The following search items were defined:

For Reviews: pharma*AND eco*, limited by "review"

For single substances: the specific search entry for each pharmaceutical (see table 1) combined with the search terms: $aqua^* OR \ eco^* OR \ tox^*$

The following research areas were seen as irrelevant and therefore excluded from the literature search:

Anesthesiology, Anthropology, Biomedical, Social Sciences, Biophysics, Business Economics, Communication, Computer Science, Criminology Penology, Critical Care Medicine, Cultural Studies, Demography, Dentist, Oral Surgery Medicine, Dermatology, Education, Educational Research, Electrochemistry, Emergency Medicine, Energy Fuels, Engineering, Ethnic Studies, Food Science Technology, General Internal Medicine, Genetics Heredity, Geography, Geology, Geriatrics, Gerontology, Government, Law, Health Care Sciences, Services, History, Imaging Science, Photographic Technology, Infectious Diseases, Information Science, Library Science, Instruments Instrumentation, Integrative Complementary Medicine, International Relations, Legal Medicine, Materials Science, Mathematical Computational Biology, Mathematics, Medical Informatics, Medical Laboratory Technology, Oncology, Ophthalmology, Orthopedics, Otorhinolaryngology, Parasitology, Pediatrics, Pharmacology, Pharmacy, Physics, Polymer Science, Psychiatry, Psychology, Radiology, Nuclear Medicine, Medical Imaging, Rehabilitation, Research Experimental Medicine, Respiratory System, Social Issues, Sociology, Sport Sciences, Substance Abuse, Surgery, Transplantation, Tropical Medicine, Urology, Nephrology, Virology

In the second part, literature searches were conducted in PubMed using the following search items:

High throughput screening (title) & review (publication type) HTS technolog* & review (publication type) HTS technolog* (title) fluorescence based techni* cell based assay* & review (publication type) cell based reporter* (title) cell based screening & review (publication type) & cell based (title) fluorescent biosensor* & review (publication type) fluorescent protein* & review (publication type) & reporter* signalling dynamics in living cell* & review (publication type)

Data published between 1995 and 2013 were analyzed.

Table 1: List of pharmaceuticals with defined priority classes. The prioritization of UBA is based on MECs obtained from the German counties, consumption rates (tendencies between 2002 and 2012), effect values of UBA-internal studies or literature, suspicion to be a PBT compound, degradability and metabolism. The prioritization of Bergmann et al. (2011) [1] is only based on literature data which resulted in partial differences between the two lists of priority substances available.

- P: high priority according to Bergmann et al. (2011) [1] and UBA;
- Pu: high priority according to UBA
- Pb: high priority according to Bergmann et al. (2011) [1]
- (P): medium priority according to Bergmann et al. (2011) [1]
- ?: substances requiring further information
- none: substances without priority according to Bergmann et al. (2011) [1]

Pharmaceutical	Search entry	Priority class	Priority class (Bergmann)	Priority class	Class	CAS
Sartanic acid						
	sartan*	Р		Pu	antihypertensive	
14-						
Hydroxyclarithromycin	hydroxyclarithromycin*	Р		Pu	antibiotic	110671-78-8
17alpha-Ethinylestradiol	ethinyl* estradiol*	Р	Р	Р	contraceptive	57-63-6
17beta-Estradiol	estradiol* AND pharma*					
			Р	Pb	hormone	50-28-2
4-N-						
Methylaminoantipyrin	methylaminoantipyrin*	Р		Pu	analgesic	519-98-2

Biomonitoring of pharmaceuticals/final report

Pharmaceutical	Search entry	Priority class	Priority class	Priority class	Class	CAS
6(carboxymethoxy)-4- (2-chlorophenyl)-5- (ethoxycarbonyl)-2- methylpyridine-3- carboxylic acid	amlodipin*	P		Pu	antihypertensive	
Acetylsalicylic acid	acetylsalic*				NSAID	50-78-2
Aciclovir	aciclovir*		?	?	antiviral drug	59277-89-3
Allopurinol	allopurinol*				xanthine oxidase inhibitor	315-30-0
AMDOPH	AMDOPH		(P)	(P)	analgesic	
Amlodipine	amlodipin*	Р		Pu	antihypertensive	88150-42-9
Amoxicilline	amoxicillin*	Р	Р	Р	antibiotic	26787-78-0
Atenolol/ Atenolol acid	atenolol*, tenormin*	Р		Pu	betablocker	29122-68-7
Azitromycin	azitromycin*, zithromax*, azithrocin*, azin*	Р		Pu	antibiotic	83905-01-5
Bezafibrate	bezafibrat*		(P)	(P)	lipid regulator	41859-67-0
Bisoprolol	bisoprolol*	Р	?	Pu	betablocker	66722-44-9
Capecitabin	capectiabin*, xeloda*	Р		Pu	cytostatic	15361-50-9
Carbamazepine	carbamazepin*	Р	Р	Р	anticonvulsant	298-46-4
Cefaclor	cefaclor* OR cefachlor*		?	?	antibiotic	53994-73-3
Cefuroxime axetil	cefuroxim* axetil*		?	?	antibiotic	64544-07-6
Chloramphenicole	chloramphenicol*		Р	Pb	antibiotic	56-75-7
Chlortetracycline	chlortetracyclin*		Р	Pb	antibiotic	57-62-5
Ciprofloxacin	ciprofloxacin*	Р	Р	Р	antibiotic	85721-33-1
Clarithromycine	clarithromycin*	Р	Р	Р	antibiotic	81103-11-9
Clindamycin	clindamycin*	Р	?	Pu	antibiotic	18323-44-9
Clopidogrel	clopidogrel*		?	?	antiplatelet agent	113665-84-2
Clotrimazole	clotrimazol*		(P)	(P)	antimycotic	23593-75-1
Deltamethrin	deltamethrin*, decamethrin*	Р		Pu	antiparasitics	52918-63-5
Desflurane	desfluran*		?	?	anesthetic	57041-67-5
Diatrizoic acid	diatriz* OR iotalam* OR amidotriz*		(P)	(P)	contrast agent	737-31-5
Diazepam	diazepam*		(P)	(P)	antidepressant/ antipsychotic	439-14-5
Diclofenac	diclofenac*	Р	Р	Р	NSAID	15307-86-5
Dienogest	dienogest*, visanne*	Р		Pu	gestagan	65928-58-7
Dipyridamole	dipyridamol*		?	?	antiplatelet agent	58-32-2
Doxycycline	doxycyclin*		Р	Pb	antibiotic	564-25-0
Duloxetine	duloxetin*, cymbalta*	Р		Pu	antidepressant/ antipsychotic	116539-59-4
Entacapone	entacapon*		?	?	catecholamin-O- Methyltransferase- Inhibitor	130929-57-6
Eprosartan	eprosartan*		?	?	antihypertensive	133040-01-4
Erythromycin	erythromycin*		P	Pb	antibiotic	114-07-8
Gabapentin	gabapentin* OR neurotin*		?	?	anticonvulsant	60142-96-3
Gemfibrozil	gemfibrozil*, lopid*	Р		Pu	lipid regulator	25812-30-0
Glutaral	glutaral*		?	?	desinfectant (Glutaraldehyd)	111-30-8
Hydrochlorothiazide	hydrochlorothiazid*		?	?	antihypertensive	58-93-5
Ibuprofen	ibuprofen*	Р	(P)	Р	NSAID	15687-27-1
Imatinib	imatinib*, gleevec*, glivec*	Р		Pu	cytostatic	152459-95-5
Indometacin	indometacin* OR indomethacin*		(P)	(P)	NSAID	53-86-1
lohexol	iohexol* OR omnipaq*	Р	(P)	Р	contrast agent	66108-95-0
lomeprol	iomeprol*	Р	(P)	Р	contrast agent	78649-41-9
lopamidol	iopamidol*		(P)	(P)	contrast agent	60166-93-0

Biomonitoring of pharmaceuticals/final report

Pharmaceutical	Search entry	Priority class	Priority class (Bergmann)	Priority class (final)	Class	CAS
lopromide	iopromid* OR ultravist*		(P)	(P)	contrast agent	73334-07-3
Ivermectin	ivermectin*, stromectol*	Р		Р	antiparasitic	70288-86-7
Lamotrigin	lamotrigin*, lamictal*	Р		Pu	anticonvulsant	84057-84-1
Levetiracetam	levetiracetam*		?	?	anticonvulsant	102767-28-2
Lincomycin	lincomycin*		(P)	(P)	antibiotic	154-21-2
Mesalazine	mesalazin*		?	?	NSAID	89-57-6
Mestranol	mestranol*		Р	Pb	contraceptive	72-33-3
Metamizole	metamizol*	Р		Pu	analgesic	68-89-3
Metformin	metformin*	Р		Pu	antidiabetic	657-24-9
Metoprolol	metoprolol*	Р		Pu	betablocker	51384-51-1
Naproxen	naproxen*	Р	Р	Р	NSAID	22204-53-1
Norethisterone	norethisteron*		Р	Pb	contraceptive	68-22-4
Opipramol	opipramol*		?	?	antidepressant/ antipsychotic	315-72-0
Oxytetracycline	oxytetracyclin*		Р	Pb	antibiotic	79-57-2
Pantoprazole	pantoprazol*		?	?	proton pump inhibitor	102625-70-7
Paracetamol	paracetamol* OR acetaminophen*		Р	Pb	analgesic	103-90-2
Piperacillin	piperacillin*		?	?	antibiotic	61477-96-1
Pregabalin	pregabalin*		?	?	anticonvulsant	148553-50-8
Primidone	primidon*		Р	Pb	anticonvulsant	125-33-7
Propranolol	propanolol* OR propranolol*		(P)	(P)	betablocker	525-66-6
Quetiapine	quetiapin*	Р	?	Pu	antidepressant/ antipsychotic	11974-69-7
Ramipril	ramipril*		?	?	antihypertensive	87333-19-5
Roxithromycine	roxithromycin*		Р	Pb	antibiotic	80214-83-1
Sevelamer	sevelamer*		?	?	phosphate binding drug	52757-95-6
Simvastatin	simvastatin*	Р	?	Pu	lipid regulator	79902-63-9
Strontium ranelate	strontium ranelat*		?	?	osteoporose agent	135459-90-4
Sulbactam	sulbactam*		?	?	antibiotic	68373-14-8
Sulfadimethoxine	sulfadimethoxin*		Р	Pb	antibiotic	122-11-2
Sulfadimidine	sulfadimidin* OR sulfamethazin*		Р	Pb	antibiotic	57-68-1
Sulfamethoxazole	sulfamethoxazol*	Р	Р	Р	antibiotic	723-46-6
Tazobactam	tazobactam*		?	?	antibiotic	89786-04-9
Telmisartan	telmisartan*		?	?	antihypertensive	144701-48-4
Tetracycline	tetracyclin*		Р	Pb	antibiotic	60-54-8
Tiamulin	tiamulin*	Р	Р	Р	antibiotic	55297-95-5
Tilidine	tilidin*		?	?	analgesic	51931-66-9
Torasemide	torasemid* OR torsemid*		?	?	diuretic drug	56211-40-6
Tramadol	tramadol*, ryzolt*, ultram*	Р		Pu	analgesic	27203-92-5
Valproic acid	valpro*		?	?	anticonvulsant	99-66-1
Valsartan	valsartan* OR angiotan* OR diovan*	Р	?	Pu	antihypertensive	137862-53-4
Venlafaxine	venlafaxin*		?	?	antidepressant/ antipsychotic	93413-69-5

2.2. Created library and data files

2.2.1 Endnote library

All references analyzed were included into an Endnote library ("EndNote Library -

Pharmaceuticals – Literature study part 1 / 2 /3") in the format ".CIW".

The three partners involved in the literature search used the following labels for their citations: BER 1-x: GWT Dresden (R-: Review, A-: Additional Information, I-: Irrelevant)

SCH 1-x: University Tübingen (R-: Review, A-: Additional Information, I-: Irrelevant)

FRE 1-x: STZ Frey

The library contains the following folders:

Part 1:

Tuebingen - relevant studies: Studies on vertebrates, plants, protozoans and bacteria, which were directly included into the database.

- Tuebingen additional information: Studies, mainly reviews, on vertebrates, plants, protozoans and bacteria, which were used as help for the general interpretation of the data, but not directly included into the database.
- Tuebingen irrelevant studies: Studies, which were analyzed but were not included into the database.
- Tuebingen- evaluation not possible: Studies, for which an evaluation was not possible because the full text was not available and the abstract did not contain sufficient information.
- Dresden relevant studies: Studies on invertebrates, which were directly included into the database.
- Dresden additional information: Studies, mainly reviews, on invertebrates, which were used as help for the general interpretation of the data, but not directly included into the database.
- Dresden irrelevant studies: Studies, which were analyzed but were not included into the database.

Part 2:

Mannheim - *in vitro* techniques: Studies on *in vitro* techniques, which were used for the second part of the literature study.

Part 3:

Additional literature used for this report

2.2.2 OEKOTOX_{upgrade}

In part 1 of the study, all publications available for the 90 substances were analyzed and evaluated with respect to the following criteria:

Name of effective substance, investigated form of the substance (salt/conjugate), synonyms, CAS number, LogK_{ow}, molar mass, substance class, field of application, effective substance or medical product, single substance or mixture, test organisms (species), biota group, field/lab/mesocosm test, effect endpoint, population relevance yes/no, standard method yes/no, test concentration in water (µg/L, mol/L), test concentration in sediment (if necessary), applied dose (if necessary), duration of test, acute/chronic/sub-acute, chemical analyses present, accumulation data present, bioaccumulation factor, NOEC (µg/L, µmol/L), LOEC (µg/L, µmol/L),

 EC_{10} (µg/L, µmol/L), EC_{50} (µg/L, µmol/L), citation, DOI, full text available, Endnote label, chemical present in priority list yes/no, comments, reliability analysis (only for selected publications). In case no effect was observed in the study, the highest concentration tested (without effect) has been included as "NOEC" without corresponding LOEC. Furthermore, numerous studies reported effects already at the lowest tested concentration. These values are recorded as LOECs without a corresponding NOEC. It should be kept in mind that there is a possibility that lower values than the reported may also cause an effect.

Besides data for the priority substances defined in table 1, data for several other pharmaceuticals is also reported in the database. These effects were reported in the analyzed publications besides the ones for the target substances, and therefore included. However, the data sets for these substances are not complete, since they are mere "byproducts" of the original search.

All data were included into the database OEKOTOX_{upgrade}

A separate row was created for every chemical, species and endpoint used in each study. Each single row was defined as a "database entry".

Because many publications report on multiple chemicals, organisms or endpoints, the total number of database entries is much higher than the number of publications.

2.2.3. "Evaluation database" for assessment of data

With the aim to evaluate the collected data with respect to (1) their suitability for the study, (2) data number per pharmaceutical and organism group, (3) most sensitive organisms, (4) lowest effect concentrations, (5) mixture toxicity, and (6) population relevance, the "evaluation database" has been created.

In data sheet (4) the lowest and second lowest effect data and measured environmental concentrations (MECs) obtained either from the MEC database of Bergmann et al. (2011) [1] or UBA were included with the aim to assess the environmental relevance of the lowest effect data. However, not for all 90 substances both values MECs and LOECs were available. As a result, only for 32 substances a risk quotient (MEC_{max}/LOEC_{min}) could be calculated as a proxy for their environmental relevance. These substances were: paracetamol, tramadol, amoxicilline, chloramphenicole, chlortetracycline, ciprofloxacin, doxycycline, erythromycin, lincomycin, oxytetracycline, sulfadimethoxine, sulfadimidine, sulfamethoxazole, carbamazepine, diazepam, clotrimazole, ivermectin, atenolol/ atenolol acid, metoprolol, propranolol, 17alpha-ethinylestradiol, 17beta-estradiol, norethisterone, diatrizoic acid, bezafibrate, gemfibrozil, acetyl cysteine, acetylsalicylic acid, diclofenac, ibuprofen, indometacin, naproxen. For tetracycline, quetiapine, venlafaxine and deltamethrin no MECs were available. With the aim to compare mixture toxicity data with LOECs for isolated substances, the "mixture toxicity" datasheet was created. The data sheet "population relevance" has been used as a basis to quantify available data with relevance for the population level.

3 Results

3.1. Effects of pharmaceuticals in vivo

3.1.1 Number of publications and database entries

Altogether, 452 publications were analyzed for 90 pharmaceuticals. 325 papers contained data for vertebrates, plants/algae, protozoa, or bacteria (analyzed by the University of Tübingen), 179 papers contained data for invertebrates (analyzed by the GWT, TU Dresden) , and 50 of them contained data for invertebrates and either vertebrates, plants, protozoa or bacteria (analyzed by both, University of Tübingen and GWT TU Dresden). 232 of these publications were of relevance for the database, 95 provided additional information, 134 were not relevant, and for 6 publications the evaluation was not possible, since the full PDF was either not available or did not contain enough information to analyze the study.

All in all, 1678 entries were included in the database OEKOTOX_{upgrade} (1434 for vertebrates, plants, protozoa and bacteria and 244 for invertebrates). Because many studies investigated multiple chemicals and endpoints, the number of database entries is much higher than the number of studies.

Figs. 1 and 2 show that most of the studies were conducted with antibiotics, followed by NSAIDs, anticonvulsants, antiparasitics, β-blockers and contraceptives, and that the number of publications per pharmaceutical class is reflected by the number of database entries.



Figure 1: Number of publications per substance class



Figure 2: Number of database entries per substance class

3.1.2 Number of publications per organism group

Most publications on effects of pharmaceuticals in biota have been found for fish, followed by mollusks, plants/algae, crustaceans, and bacteria (Fig. 3). In general, this distribution pattern for the publication number is reflected by the number of database entries per organism group (Fig. 4), however, for plants/algae more data were available per publication than for mollusks, and for bacteria more database entries have been conducted than for crustaceans.



Figure 3: Number of studies per biota group



Figure 4: Number of database entries per biota group

3.1.3. Evaluation of data

3.1.3.1 Most sensitive biota classes for pharmaceuticals

In a first step, the lowest and second lowest effect data were analyzed with respect to their relation to distinct organism classes (Fig. 5).

It became evident that most of the lowest plus second lowest effect values were recorded for protozoans (represented by only a single test species: *Tetrahymena pyriformis)*. Mollusks were shown to be very sensitive for anticonvulsants, antipsychotics, lipid regulators, analgesics (other than NSAIDs), hormones and contraceptives. The LOECs were in the range of ng/L – lower μ g/L values.

Lowest or second lowest effect values for fish (several species) were determined for hormones, mycolytica, anticonvulsants, antiparasitics and lipid regulators. Also for them, the LOECs are in the range of ng/L - lower $\mu g/L$ values.

Bacteria and plants were shown to be highly sensitive to antibiotics, crustaceans were very sensitive to antiparasitics.

In a second step, data were analyzed in a more detailed way by determining the number of pharmaceuticals for which defined LOECs (< 0.1, 1 or $10\mu g/L$) were reported for the investigated species in order to identify the most sensitive species for pharmaceuticals.

Figure 6 shows that the most sensitive species with LOECs < 0.1 for 2-8 pharmaceuticals were the ciliate *Tetrahymena pyriformis*, zebrafish *(Danio rerio),* rainbow trout *(Oncorhynchus mykiss),* medaka *(Oryzias latipes),* the mussel *Elliptio complanata* and the mudsnail *Potamopyrgus antipodarum.*

All extremely low LOECs (in the range of pg/L) for *Tetrahymena pyriformis* derive from only a single publication by Lang & Köhidai (2012) [3], the reliability of which, however, could not be

proven (chapter 3.1.5). Additionally, the conclusion on the most sensitive species is biased by the fact that these species are also among the most frequently used test organisms.



Figure 5: Number of lowest plus second lowest effect values for 13 pharmaceutical classes



Figure 6: Number of pharmaceuticals with defined LOECs for investigated species

3.1.3.2 Most sensitive effect endpoints per substance classes

In order to identify most sensitive effect endpoints for pharmaceuticals, data were analyzed with respect to the number of pharmaceuticals with defined LOECs (< 0.1, 1 or 10μ g/L) for the investigated effect endpoints. Figure 7 makes evident that the most sensitive endpoints with LOECs < 0.1 for 3-8 pharmaceuticals were chemotaxis, behavior, vitellogenin induction, growth rate, reproduction, histopathological alterations, molecular stress biomarkers, oxidative stress markers, receptor binding, and gene expression. All extremely low LOECs (in the range of pg/L) for chemotaxis derive from the above mentioned study of Lang & Köhidai (2012) [3] who used *Tetrahymena pyriformis* as test organism. As mentioned above, the reliability of this study, however, could not be proven (shown in chapter 3.1.5).



Figure 7: Number of pharmaceuticals with defined LOECs for investigated effect endpoints

3.1.3.3 Identification of pharmaceuticals of highest relevance

To identify the risk exerted by chemicals for the aquatic environment, different types of risk quotients are calculated in ecological risk assessment (PEC/PNEC ratios for industrial chemicals and pharmaceuticals, TER for pesticides). The PEC/PNEC ratio is defined by (1) the PEC, which represents the concentration of a chemical supposed to occur in the environment (predicted environmental concentration), and (2) the PNEC (predicted no effect concentration) which is calculated on the basis of the lowest observed effect concentration in any group of biota (LOEC), divided by a safety factor which itself depends on the size and quality of the data for different trophic levels. A possible risk for the environment is indicated by a risk quotient larger than 1.

In the present study, the calculation of the risk quotient slightly differs from this routine procedure: For those pharmaceuticals, for which both Measured Environmental Concentrations (MECs) (either from Bergmann et al., 2011 [1] or UBA; summarized in data sheet ""MECs & LOECs")") and LOEC values were available, MEC/LOEC quotients were calculated as a proxy for the environmental relevance of LOECs by using the highest available MEC (MEC_{max}) and the lowest LOEC (LOEC_{min}). In contrast to the established prospective risk assessment procedure, we used LOECs (lowest observed effect concentrations) instead of NOECs. This can be justified by the necessity to identify threshold values for effects as a prerequisite for pharmaceutical monitoring. A further reason is that our literature survey aims at identifying pharmaceuticals of environmental concern retrospectively, and thus had to take into consideration concentrations measured in the environment (MECs). Finally, only for 37% of the LOECs corresponding NOECs were available. If NOECs were used, the database would have become too small for the envisaged analyses.

For the calculations of $MEC_{max}/LOEC_{min}$ values also data from Bergmann et al. (2011) [1] were analysed.

Figure 8 makes evident that most pharmaceuticals with LOECs leading to risk quotients above 0.1 were antibiotics and NSAIDs followed by β -blockers and lipid regulators, analgesics different from NSAIDs and contraceptives/hormones.

When analyzing data with focus on $MEC_{max}/LOEC_{min}$ values for the respective biota classes (Fig. 9) it became evident that most pharmaceuticals, for which risk quotients >0.1 were calculated were investigated in plants/algae and invertebrates closely followed by vertebrates. For bacteria, only few risk quotient > 0.1 were found.

For the following substances, the risk quotient values were >1 (>0.1):

Vertebrates: >1: atenolol, carbamazepine, diclofenac, 17α -ethinylestradiol, 17β -estradiol, gemfibrozil, ibuprofen, metoprolol, paracetamol, propranolol (>0.1: bezafibrate).

Invertebrates: >1: bezafibrate, carbamazepine, diclofenac, erythromycin, 17α -ethinylestradiol, ibuprofen, ivermectin, oxytetracycline, paracetamol, propranolol, sulfamethoxazole (>0.1: ciprofloxacin, gemfibrozil, 17β -estradiol)

Plant/Protozoa: >1: acetylsalicylic acid, diatrizoic acid, diclofenac, erythromycin, lincomycin, metoprolol, naproxen, oxytetracycline, paracetamol, propranolol, sulfadimidine, sulfamethoxazole (>0.1: clotrimazole, ibuprofen)

Bacteria: >1: paracetamol, erythromycin, sulfadimidine, sulfamethoxazole (>0.1: diclofenac, gemfibrozil).



Figure 8: Number of pharmaceuticals per pharmaceutical class with $MEC_{max}/LOEC_{min}$ values >0.1



Figure 9: Number of pharmaceuticals with defined MEC_{max}/LOEC_{min} values per organism group

Table 2 makes evident that, according to the calculated risk quotients, the pharmaceuticals of highest relevance (with $MEC_{max}/LOEC_{min}>0.1$) in at least three different biota classes were: Paracetamol (analgesic, acetaminophen),

Diclofenac, ibuprofen (analgesic, NSAIDs - non-steroidal anti-inflammatory drugs) Propranolol (ß-blocker)

Sulfamethoxazole, Erythromycin (antibiotics)

Gemfibrozil (lipid regulator)

Table 2: Priority pharmaceuticals for which $MEC_{max}/LOEC_{min}$ values were >0.1 in 1, 2, 3 or 4 biota groups before reliability evaluation, including data from $OEKOTOX_{uograde}$ and Bergmann et al. (2011) [1]

| MEC _{max} /LOEC _{min} >0.1 in |
|---|---|---|---|
| 4 organismic groups | 3 organismic groups | 2 organismic groups | 1 organismic group |
| Paracetamol | Ibuprofen | Bezafibrate | Acetylsalicylic acid |
| Diclofenac | Propranolol | Carbamazepine | Atenolol |
| | Sulfamethoxazole | 17β-Estradiol | Ciprofloxacin |
| | Gemfibrozil | 17 α -Ethinylestradiol | Clotrimazole |
| | Erythromycin | Metoprolol | Diatrizoic acid |
| | | Oxytetracycline | Lincomycin |
| | | Sulfadimidine | Naproxen |
| | | | Ivermectin |

3.1.3.4 Data for mixtures

In general, only for a few cases, a direct comparison between LOECs for an isolated substance and its toxicity in a pharmaceutical mixture could be realized, since data differed with respect to species, effect endpoints or test designs. However, for a few substances, such a comparison was possible (summarized in Table 3)

The table makes evident that, dependent on - at least - the chemical tested and the effect endpoint under investigation, the toxicity of a single pharmaceutical could be lower, higher or equal to its toxicity when applied in a mixture with other compounds.

However, since no mode of action-based effect endpoints were investigated in any of these studies, the specific contribution of the respective substance to the toxicity of the chemical mixture cannot be quantified.

	Species	Effect	LOEC single	Effect conc. mixture	Reference
Paracetamol	Danio rerio	reproduction output	10	0,5	Galus et al., 2013a [9, 10]
	Danio rerio	mortality	0,5	10	Galus et al., 2013a [9,10]
Propranolol	Mytilus galloprovincialis	cAMP-level, PKA activity	0,0003	NOEC 0,0003	Franzellitti et al. 2011, 2013 [11,12]
Diclofenac	Dreissena polymorpha	molecular stress marker	0,3	0,1	Parolini et al, 2013 [13]

Table 3: Comparison of pharmaceutical effect concentrations applied either isolated or mixture of pharmaceuticals

3.1.3.5. Data with population relevance

All data collected in the present study were investigated with respect to their population relevance. We defined data as being relevant for the population level as containing information on effects related to (1) community changes, (2) reproduction, fecundity, fertility, embryo development, sex ratio, intersex, imposex (summarized as "reproduction"), (3) behavior including mating behavior, (4) growth, and (5) survival / mortality.

Altogether, 106 studies and 561 database entries were found to be related to populationrelevant endpoints. Figure 10 shows that most database entries contained information of influences of pharmaceuticals on survival and mortality, growth and reproduction. Studies related to behavioral and community changes were in the minority. Hormone-like acting substances did induce population-relevant effects already in relatively low concentrations.



Figure 10: Number of database entries with population-relevant endpoints

3.1.4 Reliability of publications

For those studies which contained data for the most relevant pharmaceuticals (lowest LOECs) and for the most sensitive endpoints and organisms, the reliability was investigated according to the criteria of Wright-Walters et al. (2011) [2].

These are the following:

"1) A thorough description of the experimental design, including exposure regime and replication,

2) Analytical confirmation of test concentrations

3) Description of ecologically relevant endpoints and all supplemental morphological information collected

- 4) Use of test procedures that are based, at least generally, on internationally accepted procedures and practices. Newly developed test procedures must be able to be repeated, and meet all other required criteria
- 5) Clear linkage of reported findings with the exact experimental design, and
- *6)* Sufficient reporting of results, including system performance, toxicity results, and statistical methods employed to ascertain how the data support the conclusions that are drawn "

If criteria 1, 2, 3 and 6 were met the study was designated as "reliable". If only 3 of these criteria were met, it was designated as "conditionally reliable", If less than 3 criteria were met or mistakes became obvious, the study was designated to be "not reliable". Consideration was also given to whether the studies were conducted according to GLP and whether both NOEC and LOEC values were provided; however, these criteria did not influence the decision about reliability.

Taking all together, 72 publications have been checked for reliability. Nine of them were reliable, 49 conditionally reliable and 14 not reliable (of which 4 were, in part, conditionally reliable). In table 4, the results of the reliability assessment are summarized. For studies which were not reliable, further information is given in Table 5. Except for the study of Fairchild et al. (2011) [14], none of the investigations were conducted under GLP.

Reference	Reliability criteria met	Reliabilty
Antunes et al., 2013 [15]	1,3,6	conditionally reliable
Backhaus et al., 2011 [16]	3, (6)	not reliable
Bajet, 2012[17]	1,3,6	conditionally reliable
Benstead, 2011 [18]	1,2,3	conditionally reliable
Boltes et al., 2012 [19]	1,3,6	conditionally reliable
Boonstra et al., 2011 [20]	1,2,3	conditionally reliable
Chandra et al., 2012 [21]	1,3,6	conditionally reliable
Chen et al., 2012 [22]	1,3, 6	conditionally reliable
Claessens et al., 2013 [23]	1,3,6	conditionally reliable
Contardo-Jara, 2011 [24]	1,2,3	conditionally reliable
Doyle et al., 2013 [25]	1,3,6	conditionally reliable
Fairchild et al., 2011 [14]	1,2,3,6	reliable
Feito et al., 2012 [26]	1,3,(6)	conditionally reliable (mitochondrial activity, DNA content), not
		reliable (lipid peroxidation, chlorophyll content)
Feito et al., 2013 [27]	1,3, (6)	conditionally reliable (DNA-content), not reliable (mitochondrial
		activity)
Finn et al., 2012 [28]	2,3,6	conditionally reliable
Fong & Hoy, 2012 [29]	1,3,6	conditionally reliable
Franzellitti et al., 2011 [11]	1,3,6	conditionally reliable
Franzellitti et al., 2013 [12]	1,3,6	conditionally reliable
Gagné et al., 2012 [30]	2,3,6	conditionally reliable
Galus et al., 2013 [10]	(1),2,3,(6)	not reliable
Gonzalez-Rey, 2011 [31]	1,3,6	conditionally reliable
Gust et al., 2012 [32]	1,3,6	conditionally reliable
Gust et al., 2013 [33]	1,3,6	conditionally reliable
Hallgren et al., 2011 [34]	(1),3,6	conditionally reliable
Hallgren et al., 2012 [35]	1,2,3,6	reliable
Hillis et al., 2011 [36]	1,3,6	conditionally reliable
Hoffmann & Kloas, 2012 [37]	1,3,6	conditionally reliable
Huynh Thi et al., 2012 [38]	1,2,3,6	reliable
Ings et al., 2012 [39]	1,3,6	conditionally reliable
Johns et al. 2011 [40]	1,3,6	conditionally reliable
Kaptaner et al., 2011 [41]	1,3,6	conditionally reliable
Lang & Kohidai, 2012 [3]	1,3,(6)	not reliable
Lange et al., 2012 [42]	2,3,6	conditionally reliable
Lawrence et al., 2012 [43]	1,3,6	conditionally reliable

Table 4: Publications checked for reliability, reliability criteria met, and decision on reliability

Reference	Reliability criteria met	Reliabilty
Lei et al., 2013 [44]	1,3,6	conditionally reliable
Liu et al., 2011 [45]	1,3,6	conditionally reliable
Liu et al., 2011 [46]	1,2,3,6	reliable
Madureira et al., 2011 [47]	1,3,6	conditionally reliable
Meina et al., 2013 [48]	1,3,6	conditionally reliable
Notch &Mayer, 2013 [49]	3,(6)	not reliable
Oliveira et al., 2012 [50]	1,3,6	conditionally reliable
Ozdemir et al., 2011 [51]	(1),3,6	conditionally reliable
Parolini et al., 2011 [52]	3,6	not reliable
Parolini et al., 2011 [53]	1,3,6	conditionally reliable
Parolini et al., 2013 [13]	3,6	not reliable
Ragugnetti et al., 2011 [54]	1,3,6	conditionally reliable
Reyhanian et al., 2011 [55]	1,3,6	conditionally reliable
Ribeiro et al., 2012 [56]	1,3,6	conditionally reliable
Rocco et al., 2012 [57]	(3), 6	not reliable
Rocco et al., 2012 [58]	(3), (6)	not reliable
Saravanan et al., 2011 [59]	1,3,6	conditionally reliable
Sarria et al., 2011 [60]	1,3,(6)	conditionally reliable,
Shen et al., 2012 [61]	1,2,3,6	conditionally reliable
Shi et al., 2012 [62]	1,3,(6)	conditionally reliable (chronic test), not reliable (acute test)
Silva et al., 2012 [63]	1,2,3,6	reliable
Skolness et al., 2012 [64]	1,2,3,(6)	reliable
Sponchiado et al. 2011 [65]	3,6	not reliable
Stange et al., 2012 [66]	1,3,6	conditionally reliable
Stange et al., 2012 [67]	1,3,(6)	cond. reliable (reproduction), not reliable (gene expression)
Thomas et al., 2012 [68]	1,3,6	conditionally reliable
Toumi et al., 2013 [69]	1,2,3,6	reliable
van Leeuwen et al., 2012 [70]	(1),3,6	conditionally reliable
Veach et al., 2012 [71]	1,3,6	conditionally reliable
Wang & Gunsch, 2012 [72]	1,2,3,6	reliable
Wang et al., 2011 [73]	1,3,6	conditionally reliable
Wu et al., 2012 [74]	1,3,6	conditionally reliable
Yan et al., 2013 [75]	1,2,3,6	reliable
Yergeau et al., 2012 [76]	(1), 6	not reliable
Yonar et al., 2011 [77]	1,3,6	conditionally reliable
Zhang & Gong, 2013 [78]	1,3,6	conditionally reliable
Zhang etal., 2012 [79]	1,3,6	conditionally reliable
Zhang et al, 2012 [80]	1,3,6	conditionally reliable

Table 5: Further information on pharmaceuticals tested, t	test organisms,	effect endpoints and	l reasons for l	lacking reliability
for not reliable studies				

Reference	Pharmaceutical tested	Test organism	Effect endpoint	Reason for lacking reliability
Backhaus et al 2011 [16]	Fluoxetine Propranolol Clotrimazole	Periphyton community	Inhibition of total pigment content (biomass)	Lacking replicates for some test substances, but not specified (N= 1 to N=5). Results from other studies were involved for calculations of some concentration-response-curves without representing them.
Feito et al 2012 [26]	Diclofenac	Danio rerio Polystichum spicatum	Lipid peroxidation Chlorophyll autofluorescence	Partly not reliable for lipid peroxidation in zebrafish and Chlorophyll content in <i>Polystichum</i> due to lacking concentration-effect relationships.

Reference	Pharmaceutical tested	Test organism	Effect endpoint	Reason for lacking reliability
Feito et al 2013 [27]	Venlafaxine	Polystichum spicatum	Mitochondrial activity	Partly not reliable for mitochondrial activity due to lacking concentration- effect relationships.
Galus et al 2013 [10]	Paracetamol Venlafaxine Carbamazepine Gemfibrozil	Danio rerio	Reproductive output Embryonic mortality Developmental malformations Histopathological changes Plasma estradiol level Blood 11- Ketotestosterone level	Experimental mistake obvious with high concentrations of pharmaceuticals in control treatments, for Gemfibrozil higher than the treatment "low". This information can be obtained from the "Supplementary data", and it is mentioned in one sentence of the discussion. The effective concentrations highly differ from the nominal concentrations for which results are presented. For acetaminophen, e.g., the effective concentrations were only about 10% of the nominal concentrations.
Láng & Köhidai 2012 [3]	Acetylsalicylic acid Diclofenac Fenoprofen Ibuprofen Naproxen Paracetamol Erythromycin Lincomycin Sulfamethoxazole Trimethoprim Metoprolol Propranolol Timolol Diatrizoic acid	Tetrahymena pyriformis	Growth rate Chemotactic behaviour	Calculation of EC50 values for growth inhibition unclear; no concentration- effect relationships for chemotaxis; authors recommend themselves to use the test system not as a quantitative, but a qualitative assay to prove for environmental effects of chemicals.
Notch & Mayer 2013 [49]	17alpha- ethinylestradiol	Danio rerio	Embryonic vitellogenin mRNA embryonic Cyp1a mRNA embryonic XPC mRNA (genome repair pathway) embryonic XPA mRNA (genome repair pathway)	Lacking replicates and insufficient description of exposition conditions. Effects for XPC/XPA-mRNA und CYP1a- mRNA which only occurred after 24h or 48 h. The authors wanted to show that some effects disappear after longer exposure times. Effects for vtg-mRNA remained stable for the entire exposure time.
Parolini et al 2011 a [52]	lbuprofen	Dreissena polymorpha	Molecular stress biomarkers	Lacking replicates. Chemical analysis only in stock solution.
Parolini et al 2013 [13]	lbuprofen Diclofenac Paracetamol	Dreissena polymorpha	Molecular stress biomarkers	Lacking replicates. Chemical analysis only in stock solution.
Rocco et al 2012 a [57]	Gemfibrozil	Danio rerio	Comet assay Diffusion assay RAPD-PCR	Insufficient description of experimental design (replica, organisms, test concentrations); results from controls not presented.
Rocco et al 2012 b [58]	Lincomycin Erythromycin	Danio rerio	Micronucleus test Comet assay	Insufficient description of experimental design (replica, organisms, test concentrations): no significant effects.
Shi et al 2012 [62]	Clotrimazole	Xenopus laevis	Embryo mortality Embryo body length	lacking concentration-effect relationships in acute test
Sponchiado et al 2011 [65]	17beta-estradiol	Oreochromis niloticus	Micronucleus test Nucleus abnormalities Comet assay	Lacking replicates. Exposure from 5 to 35 days, but no renewal of test substances.
Stange et al 2012b [67]	17alpha- ethinylestradiol	Potamopyrgus antipodarum	Gene expression (estrogen receptor)	Effects on gene expression of estrogen receptor only occurred after 7 days, but not after 28 days, lacking time-effect relationshin

Reference	Pharmaceutical tested	Test organism	Effect endpoint	Reason for lacking reliability
Yergeau et al 2012[76]	Erythromycin Sulfamethoxazole Sulfadimidine Gemfibrozil Erythromycin Sulfamethoxazole Sulfadimidine Gemfibrozil	Bacterial community	Microbial community composition (DNA level)	Insufficient description of experimental design (substances, solvents for stock solution, organisms). Exposure time 8 weeks without renewal of test substances.

3.1.5 Consequences of reliability assessment for the described results

Our literature study revealed *Tetrahymena pyriformis* as the most sensitive test organism for pharmaceuticals prior to the reliability check of the publications. This assessment, however, is based on the fact, that for this protozoan extremely low LOECs are reported for a total of 8 substances by Láng & Köhidai (2012)[3] who used "chemotaxis" as an effect endpoint. Since the reliability assessment could not prove the reliability of this publication, three fish species (zebrafish, rainbow trout and medaka) can be identified as most sensitive organisms for pharmaceuticals followed by bivalves and snails. The publications which provide the lowest LOECs for these species did successfully pass the reliability assessment. Table 6, however, also makes evident, that for the three fish species the database was much larger (in total 489 database entries) than for the mentioned molluskan species (only 8 database entries). Also for crustaceans which represent ecologically important species of aquatic ecosystems only few data are available compared to fish.

	Number of publications	Number of database entries	Number of database entries with LOEC ≤0.1	Number of reliable database entries with LOEC ≤0.1	Lowest LOEC [µg/L]	Lowest reliable LOEC [µg/L]
Tetrahymena pyriformis	1	28	13		1,51E-05 (Láng & Köhidai 2012) [3]	-
Zebrafish (Danio rerio)	36	333	36	21	0,0104 (Lange et al 2012) [42]	0,0104 (Lange et al 2012) [42]
Rainbow trout (Oncorhynchus mykiss)	18	92	5	5	2,66E-03 (Ings et al 2012) [39]	2,66E-03 (Ings et al 2012) [39]
Medaka (Oryzias latipes)	8	64	12	12	0,001 (Lei et al 2013) [44]	0,001 (Lei et al 2013) [44]
mussel <i>(Elliptio</i> <i>complanata)</i>	2	4	3	3	0,04 (Gust et al, 2012) [32]	0,04 (Gust et al, 2012) [32]
mudsnail <i>(Potamopyrgus</i> <i>antipodarum</i>)	2	4	4	3	0,05 (Stange et al, 2012a) [66]	0,05 (Stange et al, 2012) [66]
Daphnia magna	12	42	2	2	0,011 (Toumi et al, 2013) [69]	0,011 (Toumi et al, 2013) [69]
Gammarus spec	6	11	0	0	1 (Boonstra et al, 2011) [20]	1 (Boonstra et al, 2011) [20]

Tab. 6: Assessment of data for most sensitive species with respect to reliability of publications

With respect to the determination of the most important pharmaceuticals for environmental effects the reliability assessment had a more important influence on the final result. In chapter 3.1.4.3 all data available from the OEKOTOX and the OEKOTOX_{upgrade} databases were assessed with the result summarized in table 2. Since the publications of the OEKOTOX database analyzed by Bergmann et al. (2011) [1], however, were not at our disposal for reliability checks, we could only compare the results for the most important pharmaceuticals based on data of the OEKOTOX_{upgrade} database prior (table 7) and after reliability assessment (table 8). The results of the analyses based on data from OEKOTOX_{upgrade} only before reliability evaluation did not differ much from those when the OEKOTOX data were included. This is due to the fact that the original OEKOTOX database mainly reports on EC_{50} values and includes only a low number of LOECs. Only a slight difference becomes obvious for Metoprolol with MEC_{max}/LOEC_{min}>0.1 in only 1 organismic group instead of 2. After assessment of publication reliability, however, only for diclofenac MEC_{max}/LOEC_{min}-values >0.1 were found in >3 biota classes. As further important pharmaceuticals propranolol, sulfamethoxazole, bezafibrate, 17α -ethinlyestradiol, 17β -estradiol and oxytetracycline were identified with MEC_{max}/LOEC_{min}-values >0.1 for 2 biota classes.

| MEC _{max} /LOEC _{min} >0.1 in |
|---|---|---|---|
| 4 organismic groups | 3 organismic groups | 2 organismic groups | 1 organismic group |
| Paracetamol | Ibuprofen | Bezafibrate | Acetylsalicylic acid |
| Diclofenac | Propranolol | Carbamazepine | Atenolol |
| | Sulfamethoxazole | 17β-Estradiol | Ciprofloxacin |
| | Gemfibrozil | 17α-Ethinylestradiol | Clotrimazole |
| | Erythromycin | Oxytetracycline | Diatrizoic acid |
| | | Sulfadimidine | Lincomycin |
| | | | Naproxen |
| | | | Metoprolol |

Table 7: Priority pharmaceutical	s before reliability	evaluation, only based	1 on data from OEKOTOX _{upgrad} ,
----------------------------------	----------------------	------------------------	--

Table 8: Priority pharmaceuticals after reliability evaluation, only based on data from OEKOTOX_{upgrade}

| MEC _{max} /LOEC _{min} >0.1 in |
|---|---|---|---|
| 4 organismic groups | 3 organismic groups | 2 organismic groups | 1 organismic group |
| Diclofenac | | Propranolol | Ibuprofen |
| | | Sulfamethoxazole | Paracetamol |
| | | Bezafibrate | Carbamazepine |
| | | 17α-Ethinlyestradiol | Erythromycin |
| | | 17β-Estradiol | Gemfibrozil |
| | | Oxytetracycline | Sulfadimidine |
| | | | Atenolol |
| | | | Ciprofloxacin |

3.1.6 Summary of part 1

The literature survey conducted in the present project reviewed publications on pharmaceutical effects in the environment from 2011- July 2013, and thus completed the database OEKOTOX established by Bergmann et al. (2011) [1] with the database OEKOTOX_{upgrade}. An Endnote library

was created which contains all publications analyzed. In addition, a data evaluation sheet was created as a base for the identification of the most sensitive organisms, the most sensitive effect endpoints, and the ecologically most relevant pharmaceuticals.

For data published between 2011 and 2013, the following results were found:

- Most studies were conducted with antibiotics, followed by NSAIDs, anticonvulsants, antiparasitics, ß-blockers and contraceptives.
- Most of the research data are related to pharmaceutical effects in fish, followed by effects in mollusks, plants/algae, crustaceans, and bacteria.
- After the reliability evaluation three fish species (zebrafish, rainbow trout and medaka) were identified as most sensitive organisms for pharmaceuticals followed by bivalves and snails.
- Sensitive effect endpoints were behavior, vitellogenin induction, growth rate, reproduction, histopathological alterations, molecular stress biomarkers, oxidative stress markers, receptor binding, and gene expression

In order to evaluate the lowest and the second lowest effect values from both the OEKOTOX and the OEKOTOX_{upgrade} database with respect to their ecological relevance, risk quotients were calculated as $MEC_{max}/LOECmin$. Risk quotients >0.1 were defined as to be of ecological relevance. The analyses provided the following results:

- Most pharmaceuticals for which risk quotients >0.1 were calculated were investigated in plants/algae and invertebrates closely followed by vertebrates. For bacteria, only few risk quotient >0.1 were found.
- Antibiotics, analgesics (NSAIDs), and β-Blockers were the pharmaceutical classes for which data with the highest ecological relevance were found.
- After reliability assessment diclofenac was identified as the pharmaceutical of highest relevance (with MEC_{max}/LOEC_{min} >0.1 in > 3 biota classes). For propranolol, sulfamethoxazole, bezafibrate, 17 α -ethinlyestradiol, 17 β -estradiol and oxytetracycline MEC_{max}/LOEC_{min}-values >0.1 were calculated for 2 biota classes.

Only in a few cases, the toxicity of an isolated substance could be compared to its toxicity in a pharmaceutical mixture. It could be shown that, at least dependent on the chemical tested and the effect endpoint under investigation, the toxicity of a single pharmaceutical can either be lower, higher, or equal to its toxicity when applied in a mixture together with other compounds. Since no mode of action-based effect endpoints were investigated it was impossible to quantify the specific contribution of the respective substance to the toxicity of the chemical mixture under investigation.

106 studies and 561 database entries were found to be related to population-relevant endpoints with most data on survival/mortality followed by growth and reproduction. Studies related to community changes and behavioral endpoints were in the minority. Direct population relevance, for example induced changes in the composition of bacterial, protozoan or algal communities could only be shown in few studies.

From part 1 of this literature review the following shortcomings could be identified:
- More studies which fulfill the reliability criteria are necessary, especially more chemical analyses should be integrated
- Only few data for invertebrates are available
- Data for ecologically relevant crustaceans are lacking
- Data on sediment toxicity are lacking
- More population-relevant community data are necessary
- In order to be able to identify the contribution of isolated pharmaceuticals to chemical mixtures, mode of action-based effect endpoints have to be investigated.

3.2 *In vitro* test systems

3.2.1 Introduction

In vitro assays developed for the monitoring of pharmaceuticals can be broadly categorized into biochemical assays and cell-based assays. In biochemical and cell based assays, homogeneous assay readouts are preferred over non homogeneous readouts since they do not require washing steps.

Biochemical assays are developed to characterize compounds that interact with an isolated target in an artificial environment. Biochemical assays are target-based and historically have been the mainstay of substance screening in the pharmaceutical industry. Such *in vitro* assays include the assessment of enzymatic activity (e.g., for kinases [1], proteases [2], transferases [3]), receptor-ligand binding (e.g., for G-protein coupled receptors (GPCRs) [4] and nuclear receptors [5]) or protein-protein interactions [6]. Biochemical assays are often directed to and specific for a recombinant purified target of interest and are conducted in homogeneous reaction media. However, not all targets can be purified or prepared suitably for biochemical measurement. For example methods that measure ion channel activities across a biological membrane require two ionic compartments with different ion concentrations separated by a lipid bilayer in which the ion channels reside. Ion channel assays are therefore conducted in cell based assay systems. Additionally, the activity of a small molecule in a reconstituted *in vitro* assay does not always translate into the same activity in a cellular context, due to requirements for cellular cofactors, issues of membrane permeability and other reasons.

Cell-based assays have emerged as a more physiological alternative to assays involving purified proteins. Cell-based assays interact with regulatory networks and feedback control mechanisms. In contrast to biochemical target-based assays, cell-based assays can target an outcome of a pathway in the physiological environment of a cell enabling amplification of a specific signal. Examples of cell-based assays include functional assays (e.g., second messenger mobilization after GPCR activation [7, 8]) and reporter gene assays [9, 10]. Cell-based assays allow for the selection of compounds that can cross cellular membranes, a prerequisite for biological activity towards multicellular organisms.

Homogeneous assays can be categorized into radioactive and non-radioactive assays. Although radioactive assays are decreasing on the market due to the need of radioactive waste disposal, this technology is unlikely to disappear completely. Homogeneous radioactive assay technologies include scintillation proximity assay (SPA) (Amersham Pharmacia Biotech) and FlashPlates[™] (NEN Life Science Products, Boston, MA). With these approaches, the target of

interest is immobilized onto a solid support (e.g. SPA beads or FlashPlate[™] surface) that contains a scintillant [11]. When a radiolabelled molecule binds to the target molecule, the radioisotope is brought in close proximity to the solid support, and the energy transfer between the emitted beta particle and the scintillant results in the emission of light. Radioisotopes which remain unbound to the target and thus free in solution are too distant from the scintillant and, consequently, the released beta particles dissipate their energy solely into the aqueous environment. SPA has been used in a wide variety of applications and has become a standard technique in high throughput screening labs. The technology has been applied to kinases [12,13] and other enzymes [14] and is widely used for the detection of ligand-receptor interactions [15]. FlashPlate[™] technology is similar to SPA but the solid surface is a microtiter plate rather than a bead. FlashPlate[™] applications include the detection of cAMP levels [16] and ligand-receptor interactions [17]. Radiometric assays have the advantage to be relatively sensitive but, however, show also numerous disadvantages including limited reagent stability, relatively long read-times and waste disposal.

Most common homogeneous assay readouts are non-radioactive and optical, measuring absorbance and luminescence. Luminescence is the emission of light from any substance, and occurs from electronically excited states. Luminescence is formally split up into two categories (fluorescence and phosphorescence) depending on the nature of the excited state. In excited singlet states, the electron in the excited orbital is paired (by opposite spin) to the second electron in the ground-state orbital. Return to the ground state is spin-allowed and occurs rapidly by emission of a photon. Typical fluorescence lifetimes are around some nanoseconds. Many fluorophores display sub-nanosecond lifetimes. Because of the short timescale of fluorescence, measurement of the time-resolved emission requires sophisticated optics and electronics. In spite of the added complexity, time-resolved fluorescence is widely used because of the increased information available from the data, as compared with stationary or steadystate measurements. Additionally, advances in technology have made time-resolved measurements easier, even when using microscopes. Phosphorescence is emission of light from triplet excited states, in which the electron in the excited orbital has the same spin orientation as the ground-state electron. Transitions to the ground state are forbidden so that phosphorescence lifetimes are typically milliseconds to seconds. Transition metal-ligand complexes, which contain a metal and one or more organic ligands, display mixed singlettriplet states. These transition metal-ligand complexes display intermediate lifetimes of hundreds of nanoseconds to several microseconds. These optical properties can be exploited by a number of detection methods.

3.2.2 Fluorescence Spectroscopy

Fluorescence spectroscopy can be applied to a wide range of questions in the chemical and biological sciences. The measurements can provide information on a wide range of molecular processes, including the interactions of solvent molecules with fluorophores, rotational diffusion of biomolecules, distances between sites on biomolecules, conformational changes, and binding interactions. Recently, the usefulness of fluorescence has been expanded by advances in technology for development of stable cell lines and cellular imaging. These advances in technology have been decreasing costs for previously complex *in vitro* assay development.

3.2.2.1 Steady-State and Time-Resolved Fluorescence

Fluorescence measurements can be broadly classified into two types of measurements: steadystate and time-resolved. Steady-state measurements, the most common type, are those performed with constant illumination and observation. Here, the sample is illuminated with a continuous beam of light, and the fluorescence intensity or emission spectrum is recorded. The second type of measurement is time-resolved, which is used for measuring intensity decays or anisotropy decays. For these measurements, the sample is exposed to a pulse of light, where the pulse width is typically shorter than the decay time of the sample. Time-resolved measurements contain information on the timescale of conformational changes. Time resolved fluorescent measurements furthermore enable separation of signals emitted by long-lived transition metalligand complexes from other unspecific signals generated by short-lived intermediates.

3.2.2.2 Fluorescence Polarization (FP)

Fluorophores absorb light along a particular direction with respect to the molecular axes. The extent to which a fluorophore rotates during the excited-state lifetime determines its polarization or anisotropy. The phenomenon of fluorescence polarization can be used to measure the apparent volume (or molecular weight) of proteins. This measurement is possible because larger proteins (or protein complexes) rotate more slowly than small ones. Hence, if a protein binds to another protein, the rotational rate decreases, and the anisotropies increase. The application of plane-polarized light in fluorescence polarization can be used to measure a probe's rotational perturbations, thereby enabling simple assay designs dependent on a single labelled ligand [18, 19]. For example fluorescein- and rhodamine-labelled small molecules are suitable for quantifying the associations with proteins via changes in fluorescence polarization.

3.2.2.3 Fluorescence Resonance Energy Transfer (FRET)

FRET is an electrodynamic phenomenon occurring between a donor molecule in the excited state and an acceptor molecule in the ground state. The donor molecules typically emit radiation at shorter wavelengths that overlap with the absorption spectrum of the acceptor (Figure 1). The extent of energy transfer is determined by the distance between the donor and acceptor, and the extent of spectral overlap. The distance at which FRET is 50% efficient is called the Förster distance, which is typically in the range of 20 to 60 Å. Förster distances ranging from 20 to 90 Å are convenient for studies of biological macromolecules. These distances are comparable to the size of biomolecules and/or the distance between sites on multi-subunit protein complexes. Any condition that affects the D–A distance will affect the transfer rate, allowing the change in distance to be quantified. FRET can be used to measure the distance between a site on a protein and a membrane surface, the association between protein subunits, and the lateral association of membrane-bound proteins.



Figure 1: Fluorescence Resonance Energy Transfer (FRET). FRET occurs between a donor in an exited state and an acceptor in the ground state. The extent of energy transfer is determined by the distance between the donor and acceptor, and the extent of spectral overlap.

3.2.3 Fluorescence Sensing

There are several observables for fluorescence sensing (Figure 2). The most direct sensing method uses changes in the fluorescence intensity of the probe in response to the analyte. However, it is often inconvenient to use changes in fluorescence intensity for measurements due to the following reasons: Measurements which are independent of fluorophore concentration can be accomplished using wavelength-ratiometric probes, which display shifts in the absorption or emission spectra upon binding of the analyte. Another ratiometric method is fluorescence polarization or anisotropy. In this case the analyte causes a change in the anisotropy of the label. Anisotropy measurements are frequently used in competitive immunoassays, in which the actual analyte displaces a labeled substitute that is bound to specific antibody. This results in a decrease in the anisotropy of the analyte. Anisotropy values are calculated using the ratio of polarized intensity measurements. The use of an intensity ratio makes the anisotropy measurements independent of fluorophore concentration as long as the measurements are not distorted by auto-fluorescence or poor signal-to-noise ratio. Fluorescence lifetimes can also be used for sensing.



Figure 2: Spectral observables for fluorescence sensing. Sensing is performed using intensities, intensity ratios, anisotropies and time-domain lifetimes [20].

3.2.3.1 Low molecular weight Sensors

Low molecular weight sensors are available for sensing pH, chloride, oxygen, carbon dioxide, Ca²⁺, Mg²⁺, and other parameters. A survey of the literature revealed that a large number of chelate molecules can be used for imaging intracellular concentrations of cations. Calcium probes are perhaps the most widely used intracellular indicators. The salt forms of these dyes do not diffuse across cell membranes, so that the cells need to be labelled by microinjection or electroporation. 1,2-bis(o-aminophenoxy)ethane-N,N,N',N'-tetraacetic acid based chelates are also available with esterified carboxy groups, the so-called acetoxymethyl esters. In this form, the dyes are less polar, and, hence, passively diffuse across cell membranes. Once being inside the cell the acetoxymethyl esters are cleaved by intracellular esterases, and the negatively charged probe is trapped in the cells.

5', 6' Dicarboxy-2',7'-dichlorodihydrofluorescein (CDCFDA) is a fluorogenic probe which is oxidized by reactive oxygen species in living cells. The cell-permeable reagent is very weakly fluorescent while in a reduced state and upon oxidation exhibits strong fluorogenic signal. After uptake into the cell cytoplasmic esterases cleave the esters trapping the resultant charged CDCF inside the cell. CDCF is a co-substrate for peroxidases and could be used for sensing of peroxidase activities [21].

3.2.3.2 Protein Sensors

The combination of recombinant technologies with chemical cross linking enables manifold creation of protein based fluorescent sensors. One example is a MBP (maltose binding protein from E. coli) based sensor based on FRET (22). This example also illustrates the increasingly sophisticated chemistry of sensors. This maltose sensor is bound to a neutravidin surface by a biotinylated linker. MBP is labelled with a non-fluorescent acceptor quenching dye. The donor

is a signalling dye, which is also bound to the surface by a specialized linker. This linker contains a cyclodextrin that binds to MBP. These components are bound to the surface by a biotinylated DNA linker arm that allows changes in rigidity. When the linked cyclodextrin is bound to the binding pocket of MBP the donor is quenched because this binding brings the donor dye in close proximity to the quenching dye. Addition of maltose displaces cyclodextrin from MBP, resulting in increased donor intensity (Figure 3). These results describe a general strategy for surface-bound sensors that are adjustable and yield large changes in intensity. This approach is likely to be used in sensors for a wide variety of analytes.



Figure 3: Surface bound maltose sensor. Schematic of a surface-bound maltose sensor based on the maltose-binding protein from E. coli and FRET [22].

3.2.3.3 Green fluorescent protein (GFP) variants

A revolution in the field of live cell imaging occurred following the development of genetically encodable fluorescent tags to specifically label a protein-of-interest within the cellular milieu. This allowed researchers to develop fluorescent biosensors that are able to track signalling molecules within their endogenous environment. This "enlightenment" in the field of cell biology was sparked by the discovery, and subsequent cloning, of the green fluorescent protein (GFP) from the jellyfish *Aequorea victoria*. For simplifying the emission spectra and improving the spectral properties of *A. victoria* GFP for biosensor applications, the wild-type protein has been mutated in several ways. There are several important factors that must be considered when employing GFP family members as genetically encodable fluorescent tags. These include both physical and biological parameters like protein folding efficiency, stability, intrinsic brightness, photo-stability, spectral properties and environmental sensitivity.

The discovery and development of GFP family members exhibiting either red- or blue-shifted excitation/emission spectra allows fluorescence proteins to be used for a number of FRET-based applications. If prior excitation of the donor fluorophore is generated as by-product of an enzymatic reaction, such as the oxidation of luciferin, this is termed "bioluminescence resonance energy transfer" (BRET). These processes represent highly sensitive methods for measuring protein-protein interactions and conformational changes within individual proteins.

Although BRET signals are generally dimmer than those generated by FRET, this approach offers several potential advantages over FRET for live cell applications [23]. Because BRET does not require excitation of the donor fluorophore for energy transfer to occur, it does not elicit cellular auto-fluorescence. As a consequence, BRET exhibit signal-to-noise ratios that are much higher than those obtained using FRET. Likewise, the chemiluminescent nature of the excitatory light used for BRET simultaneously eliminates two of the primary sources of error associated with FRET-based measurements: spectral bleed-through (donor emission into the

acceptor channel and excitation of acceptor molecules by the donor excitation wavelength) and photobleaching of the donor molecules by excitatory light. Finally, since the donor and acceptor molecules used for BRET generate luminescent and fluorescent signals, respectively, the expression levels of each component can be measured independently of the other. Thus BRET-based probes hold potential for live cell applications, including some unique applications in cell based compound screening [24] and *in vivo* pharmacodynamics studies for drug candidates.

Relatively large fluorescent proteins could be replaced by short peptide sequences designed to bind small molecule probes [25]. This can result in restoring activity which was lost due to bulky fluorescent protein fusion tags. However the chemical modifying of the peptide sequences for generation of fluorescent peptides is laborious and modifying reagents need to cross cell membranes. Therefore it is more convenient to use genetically encodable fluorescent proteins to build fluorescent biosensors for probing cellular components within their native environment. Such biosensors have been used to monitor a multitude of cellular processes, offering valuable insights into the dynamic nature of the signalling, metabolic and other regulatory networks that govern cell function. The design and application of a series of fluorescent biosensors range from relatively simple sensors designed to measure changes in the expression to more complex reporter systems designed to probe biochemical processes, such as second messenger turnover and enzyme activities within their native cellular environment [26].

Monitoring Protein Expression

Fluorescence protein based transcriptional reporters have proven to be valuable tools for studying transcriptional activities. One of earliest uses of GFP as a biological probe involved the *in vivo* visualization of promoter activation in the nematode *C. elegans*. During these studies, GFP DNA was placed under the control of a promoter and GFP gene expression was measured during different stages of nematode development [27]. During these studies, GFP DNA was placed under the control of a promoter and GFP gene expression was measured during different stages of nematode development. Over the years, similar studies have been conducted in a variety of cellular contexts, offering valuable information about the activation and regulation of cellular promoters. The stability of GFP inside the cell (t1/2 = -1 day) [28] allows the activation of weak promoters to be measured. The persistence of GFP molecules long after transcription mask transient changes in gene expression. One way to overcome this limitation is to fuse a degradation sequence, such as a domain from mouse ornithine decarboxylase, to the fluorescence protein indicator [29]. Using this strategy, the fluorescent half-life of an EGFPmouse ornithine decarboxylase chimera was reduced nearly 12-fold, to approximately two hours. The increased temporal resolution afforded by this probe allowed the observation of transient changes in NF- κ B-mediated gene expression that could not be observed using longerlived EGFP reporters.

Monitoring Biochemical Changes within Cellular Environment

Through protein engineering efforts, fluorescent protein can be generated in such a way that its spectral properties are altered in response to specific cellular factors. Fluorescent protein based reporters that change their spectral properties in response to cellular parameters have also been used extensively to study the regulation of many cellular processes. While in some cases these types of sensors exploit photophysical properties intrinsic to the fluorescent protein tags themselves, in others, the spectral properties of the chromophore are altered by distorting the architecture of the fluorescent protein using integrated protein sequences that are sensitive to the signalling molecule under study. Engineered molecular switches, which are constructed based on a modular design, consist of a "receiver" module that specifically recognizes the small-molecule-of-interest linked to a "switching" module that converts the binding event into a conformational change. By combining the molecular switch with an appropriate reporter unit (either by grafting it into the fluorescence protein itself or by flanking the switch region with complementary FRET pairs) small molecule-dependent changes in the sensor unit can be translated into fluorescence readout from the reporter unit. Using this basic design, researchers have constructed a diverse set of fluorescence protein based biosensors capable of probing a large number of small molecule analytes involved in cellular signalling.

3.2.3.3.1 Single Fluorophore Sensors

3.2.3.3.1.1 Halide and pH Sensors

Unlike wild-type *A. victoria* GFP, whose spectral properties remain relatively unchanged over a wide range of physiological conditions, the mutations introduced into many engineered fluorescence protein variants often render them sensitive to fluctuations in the cellular environment. For example, because of changes in their internal hydrogen bonding networks, the chromophores of EGFP and several EYFP family members fluoresce very weakly in the protonated state. As a consequence, cellular components that promote protonation, such as protons or halide ions, can lead to markedly reduced fluorescence intensities. This intrinsic property has been exploited to measure several important cellular parameters, including pH and chloride ion concentrations.

The YFP variant YFP(H148Q) exhibits a dramatic reduction in its fluorescence intensity at elevated halide concentrations. This property has been exploited to monitor fluctuations in chloride ion concentration in cells overexpressing the chloride transporter cystic fibrosis transmembrane regulator. These studies demonstrated that YFP(H148Q) faithfully reports chloride flux through the cystic fibrosis transmembrane regulator in response to elevations in intracellular cyclic AMP, providing an attractive alternative to chemical halide dyes [30]. While EGFP (pKa = 6.15) is well-suited for the detection of pH fluctuations within acidic organelles, EYFP (pKa = 7.1) is preferable in more basic environments like the mitochondrial matrix where equilibrium pH values are closer to 8.0 [31].

3.2.3.3.1.2 Redox Potential Sensors

Sensor units can be constructed by introducing residues on the surface of a fluorescence protein variant that renders it sensitive to a particular cellular parameter. This is the basis for redoxsensitive fluorescence protein indicators, termed roGFPs, which use disulfide bond formation between pairs of strategically placed cysteine residues on the surface of GFP to alter the β -barrel architecture of the protein [32]. In the case of roGFP2, structural analysis suggests that the formation of a disulfide bond between the engineered cysteine residues promotes reorganization of two β -strands in close proximity to the chromophore, causing subtle rearrangements in the residues surrounding the chromophore. These conformational changes ultimately shift the equilibrium between the neutral and phenolic states of the chromophore, resulting in reciprocal changes in the excitation maxima at 400 and 480 nm in response to changes in the redox potential [33]. As genetically encodable and genetically targetable redox sensors, roGFP family members as well as the related rxYFPs, have proven to be valuable tools for measuring the redox potential of many subcellular compartments, including the cytosol [34]

3.2.3.3.1.3 Calcium Sensors

The circularly permutated EGFP-based GCa-MPs [35] are among the most popular genetically encodable Ca2+ indicators. These fluorescence proteins were generated by fusing the carboxy terminus of calmodulin with the calmodulin binding peptide M13. In these reporters the switch is grafted into a single fluorescence protein variant. In the presence of Ca²⁺, the CaM-M13 interaction induces conformational changes in the reporter that result in a change in fluorescence intensity of the chromophore. Such Ca²⁺ sensors have been used to measure Ca²⁺ fluxes in a variety of cell types, offering valuable insights into the timing and regulation of Ca²⁺ transients during many cellular processes.

3.2.3.3.1.4 cGMP Sensors

A series of fluorescent indicators of cGMP (FlincGs) based on the regulatory domain of protein kinase G was developed [36]. These sensors consist of a circularly permutated EGFP variant fused to two cGMP binding sites derived from various regions of the protein kinase G regulatory domain. To probe changes in cGMP concentration over a wide range of conditions, FlincG family members have been engineered to bind cGMP with KD's ranging from 35 nM to 1.1 μ M. Nucleotide binding induces conformational changes in the protein kinase regulatory domain that lead to a corresponding increase in the fluorescence intensity of the reporter. Importantly, each of the FlincG probes also exhibits a high degree of selectivity for cGMP relative to the chemically similar second messenger cAMP. This specificity for cGMP, coupled with their rapid and reversible binding kinetics and relatively good pH stability, makes FlincGs well-suited for measuring cGMP inside cells.

3.2.3.3.1.5 Reactive Oxygen Species Sensor

A hydroperoxide sensor was generated by inserting circularly permutated EYFP into the regulatory domain of a hydroperoxide sensing bacterial protein [37]. This hydroperoxide sensor exhibits reciprocal changes in the emission intensity at 520 nm when excited by 405 and 488 nm light. Using this sensor, fast and reversible changes in the fluorescence intensity of circularly permutated EYFP in a cell line after exposure to 50 μ M hydroperoxide was observed.

A circularly permuted yellow fluorescent protein can be used for sensing superoxide radical anion, the primal reactive oxygen species generated by the electron transport chain. The fluorescence emission (at 515 nm) of purified circularly permutated YFP when excited at 488 nm is five times brighter under strong oxidizing conditions compared to strong reducing conditions, indicative of a large dynamic range. Extensive in vitro experiments revealed the superoxide selectivity of circularly permutated YFP over other physiologically relevant oxidants and metabolites. Compared to the fully reduced state, circularly permutated YFP fluorescence displays a 250% increase by oxygenation and a full 420% increase by the superoxide radical anion. The superoxide radical anion associated increase in circularly permutated YFP fluorescence is completely reversed by subsequent addition of copper/zinc-superoxide dismutase. By contrast, circularly permutated YFP emission is unchanged by hydroxyperoxide and peroxynitrite, and is decreased by hydroxyl radical and nitric oxide. Other metabolites tested, including physiological levels of Ca²⁺, ATP, ADP, NAD(P)⁺, and NAD(P)H, all exert negligible or only marginal effects [38]. Unlike GFP-based redox biosensors the fluorescence emission of circularly permutated YFP is unaltered when the redox potential varies between -319 mV and -7.5 mV. The reversibility of circularly permutated YFP permits real-time measurements of dynamic changes in superoxide levels in living cells.

3.2.3.3.1.6 ATP/ADP ratio sensing

Perceval, an ATP/ADP indicator is based upon the bacterial ATP-binding protein, GlnK1, and a circularly permutated version of Venus [39]. In the presence Mg-ATP, the T-loop is converted from an extended structure to a highly compact form that is believed to relieve strain on the integrated circularly permutated Venus module. In Perceval, T-loop closure leads to reciprocal changes in circularly permutated Venus emission intensity when excited with 405 and 490 nm light. Therefore, Perceval offers the advantage of a radiometric readout. Mg-ADP also elicits a change in the emission ratio of the probe; however, because of incomplete loop closure, this change is only about half that observed upon Mg-ATP binding. As a consequence, Perceval can be calibrated to measure the relative ratio of ATP to ADP inside cells based upon competition between the two molecules for T-loop binding. As a proof of principle, the authors demonstrated that Perceval could detect global metabolic changes caused by the inhibition of glycolysis or the modulation of external glucose levels [39].

3.2.3.3.1.7 Membrane Potential sensing

Fluorescence protein based voltage sensors modulate fluorescence intensity of the fluorescence protein reporter module by placing it between regions of an ion channel or voltage-sensitive protein that undergoes a conformational change in response to changes in the membrane potential [40]. A Cerulean-based voltage sensor, that utilizes conformational changes in the voltage-sensitive domain of the voltage sensor containing phosphatase from *Ciona intestinalis* to affect the fluorescence intensity of Cerulean [41]. These properties make this voltage sensor well-suited to measure the fast neuronal electrical signals often observed during signal propagation.

Beside single fluorophore sensors like the above mentioned, more complex sensor units are required to convert changes in other cellular parameters into a fluorescence output. A "molecular switch" can be derived from a conformational change intrinsic to an endogenous protein or it can be generated via an engineered switch (Figure 4).



Figure 4: Molecular switch. A molecular switch can be generated by either (A) a conformational change intrinsic to a protein domain or (B) by an engineered conformational change driven by interactions between a receiver module (grey block) and a switching module (green block). In each case, the conformational change is converted to a change in FRET efficiency (blue arrow) by altering the distance and/or orientation of the attached fluorescence protein reporter units.

3.2.3.3.2 FRET / BRET based Sensors

FRET-based sensors exist as uni- and bimolecular reporter systems, each of which utilizes a molecular switch to convert activity-dependent changes in the reporter into a measurable FRET response. In each case the molecular switch is designed to alter the distance or relative orientation of a FRET pair in response to specific cellular conditions. For instance, whereas most unimolecular sensors rely upon a conformational change to reposition their fluorophores in space, bimolecular probes typically bring their FRET pairs into close proximity via protein-protein interactions. An advantage of a bimolecular design is that the reporter typically exhibits a larger dynamic range than its unimolecular counterpart. In the case of bimolecular reporter systems the stoichiometries between the donor and acceptor are variable. Therefore more sophisticated measures of FRET efficiency, such as donor fluorescence recovery after acceptor photo bleaching and fluorescence lifetime imaging, must be used.

3.2.3.3.2.1 FRET based cAMP Sensor

Several FRET-based biosensors have been developed to better understand how the concentrations of cAMP fluctuate over time and throughout the cell. These include the unimolecular reporter system derived from various portions of the guanine nucleotide exchange factor, exchange protein directly activated by cAMP (Epac) [42]. Several intramolecular cAMP reporters have been developed based upon cAMP effector Epac1 and Epac2. These reporters using Epac (ICUE) [43, 44], CFP-Epac (δ DEP-CD)-YFP [45], Epac1-camps and Epac2-camps [42] all exhibit decreasing FRET following cAMP binding. Presumably, the binding of cAMP induces an intrinsic conformational change in Epac isoforms that liberates the catalytic domain from intrasubunit allosteric inhibition, thereby altering the distance and relative orientation of their flanking CFP/YFP FRET pairs. Furthermore, mutations have been introduced to abolish the guanine exchange factor activity of Epac, thereby reducing the effects that overexpression of the biosensor may have on cellular signalling pathways. cAMP sensor studies have begun to uncover important details about the role of subcellular compartmentalization in shaping cAMP dynamics during GPCR activation.

3.2.3.3.2.2 BRET based cAMP Sensor

By replacing the CFP donor in Epac-based sensors with luciferase, cAMP reporters were created that utilize BRET as the fluorescent readout [46]. Though their emission intensity has prevented BRET-based sensors from being used to measure biochemical changes at the subcellular level, BRET-based cAMP reporters have proven to be powerful tools for examining drug effects in cell populations in a medium- to high-throughput manner and for screening fluorescent compounds whose excitation/emission profiles preclude the use of FRET-based probes. For instance, using a BRET-based cAMP sensor derived from the second generation ICUE reporter, ICUE2, the impact of nine clinically effective antipsychotics on D2 class dopamine receptor activity was examined [47].

3.2.3.3.2.3 FRET based GPCR activation Sensor

The GPCR superfamily, which represents the largest family of proteins involved in signal transduction, is composed of structurally-similar receptor proteins characterized by seven R-helical membrane-spanning domains. As the primary upstream activators of many intracellular signalling pathways, GPCRs play a key role in converting extracellular stimuli, such as hormones and neurotransmitters, into an intracellular response. The deregulation of GPCRs is also critical to the etiology of many diseases. In fact, roughly one half of the drugs on the market today target GPCRs. For gaining insights into spatial and temporal aspects of GPCR

activation and signalling as well as into the mechanistic basis of the signalling process itself, several FRET-based sensors have been created to study receptor activation in the context of single, living cells [48]. Upon ligand binding, GPCR family members act as quanine nucleotide exchange factors that facilitate the exchange of GDP for GTP in the GR subunit of associated trimeric G-protein complexes. Nucleotide exchange promotes the dissociation of the GR subunit from the $G\beta_{Y}$ subunits of the complex which, in turn, leads to the activation of downstream effectors, such as phospholipase C and transmembrane adenylate cyclases involved in phosphoinositide and cAMP metabolism. To measure activation-induced conformational changes in GPCR family members, several groups have fused CFP and YFP color variants to the third intracellular loop and the C-termini of the receptor molecule to yield sensitive GPCR activation sensors [49]. These sensors, which all exhibit reproducible decreases in emission ratio following agonist stimulation, have been used to study GPCR activation kinetics in the presence of full agonists, partial and even inverse agonists. In contrast to agonists generating a decrease in FRET inverse agonists appear to cause an increase in FRET that may be attributed to distinct conformational states of the receptor [50]. Effects of beta blockers bisoprolol, metoprolol and carvedilol in inhibition β 1-adrenoceptor isoforms in living cells was measured in real time by FRET using a cyan fluorescent protein, Cerulean fused to the carboxy terminus of the human β 1-adrenoceptor and a yellow fluorescent protein inserted into the third intracellular loop [51].

3.2.3.3.2.4 FRET based protein kinase activity Sensor

Protein kinase activity reporters utilize an engineered molecular switch based upon a modular design. A consensus phosphorylation site specific for the kinase of interest serves as the "receiving segment" while a phosphoamino acid binding domain functions as the "switching segment". These regions are concatenated together by a flexible linker and sandwiched between a fluorescence protein FRET pair. Whereas the length of the linker and the choice of FRET pairs influence the dynamic range of the reporter, the "receiving and switching segments" contribute to its specificity and reversibility. This basic modular design has been applied to a number of protein kinases, including protein kinase A. Genetically encoded A-kinase activity reporters are reversible and targetable reporters allowing real-time imaging of protein kinase A activity, and are valuable for analyzing compartmentalized kinase activities [52].

3.2.3.3.2.5 Coupled FRET based NO Sensor

A coupled fluorescent indicator system was used for reporting NO release. The FRET based cGMP indicator "CGY" exhibits a rapid and reversible decrease in emission ratio in response to elevations in cGMP produced by NO-dependent activation of soluble guanylyl cyclase. Since it is estimated that a single NO molecule generates nearly 6,000 cGMP molecules/min, this indicator system is extremely sensitive to NO release [53]. In fact, this indicator system is able to detect NO concentrations in the picomolar range, several orders of magnitude lower than direct NO fluorescent indicators.

3.2.3.3.3 Protein Fragment Complementation Sensor

Fluorescence based approaches has paralleled the study of GPCR oligomerization. GPCRs are the largest family of cell surface receptors and are the targets many clinical drugs (e.g. β-blockers and antipsychotics). Bimolecular fluorescence complementation relies on the generation of a fluorescent signal from two non-fluorescent fragments of a fluorescence protein when brought in close proximity by fusion partners. Receptor activation through binding of a ligand agonist proceeds by conformational rearrangement within the transmembrane helical

domain as the receptor switches from an inactive to an active state, which in turn activates heterotrimeric G proteins. Activated G proteins regulate the levels of intracellular second-messenger molecules (Ca²⁺, cAMP, phoshoinositides, cGMP), which modulate signalling cascades involving kinases such as PKA, PKC and others. The 2012 Nobel Prize in Chemistry was awarded to Brian Kobilka and Robert Lefkowitz for their work that was crucial for understanding GPCR function.

Proteins interacting with GPCRs have been shown to modulate receptor expression, membrane targeting, and desensitization. Among these, arrestins are well characterized scaffolding proteins that are notably involved in receptor internalization after ligand activation. As part of a high-content protein fragment complementation assay study based on fluorescence protein complementation, interaction between β -arrestins and the β 2-adrenoceptor was detected in cells treated with β 2-adrenoceptor agonists, thus providing a measure of receptor activation and internalization [54]. These studies demonstrate the applicability of fluorescent protein fragment complementation assays for the detection of drug-induced changes in GPCR interactions. Likewise, non-fluorescent protein fragment complementation assays based on β -galactosidase fragment complementation can also be used for the detection of GPCR- β -arrestin interactions.

3.2.4 Immunoassays

Immunoassays constitute a large and diverse family of assays. The basic idea is to couple the association of antibody with antigen to some other event that yields an observable spectral change. Various mechanisms are possible, including energy transfer, anisotropy, delayed lanthanide emission, or the use of enzymes to amplify the signal. The use of antibodies as analytical tools can be traced to the development of radioimmunoassays by Berson and Yalow, which resulted in a Nobel Prize. Since then immunoassays have been widely used, but are now based mainly on fluorescence detection.

3.2.4.1 cAMP AlphaScreen® (PerkinElmer)

The acronym ALPHA stands for Amplified Luminescent Proximity Homogeneous Assay. The assay contains two bead types, donor beads and acceptor beads. Both bead types provide reactive aldehyde groups for conjugating biomolecules to the bead surface. Donor beads contain a photo-sensitizer, phthalocyanine, which converts ambient oxygen to an excited form of O2, singlet oxygen, upon illumination at 680 nm. Singlet oxygen has a limited lifetime prior to falling back to ground state. Within its 4 µsec half-life, singlet oxygen can diffuse approximately 200 nm in solution. If an acceptor bead is within that proximity, energy is transferred from the singlet oxygen to thioxene derivatives within the acceptor bead, subsequently culminating in light production at 520– 620 nm (Figure 5). In the absence of an acceptor bead, singlet oxygen falls to ground state and no signal is produced.



Figure 5: AlphaScreen cAMP assay. Competitive assay type where endogenous cAMP produced in whole cells competes with biotinylated cAMP for binding to an anti-cAMP antibody conjugated to the Acceptor beads.

3.2.4.2 cAMP LANCE® (PerkinElmer)

LANCE stands for Lanthanide chelate exite. This assay is a homogeneous time-resolved fluorescence energy transfer immunoassay designed to measure cAMP produced upon modulation of adenylyl cyclase activity. The assay is based on the competition between a Europium-labelled cAMP tracer and sample cAMP for binding sites on cAMP-specific antibodies labelled with a FRET acceptor dye (ULight). When the ULight anti-cAMP antibody is bound to the Eu-cAMP tracer, excitation at 340 nm excites the Europium. The energy is transferred to the ULight-labelled antibody. The fluorescence measured at 665 nm will decrease in the presence of cAMP from test samples, and resulting signals will be inversely proportional to the cAMP concentration of a sample (Figure 6).



Figure 6: LANCE cAMP assay. Competitive assay type where endogenous cAMP produced in whole cells competes with a cAMP antibody complex.

3.2.4.3 cAMP HTRF (CisBio International)

HTRF (homogeneous time resolved fluorescence) combines standard FRET technology with time-resolved measurement of fluorescence. The cAMP assay is based on a competitive immunoassay using cryptate-labelled anti-cAMP antibody and d2-labelled cAMP (Figure 7).



Figure 7: HTRF cAMP assay. Competitive assay type where endogenous cAMP produced in whole cells competes with a cAMP d2 complex. Donor: Europium cryptates (Eu3+cryptate), Acceptor: d2 (an organic motif of approximately 1 kDa).

3.2.4.4 cAMP Screen (Life Technologies)

The cAMP Screen is a competitive immunoassay. Cell lysates are incubated with a cAMP-AP conjugate and an anti-cAMP antibody in the secondary antibody–coated assay plate. The resulting immune complexes are captured in the plate. The captured immune complexes are washed to remove unbound cAMP-AP, and the resulting signal is measured in a luminometer.

3.2.5 Reporter Assays

Cell based reporter assays provide a cost effective platform for sensing promoter activities. Reporter gene constructs contain a promoter element which regulates the expression of a selected reporter protein. Commonly used reporters are enzyme proteins with activities linked to colorimetric or luminescent readouts such as luciferase, alkaline phosphatase, β -galactosidase, β -lactamase or fluorescent proteins such as GFP variants. The advantages of reporter gene assays include the wide linearity and sensitivity of the technique and a large signal to background ratio, making them suitable for amplification of a single signal induced by a receptor agonist.

GPCR activation is well known to alter gene transcription via responsive elements for second messengers including the cAMP response element (CRE), the nuclear factor of activated T-cells response element (NFAT-RE), the serum response element (SRE) and the serum response factor response element (SRF-RE, a mutant form of SRE), all of which are located within gene promoter regions. G-protein-dependent reporter gene assays were developed using second messenger responsive elements upstream of a minimal promoter, which in turn regulate the expression of a selected reporter protein (Figure 8).

Reporter gene assays are also easy to set up and can be scaled down to extremely low assay volumes. Despite these advantages, some concerns have been raised, such as the requirement for long incubation periods, difficulty in antagonist detection due to reporter accumulation and the higher potential for false positives because the signal event is distal from receptor activation.

Concerns about the long incubation time and accumulation of reporter have been addressed through the use of destabilized reporters. The higher false positive rate due to the distal signalling event could be partially resolved with by the co-expression of a constitutively expressed internal control, so compounds non-specifically affecting gene transcription could be ruled out [55].



Figure 8: Receptor binding and G-protein-dependent reporter gene assays. Schematic representation of receptor binding and major pathways activated by different G proteins. RE symbols a specific second messenger responsive element upstream of a minimal promoter, which in turn regulate the expression of a selected reporter protein.

3.2.6 β -adrenoceptor assays

Activation of cardiac β -adrenoceptors by endogenous catecholamines plays a key role in the regulation of cardiac function. The heart contains at least 2 β -adrenoceptor subtypes, termed β 1-AR and β 2-AR, and may also contain β 3-ARs. Stimulation of the β 1-AR represents the strongest endogenous mechanism for increasing contractility and beating frequency of the mammalian heart [56]. In human heart failure, which has become one of the leading causes of death and hospitalization, the sympathetic nervous system is chronically activated to overcome the loss of cardiac output. While this initially leads to compensation through a short-term increase in cardiac function, chronic stimulation of the cardiac β -AR system contributes to progression of the disease. These detrimental effects of chronic β -adrenergic signalling are attributed to the β 1-AR subtype. Consequently, the development of receptor antagonists has resulted in the single most effective therapeutic regimen to treat heart failure [57]. Beta blockers are part of most commonly prescribed drugs in Germany.

 β -adrenoceptors are a class of GPCRs. The subtype β 1-AR is linked to Gs protein which in turn is linked to adenylate cyclase (Figure 8). Downstream effectors of cAMP include cAMP response element and cAMP-dependent protein kinase, which mediate some of the intracellular events following hormone binding.

Measuring of β -adrenoceptor activity could be done using *in vitro* assays like tagged-ligand binding, GTP γ S binding, CRE reporter, cAMP or conformation based FRET assays.

3.2.6.1 Tagged ligand binding assays

Tagged ligand binding assays are based on antibodies binding to cAMP. Commercial available are several competitive assays like cAMP AlphaScreen (3.2.4.1), cAMP LANCE (3.2.4.2), cAMP HTRF (3.2.4.3) or cAMP screen (3.4.4). All assays measure cAMP in cellular extracts, therefore cells expressing β -adrenoceptor has to be lysed first.

3.2.6.2 cAMP luminescent biosensor assay

A novel protein biosensor is described, which uses a cAMP-binding domain from PKA coupled to a circularly permuted form of *Photinus pyralis* luciferase. Upon binding of cAMP the luciferase undergoes conformational change from an open less reactive to a closed form resulting in high luciferase activity (Figure 9) [58].



Figure 9: cAMP luciferase assay. Schematic representation of the cAMP luciferase assay

The real-time cAMP luminescent biosensor assay allows the sensitive detection of GPCR mediated signalling through the second messenger cAMP. In contrast to lytic assays with multiple samples and procedural steps, cAMP dynamics can be followed from a population of living cells with greatly reduced time and labour. When compared to existing biosensor designs, this evolved biosensor construct provides increased dynamic range, making it better suited to monitor a broad range of receptor behaviours such as treatment with full, partial, or inverse agonists.

3.2.6.3 cAMP CRE reporter assays

After activation of the β -adrenoceptor adenylyl cyclase is activated by a coupled Gs protein. Adenylyl cyclase catalyzes the formation of cAMP which activates PKA. PKA in turn initiates the downstream kinase cascade. One of the end steps of this cascade is the phosphorylation of the cAMP response element binding protein. The phosphorylated CREB can bind promoter regions containing the 5'-TGACGTCA-3' sequence i.e. the cAMP response element, and in consequence induces downstream gene transcription. An externally-introduced reporter gene usually contains a specific promoter and a reporter gene DNA. The promoter can be made artificially by fusion of CRE elements with a minimal promoter sequence. For real time imaging a CRE-EGFP reporter system was developed, which could be used for monitoring real time transcription under a fluorescent microscope or by flow cytometry [59].

3.2.6.4 Receptor conformation based FRET assay

Beside FRET based (3.2.3.3.2.1) and BRET based cAMP sensors (3.2.3.3.2.2) a FRET approach is described were the human β 1-adrenoceptor was fused at the carboxyterminus to a mutant CFP (Cerulean) and a YFP inserted into the third intracellular loop (Figure 10). After binding of the agonist norepinephrine a conformational shift leads to a decrease of FRET signal. The β -blocker bisoprolol, metoprolol, and carvedilol led to an active change of the receptor conformation resulting in an increase in the FRET ratio, suggesting inverse agonist behaviour [51].



Figure 10: Development of a madrenoceptor FRET sensor. After binding of the agonist norepinephrine a conformational shift leads to an increase of distance of the two fluorescent proteins resulting in a decreasing FRET effect [51].

3.2.6.5 cAMP gated ion channel

cAMP produced in subcellular compartments near the plasma membrane can be monitored in HEK cell lines expressing the rat wild-type (GenBank CAA39135.1) or several mutant olfactory CNG channel proteins after recombinant adenovirus infection [60]. Forskolin induced cAMP synthesis resulted in Ca²⁺ influx in the cells. Using the wild-type channel, a forskolin induced modest Ca²⁺ influx could be measured after pre-treatment of the cells with phosphodiesterase inhibitors only. Site specific mutations of the wild-type protein resulted in special mutant channel proteins with enhanced cAMP sensitivity and specificity [60].

3.2.6.6 HCN2 cAMP FRET assay

A cAMP FRET sensor based on a single cAMP binding domain of the hyperpolarizationactivated cyclic nucleotide-gated channel 2 (HCN2) enabled studies of spatial and temporal cAMP dynamics after β 1- and β 2- adrenoceptor stimulation in freshly isolated adult cardiomyocytes [61].

3.2.7 Cyclooxygenase assays

Cyclooxygenase-1 and cyclooxygenase-2 are bifunctional enzymes that carry out two sequential reactions in spatially distinct but mechanistically coupled active sites: the double dioxygenation of arachidonic acid to prostaglandin G2 and the reduction of prostaglandin G2 to prostaglandin H2. Arachidonic acid oxygenation occurs in the cyclooxygenase active site, and prostaglandin G2 reduction occurs in the peroxidase active site. Prostaglandin H2 diffuses from the COX proteins and is transformed by different tissue specific isomerases to prostaglandins (PGE2, PGD2, PGF2 α , PGI2) and thromboxane A2 (TxA2)

Although crude and purified preparations of cyclooxygenase isoenzymes have been used in characterizing inhibitors, several studies indicate that for unknown reasons the potency and selectivity of inhibitors determined using intact cells expressing cyclooxygenase isoenzymes differ from values established using cell-free cyclooxygenase preparations. For example,

ibuprofen is approximately 10-fold more potent against cyclooxygenase-2 in intact cells than against cyclooxygenase-2 activity in broken cells [62].

Several functional COX assays using purified or recombinant enzymes have been reported and include an oxygen consumption assay, a peroxidase co-substrate oxidation assay, a radiolabelled chemical inhibition assay, and an enzyme-linked immunosorbant assay.

3.2.7.1 Oxygen consumption assay

COX enzymatic activity was determined by measuring oxygen consumption at 37°C in an oxygraph chamber using an YSI Model 53 oxygen monitor [63]. Tests were performed using isolated protein.

3.2.7.2 Peroxidase co-substrate oxidation assay

Peroxidase activity of the cyclooxygenase could be measured in a peroxidase co-substrate oxidation assay using guaiacol, heme and hydroperoxide [64]. Assays were performed with recombinant purified protein.

3.2.7.3 COX Immunoassay

COX activity in protein solutions can be measured using a commercial Prostaglandin E2 competitive immunoassay kit. The kit uses a monoclonal antibody to Prostaglandin E2 in a competitive manner. Prostaglandin E2 from the sample competes with an alkaline phosphatase Prostaglandin E2 fusion molecule (#ADI-901-001 Enzo Life Sciences).

3.2.7.4 COX fluorescence assay

During a cellular assay the peroxidase substrate, 5- (and 6)-carboxy-2',7'dichlorodihydrofluorescein (CDCF) was passively incorporated into cells as the acetoxymethyl ester, 5- (and 6)- carboxy-2',7'-dichlorcdihydrofluorescein diacetate (CDCFDA). Following partitioning into the cell, cytoplasmic esterases cleave the methyl esters and trap the resultant charged CDCF. CDCF serves as a reducing substrate for the peroxidase activities of cyclooxygenase-1 and cyclooxygenase-2. PGG2 generated by the cyclooxygenase activity upon the addition of arachidonate is reduced to PGH2 by the peroxidase activity with resultant oxidation of CDCF to a fluorescent product that can be detected by fluorescence microscopy (Figure 11) [65].



Figure 11: COX fluorescence assay. Fluorescence assay for cyclooxygenase activity in living cells. After loading of the cells with CDCFDA arachidonate (RH) is added and oxidised. Peroxides are reduced by COX peroxidase activity with resultant oxidation of CDCF to a fluorescent product that can be detected by fluorescence.

3.2.8 Sensitivity of in vitro and in vivo assays

Different sensitivity of substances in *in vivo* compared to *in vitro* tests may be related to metabolic activities. The metabolic capacity of a living organism could differ from the metabolic activity of single cells. Dependent on this difference an *in vitro* test system could have an enhanced sensitivity over the *in vivo* test system. It is also possible that a metabolite from a substance (e.g. the endocrine disrupting chemical flutamide) exert a higher binding activity on the mammalian hormone receptor as the substance itself. The binding constant of a substance to its biological target molecule influences the lowest observed effect concentration in an *in vitro* assay system. The detection limit of much pharmaceuticals in *in vitro* assays is in the range of nM to μ M concentration due to the corresponding IC₅₀ concentrations of 10⁻⁶ to 10⁻⁹ mol/L. However, when accumulation occurs in an animal, *in vitro* test systems can be less sensitive.

3.2.8.1 Comparative in vivo / in vitro analysis of endocrine disrupting chemicals

Endocrine disrupting chemicals are defined as exogenous agents that interfere with the production, release, transport, metabolism, binding, action or elimination of natural hormones in the body. A variety of developmental and reproductive disorders observed in wildlife species have clearly been linked to the exposure to endocrine disrupting chemicals. The importance of identifying potential endocrine disruption has been recognized by regulatory bodies. In order to consider potential endocrine disrupting effects the US Environmental Protection Agency established the Endocrine Disruptor Screening Program. This program employs an approach including a combination of *in vitro* and *in vivo* mammalian and ecotoxicological screens for identifying and characterizing endocrine effects of pesticides, industrial substances, and environmental contaminants. Part of the tier 1 screening is a Fish Short-Term Reproduction test and an Amphibian Metamorphosis Assay. These *in vivo* tests are used to identify potential endocrine disrupting endocrine states are used an Amphibian Metamorphosis Assay. These *in vivo* tests are used to identify potential endocrine disrupting chemicals. Because this regulatory approaches focus on screening assays that use animals and are several weeks in duration shorter-term *in vitro* tests that reduce the number of animals and time are required.

In vitro screening for endocrine disruption could address many different hormone systems, such as reproductive hormones (e.g. estrogens, androgens, progesterons), or thyroidal hormones, corticosteroids, growth hormone and their associated hypothalamus/pituitary releasing and stimulating hormones. To date, most alternative assays for detecting potential disruption of estrogen, androgen and thyroid pathway regulation evaluate receptor binding/transactivation, receptor mediated gene/protein expression or hormone synthesis.

For a comparative analysis published *in vivo* data on reproductive effects and metamorphosis were compared with published data of *in vitro* receptor binding or reporter gene assays [66]. For nearly every compound, for which *in vivo* data were available, some alternative *in vitro* assays showed a sensitivity similar to or greater than these *in vivo* data. However, the correlation to *in vivo* data was biased *by* using mainly *in vivo* data for compounds with affinity to the estrogen receptor [66]. Comparative *in vivo* and *in vitro* data for substances with other mode of actions (e.g. COX-inhibitors, β -adrenoceptor blockers) are still lacking.

3.2.9 Evaluation

3.2.9.1 Selection of substances with high priority

There have been many instances in which pharmaceuticals and their metabolites have been identified in water effluents. For developing an *in vitro* assay measuring biological activity of a group of pharmaceuticals first a target molecule has to be defined. An assay could then be developed where the target molecule itself or a downstream signalling event leads to a signal read out after binding of a pharmaceutical to its biological target.

Pharmaceuticals with high priority according to UBA are the β -blocker Atenolol, Bisoprolol, Metoprolol and Propranolol. All this pharmaceuticals as well as metabolites having similar biological activity could be measured by an *in vitro* assay monitoring inhibition of the β -1 adrenoceptor.

Another group with high priority according to UBA are analgesic drugs including Diclofenac, Ibuprofen, Naproxen, Metamizole and 4-N-Methylaminoantipyrin. All these substances as well as the lower priority drugs Indometacin and Mesalazine inhibit the activity of the cycloxygenase 1 and/or 2 (Cox1, Cox2). Therefore an *in vitro* assay measuring inhibition of Cox activity could be used for monitoring a biological effect caused by one or more pharmaceuticals characterized by this mode of action.

3.2.9.2 Evaluation of published *in vitro* assays

After selection of β -blocker and analgesic drugs for *in vitro* assay development, corresponding published *in vitro* assays are evaluated in the following chapter. A practicable solution for *in vitro* assay systems is a cell based assay system using a stable transfected cell line expressing its recombinant gene of interest in an inducible fashion. In such a case false positive results could be excluded by using the non-induced cell line in parallel. Most of the published cell based *in vitro* assays use cell lines expressing the recombinant protein continuously. There are several inducing expression systems published and it should be possible to use an inducible expressing system for the development of cell line dependent *in vitro* assay systems.

3.2.9.2.1 β-adrenoceptor compatible *in vitro* assays

Biosensors for β 1-adrenoceptor activation/inactivation are published for direct and coupled downstream activities (Table 1.)

The most specific β -1 adrenoceptor (β 1-AR) inhibition assay measures the β 1-AR conformation change by a conformation based FRET assay (β 1-AR sensor, 3.2.6.4). Concentration-response curves of β 1-AR sensor activation for norepinephrine and isoproterenol yielded half-maximal effective concentration (EC₅₀) of 1.800 ± 200 nM and 230 ± 40 nM. Due to a direct conformational effect in transiently transfected cells the FRET signal (Δ FRET) was only 5%. This signal was reversible by addition of the antagonist propranolol (10µM).

Downstream effects of β -1 AR activation like cAMP generation or gene activation can be utilized to monitor β -1 AR inhibition. These assays would have the advantage of higher signal read out due to signal amplification. One have to have in mind that levels of intracellular cAMP are tightly regulated, with degradation controlled via the cAMP phosphodiesterase enzymes. When cAMP is produced it binds to protein kinases within the cell, initiating phosphorylation events that regulate target enzymes and transcription factors. There are a variety of cAMP phosphodiesterase enzyme isoforms, which are generally activated by cAMP dependent protein kinases, thus providing an important negative feedback system on the receptor-mediated signalling cascade and regulating the extent of changes in intracellular cAMP concentrations.

Most sensitive cAMP immunoassays (3.2.4) are competitive assays and share the disadvantage that increases in cAMP produce a decrease in signal, making them liable to false positives.

A very sensitive *in vitro* assay utilizes reporter genes that contain a cAMP response element that regulates the transcription of an enzyme or a fluorescent/bioluminescent protein (3.2.6.3). Synthetic promoters made up of multiple copies of these sequences are routinely employed in reporter genes and have been extensively used to study GPCRs. The reporter protein needs to have a short half-life to minimize basal accumulation of reporter proteins that can restrict the sensitivity of the final readout. In the case of GFP, this can be achieved by creating a destabilized version of GFP by fusing a degradation domain from mouse ornithine decarboxylase to the C-terminal of GFP. As a result of substantial signal amplification between ligand binding and the final measured response ligands with partial agonist activity are likely to manifest themselves as full agonists in most reporter gene *in vitro* assays. This results in the observation that many β -blockers in common clinical practice produce substantial agonist effects at β 1- and β 2-adrenoceptors when measured at the level of gene expression.

Kinetic studies of the time course of agonist-stimulated gene expression have been undertaken using reporter genes. These studies have shown that a minimum of 30 min of agonist exposure is required to detect a measureable change in reporter gene activity and that it is the duration of cAMP elevation rather than the total quantity of cAMP produced that is the major determinant of the final response. The resulting requirement for sustained stimulation needs to be taken into account when designing an inhibitory assay and the potential for receptor desensitization with highly efficacious agonists during the time course of the assay means that lower efficacy agonists should be employed.

Direct cAMP detection is possible using firefly luciferase-based biosensors. Genetic manipulation of firefly luciferase into a reversible biosensor of cAMP generation (3.2.6.2) resulted in broad linearity coupled with high sensitivity of intracellular cAMP concentration determination. The firefly luciferase based cAMP biosensor has a pEC₅₀ for cAMP of 6.3 and a large signal-to-noise window of approximately 70-fold. When expressed in HEK293 cells, the addition of 10 μ M forskolin, a direct activator of adenylyl cyclase, can mediate a 25-fold

increase in the luminescent signal within 3.5 min. This sensor represents a powerful method to detect the kinetics of cAMP generation. This cAMP luminescent biosensor assay is covered by a patent application (EP2281046). However, when interpreting kinetics of the luminescent signal, consideration must be given towards the potential for a delay between the real-time cAMP dynamics and the generation of the active form of firefly luciferase.

Another assay type for detecting intracellular cAMP concentration uses FRET based cAMP sensors (3.2.3.3.2.1.). These sensors are used for investigating the spatial and temporal characteristics of cAMP signalling at a single cell level.

Biosensor	Signal	Advantages/Disadvantages	Ref.
β 1-adrenoceptor	FRET	Highly specific	[51]
cAMP luminescent biosensor	Luminescence	cAMP EC50= 6µM, large signal to noise window / patent pending	[58]
cyclic nucleotide- gated channels			
CNGA2	Ca ²⁺ current	cAMP EC50= 1µM, conversion of cAMP into Ca ²⁺ signal, rapid transient signal, complicated read out.	[60]
HCN2-camps	FRET	cAMP EC50= 6µM, good for high basal cellular cAMP	[61]
PKA based cAMP			
sensor			
R-CFP, C-YFP	FRET	cAMP EC50= 0,5-0,9 µM, multimeric	[69]
PKA-camps	FRET	cAMP EC50= 1,9µM, single chain (faster kinetics)	[42]
Epac based cAMP			
SEIISUI Enaci/2-commo	EDET	$AMD ECEO = 2.4 / 0.0 \mu M$ single shain	[42]
		CAMP EC50- 2,47 0,9 µm, single chain	[42]
		CAMP ECSUE 0,3 µm, single chain, nigh sensitivity	
СЕР-Ерас-ҮЕР	FREI	cAMP EC50= 50 µM, low sensitivity	[45]
ICUE1/2	FRET	cAMP EC50= 10-50µM, low sensitivity	[44]
cAMP Reporter gene			
CRE EGFP	Fluorescence	basal CRE signal activity essential for cell viability	[59]
CRE Luciferase	Luminescence	reporter accumulation	[59]

Table 1: [-adrenoceptor biosensors. Published biosensors for adrenoceptor and coupled downstream activities

PKA and EPAC-based FRET sensors have a similar affinity for cAMP of approximately $0.3-3 \mu$ M and a dynamic range of approximately $0.1-10 \mu$ M. As a consequence of their relatively high sensitivity, these biosensors are likely to be quickly saturated in cell types that have particularly high concentrations of cAMP. When expressed in HEK cells, the EPAC-based cAMP probe had a uniform, cytosolic distribution. However, the compartmentalization of cAMP responses results in cAMP signalling targeted to discrete microcellular domains. Generally, the activation kinetics of plasma membrane targeted cAMP sensors are more rapid and of greater amplitude than those of their cytosolically distributed equivalents. This may reflect the fact that the plasma membrane is the site of cAMP production and/or the restriction of cAMP diffusion within microdomains located near the plasma membrane.

cAMP gated ion channels like the rat oCNG channel (3.2.6.5) can induce a Ca^{2+} influx as a result of local increase of cAMP concentration [60]. The Ca^{2+} current was measured by patch clamp recording and fura-2 fluorescence measurement.

3.2.9.2.2 Cyclooxygenase compatible *in vitro* assays

From published Cyclooxygenase assays (Table 2) only the COX fluorescence assay is suitable for application in a homogeneous assay system without the need of using purified protein preparations.

The cyclooxygenase reaction is part of an enzymatic cycle after generation of a tyrosyl radical in the cyclooxygenase active side. This enzymatic reaction converts arachidonic acid into prostaglandin G2. The lipid hydroperoxide prostaglandin G_2 is then reduced to prostaglandin H2 (Figure 13). Leakage of the peroxyl radical from the cyclooxygenase active side leaves the enzyme in a catalytically inactive form. Reactivation of the cyclooxygenase activity requires reaction of the heme prosthetic group with another molecule of hydroperoxide explaining the need for continued presence of hydroperoxide in cyclooxygenase-arachidonic acid reactions [67]. The reduction of hydroperoxide prostaglandin G2 to prostaglandin H2 needs a cellular electron donor. It is likely that other cellular enzymes take part in the reduction of prostaglandin G2. A possible candidate enzyme is thioredoxin reductase. The mammalian thioredoxin reductase shows reducing activity towards a variety of substrates. It was shown that mammalian thioredoxin reductase reduces arachidonic acid hydroperoxides [68]. Thioredoxins are small (Mw 12.000) disulfide-containing redox proteins known to be present in all eukaryotic and prokaryotic organisms. Thioredoxins are members of the ubiquitous thiol-disulfide oxidoreductase family representing the major ubiquitous disulfide reductases responsible for the maintaining proteins in reduced state in the cytoplasm. Thioredoxins are reduced by electrons from NADPH via thioredoxin reductase.

COX assay	Signal	Advantages/Disadvantages	Ref.
Oxygen consumption assay	l (current)	Clark type oxygen electrode, test use purified COX protein	[63]
Peroxidase co- substrate oxidation assay	Photometric (E436)	Guaiacol oxidation assay with purified COX protein	[63]
COX Immunoassay	Photometric (E405)	Competitive PGE2 immunoassay kit, use of purified protein	(3.2.7.3)
COX fluorescence assay	Fluorescence	Cellular assay using stable cell lines, light and oxygen sensitive reagent	[65]

Table 2: Cyclooxygenase activity assays. Published biosensors for measuring cyclooxygenase activity activities

A frequent protein modification as result of oxidative stress is the oxidation of sulfhydryl groups. The redox active protein component cystein would be oxidized completely under normal atmospheric conditions to form a disulfide bond. An unwanted formation of disulfide bonds in the cytoplasm also called "disulfide stress" could be reversed by redox regulation. The simplest scheme of modulation redox state is depicted in Figure 12.



Figure 12: Modulation of redox regulation.



Figure 13: Arachidonic acid metabolism through the COX pathway. COX-1 and COX-2 convert arachidonic acid to the intermediate prostaglandin PGG2 and then to PGH2 which either spontaneously decomposes or is converted by other enzymes to form primarily PGE2 and PGD2

It was shown by Kargman et al. [65] that in CHO cell lines stably expressing cyclooxygenase isoforms exogenous arachidonic acid was metabolized. Cyclooxygenase activity was measured after loading of cells with dichloro-dihydrofluorescein diacetate by fluorescence after converting of the ester into the carboxylic acid by cellular esterases (3.2.7.4). Due to instability of the reagent ester, inhomogeneities in cellular loading and hydrolysis of the reagent into the fluorescent dye, this assay seems to be not convenient for a robust *in vitro* assay method.

3.2.9.3 Suggestion for in vitro assay development

For development of homogenous *in vitro* assay systems with high sensitivity and specificity it is recommended to engineer β -blocker and NSAID biosensor cell lines. The cell lines should stably express their transgenes in an inducible form. The tests should be conducted with induced and non-induced cells in parallel, thus enabling test controls excluding false positive signals. β blocker biosensor cell lines should monitor the β -adrenoceptor dependent cAMP signal. cAMP should be measured by the FRET sensor Epac (3.2.3.3.2.1) and in parallel by the calcium sensor GCaMP (3.2.3.3.1.3) after converting the cAMP signal into a Ca²⁺ current. NSAID biosensor cell lines should be based on the detection of lipid peroxides. The cyclooxygenase dependent lipid peroxide generation should be monitored by the redox potential sensor roGFP (3.2.3.3.1.2).

3.2.9.3.1 β -adrenoceptor *in vitro* assay suggestion

In a first step, two parallel approaches are recommended:

1. A FRET based microscopic assay (as described in 3.2.3.3.2.1). This assay enables detection of space resolved signals. Although new microtiter plate readers with higher sensitivity are commercially available it is not clear whether this sensitivity is sufficient for measuring FRET signals without spatial resolution. cAMP is restricted into microdomains located near the plasma membrane. This could result in a limited signal range. Because of this uncertainty it is suggested to develop a second approach:

2. An assay system where a membrane located cAMP signal is converted into a cytoplasmic Ca^{2+} signal as described in 3.2.6.5. Such a signal could be measured in a microtiter assay format without the need of spatial resolution. This should be possible cloning a mutant of the cAMP gated rat olfactory CNG channel [60]. The Ca^{2+} influx could be measured by a Ca^{2+} dependent fluorescent dye [60] or by a genetically encoded Ca^{2+} biosensor (3.2.3.3.1.3).

For development of a homogeneous cell based assay system, a cell line expressing the β -1 adrenoceptor together with an Epac1/2 cAMP sensor in an inducible form should be developed and tested whether the cAMP coupled FRET signal is sufficient for measuring in a microtiter plate assay (approach 1).

In addition, a cell line has to be engineered in such a way that the cell line expresses the β -1 adrenoceptor, a specific mutant of the cyclic nucleotide gated channel CNGA2 and the Ca²⁺ sensor GCaMP3. All proteins should be expressed stably in an inducible form (approach 2).

For saving costs it is recommended to prove in principal functionality on a very early development step for the two *in vitro* test systems and to define appropriate milestones. After reaching the first milestone it has to be decided on the test system which has to be further developed.

If β -blockers are present in the test sample the inhibition of the β -adrenoceptor leads to a reduced cAMP generation and thus to an increase of the Epac FRET signal (approach 1) or a decrease of Ca²⁺ dependent fluorescence (approach 2) both relative to the control.

3.2.9.3.2 Cyclooxygenase in vitro assays suggestion

For screening of COX inhibitors, a test based on the detection of redox potential changes described in 3.2.3.3.1.2 is recommended. The inducible expression of cyclooxygenase in a genetically engineered cell line should result in generation of lipid peroxides after addition of arachidonic acid. Due to this inducible peroxide generation the redox state should be

influenced resulting in some disulfide stress. This disulfide stress could be monitored by redox sensitive fluorescence protein indicators.

For development of a homogeneous cell based assay system one would have to engineer a cell line in such a way that the cell line expresses both, a cyclooxygenase and a redox sensitive fluorescent GFP protein variant in parallel. Both proteins should be expressed stably in an inducible form.

If COX inhibitors are present in the test sample, a diminished lipid peroxide generation will result in a decreased roGFP fluorescence signal relative to the control.

4. Conclusions

As a basis for the development of an effect-based strategy to biomonitor pharmaceuticals, a literature review has been conducted which revealed the necessity to develop mode of actionbased biotests for routine monitoring of distinct pharmaceutical classes. In this context, a strategy to monitor entire classes of pharmaceuticals with the same mode of action is given preference because a monitoring programme comprising different biotests for each single substance would cause unrealistically high costs, and a monitoring programme which focusses on just a few lead substances would drastically underestimate the risk exerted by a plethora of pharmaceuticals excluded from analysis. Consequently, priority should be given to develop effect-oriented *in-vitro* tests for pharmaceutical classes and, within those, to analgesics, βblockers, and antibiotics. Such *in-vitro* assays need to be evaluated with *in-vivo* test systems in parallel. In view to incorporate living organisms here, the literature review revealed ciliate, fish, and mollusk species to exhibit particular sensitivity to pharmaceuticals. Most sensitive endpoints were behavior (chemotaxis), vitellogenin synthesis, growth, reproduction, histological responses, biochemical stress markers, changes in gene expression profiles, receptor binding, and, with reservation, the heart rate. Numerous pharmaceuticals have also been tested for their impact on parameters that directly influence population development: reproduction, fecundity, ontogeny, mating patterns. Here, most significance has been assigned to pharmaceuticals with endocrine action. In contrast to single substances, effects of mixtures of pharmaceuticals can hardly be assessed reliably, since data are not consistent. Consequently, mode of action-based biotests are necessary to decipher the contribution of single substances to the toxicity exerted by a mixture of pharmaceuticals. Endpoints which may be candidates to track mode of action-specific effects of pharmaceuticals are receptor affinity, vitellogenin induction (for estrogenic hormones), specific induction of gene expression, specific repression of enzymes (e.g. COX) or the formation of biochemical secondary products (e.g. lipid peroxides). Overall, the constructed database represents an extensive compilation of recently generated data on pharmaceutical effects, which will help researchers orienting in this quickly growing field. Besides the identification of certain promising test species and effect endpoints, there is growing evidence that several pharmaceuticals bear the potential to exhibit effects at environmentally relevant concentrations. They may therefore pose serious risks towards aquatic ecosystems and further studies are urgently needed.

The evaluation also showed that researchers need to put a higher effort into improving the reliability of their reported data; this includes a conclusive experimental design as well as a comprehensibly documentation and interpretation of the results.

5. Identification of shortcomings

The literature review generally revealed a strong heterogeneity of data, frequently resulting in just a single dataset for a species. Furthermore, in fish, most studies have been conducted with model species or with species of only local relevance. Consequently, single studies providing numerous data for a large number of chemicals which have been conducted with a single test organism in a single test run will bias the empirical evidence, particularly if the reliability of this test or study is in question.

In view to the relevance of data for Germany, a major shortcoming is the scarcity of data on ecologically relevant invertebrates (gammarids or biota of the sediment). Thus, the sensitivity of key species of home waters cannot be reliably assessed. Generally, data on sediment toxicity are limited in number. Only few publications report on studies that have used the same endpoint in the same test organism for different pharmaceuticals, and mode of action-specific endpoints are rarely used both in studies on single substances and mixtures. Quite often, chemical analytics supplementing biological tests is missing, and only about 70% of publication reviewed met the reliability criteria of Wright-Walters et al. (2011) to a sufficient extent. In general, a mode of action-based *in vitro* test for non-hormonal pharmaceuticals does not exist.

6. Future perspectives in the biomonitoring of pharmaceuticals

(1) We suggest to coin biotests that are based on mode of action-specific mechanisms and thus are specific for pharmaceutical classes and can be implemented in monitoring programs. Advantages of such biotests would be

- the integration of overall chemicals belonging to an mode of action-specific class of pharmaceuticals, irrespective of their accessibility by chemical analytics which can be limited by constraints posed by methodological detection limits, laboratory capacities, or budgetary limits,
- a pre-adaptation for the monitoring of future pharmaceuticals that exhibit the same mode of action as those the test has been developed for,
- and the integration of combinatory effects of mixtures of pharmaceuticals.

(2) Suitable prototypes implementing this idea would be *in vitro* tests for analgesics, like NSAIDs, and *in vitro* tests for β-blockers.

(3) The development of these *in vitro* biotests must go in line with *in vivo* experiments on ecologically relevant species which represent water and sediment biota in order to validate the sensitivity of the novel *in vitro* tests and to "ecologically calibrate" their signals. In these *in vivo* studies, identical endpoints shall be investigated in the laboratory and in field-relevant exposure systems, both for single pharmaceuticals and their mixtures. This strategy will provide necessary information regarding

- the relevance of *in vitro* test signals for the situation *in vivo*,
- the necessity to artificially concentrate water samples,
- the relevance of laboratory studies for the field situation,
- the significance of mixture toxicity, and
- differences in the toxicity of pharmaceuticals to water- and sediment-living biota.

These suggestions are completely in line with the postulations of Ankley et al. (2007) who emphasize the impotance of mode of action-based studies for pharmaceutical monitoring, and of Brausch et al. (2012) [6] who stress the necessity of

"...(a) chronic toxicity data for individual pharmaceuticals to benthic invertebrates,

including bivalves, and fish is lacking;

(b) Effects of pharmaceuticals on threatened or endangered species, which warrant protection at the individual level of biological organization;

(c) MOA-based studies, in which biochemical and histological alterations are investigated or studies in which genetic alterations are monitored in response to long-term pharmaceutical exposure;

(d) Techniques capable of detecting sensitive endpoints in aquatic organisms, such as in vitro and computational toxicology, for prioritizing chemicals and pathways for future studies;

(e) Data on complex mixtures of pharmaceuticals that found in WWTP effluents"

7 References

Part 1:

- [1] Bergmann, A., Fohrmann, R., Weber, F.A. (2011): Zusammenstellung von Monitoringdaten zu Umweltkonzentrationen von Arzneimitteln. UBA Texte 66, 2011.
- [2] Wright-Walters, M., Volz, C., Talbott, E., Davis, D. (2011): An updated weight of evidence approach to the aquatic hazard assessment of Bisphenol A and the derivation a new predicted no effect concentration (Pnec) using a non-parametric methodology. Sci. Tot. Env. 409: 676–685.
- [3] Lang, J., Kohidai, L. (2012): Effects of the aquatic contaminant human pharmaceuticals and their mixtures on the proliferation and migratory responses of the bioindicator freshwater ciliate Tetrahymena. Chemosphere 89, 592-601.
- [4] Fent, K. Weston; A.A., Caminada, D. (2006): Ecotoxicology of human pharmaceuticals.Aquat Toxicol. 76(2):122-59.
- [5] Brauch, H.-J. (2011): Organische Spurenstoffe in Gewässern. Vorkommen und Bewertung. Gwf-Wasser/Abwasser, 12, 1206-1211
- [6] Brausch, J.M., Connors, K.A., Brooks, B.W., Rand, G.M. (2012): Human Pharmaceuticals in the aquatic environment: A review of recent toxicological studies and considerations for toxicity testing. in: Whitacre, D.M. (Ed.), Reviews of Environmental Contamination and Toxicology, Vol 218. Springer, New York, pp. 1-99.
- [7] Eggeling, C., Brand, L., Ullmann, D., Jäger, S. (2003): Highly sensitive fluorescence detection technology currently available for HTS. Drug Discov Today. 8(14): 632-41.
- [8] Escher BI, Bramaz N, Eggen RIL, Richter M (2005): In vitro assessment of modes of toxic action of pharmaceuticals in aquatic life. Environ Sci Technol 39:3090–3100
- [9] Galus, M., Jeyaranjaan, J., Smith, E., Li, H., Metcalfe, C., Wilson, J.Y. (2013a): Chronic effects of exposure to a pharmaceutical mixture and municipal wastewater in zebrafish. *Aquatic toxicology (Amsterdam, Netherlands)* 132-133, 212-222.
- [10] Galus, M., Kirischian, N., Higgins, S., Purdy, J., Chow, J., Rangaranjan, S., Li, H.X., Metcalfe, C., Wilson, J.Y. (2013b): Chronic, low concentration exposure to pharmaceuticals impacts multiple organ systems in zebrafish. *Aquat. Toxicol.* 132, 200-211.

- [11] Franzellitti, S., Buratti, S., Valbonesi, P., Capuzzo, A., Fabbri, E. (2011): The beta-blocker propranolol affects cAMP-dependent signalling and induces the stress response in Mediterranean mussels, Mytilus galloprovincialis. *Aquat. Toxicol.* 101, 299-308.
- [12] Franzellitti, S., Buratti, S., Valbonesi, P., Fabbri, E. (2013): The mode of action (MOA) approach reveals interactive effects of environmental pharmaceuticals on Mytilus galloprovincialis. *Aquat. Toxicol.* 140, 249-256.
- [13] Parolini, M., Pedriali, A., Binelli, A. (2013): Application of a Biomarker Response Index for Ranking the Toxicity of Five Pharmaceutical and Personal Care Products (PPCPs) to the Bivalve Dreissena polymorpha. *Arch. Environ. Contam. Toxicol.* 64, 439-447.
- [14] Fairchild, W., Doe, K., Jackman, P., Akaishi, F., Arsenault, J., Aube, J., Losier, M., Cook, A. (2011): Acute and chronic toxicity of two formulations of the pyrethroid pesticide deltamethrin to amphipods, sand shrimp and lobster larvae (PO). *Canadian Technical Report* of Fisheries and Aquatic Sciences 2949, 64-65.
- [15] Antunes, S.C., Freitas, R., Figueira, E., Goncalves, F., Nunes, B. (2013): Biochemical effects of acetaminophen in aquatic species: edible clams Venerupis decussata and Venerupis philippinarum. *Environ. Sci. Pollut. Res.* 20, 6658-6666.
- [16] Backhaus, T., Porsbring, T., Arrhenius, A., Brosche, S., Johansson, P., Blanck, H. (2011): Single-substance and mixture toxicity of five pharmaceuticals and personal care products to marine periphyton communities. *Environ. Toxicol. Chem.* 30, 2030-2040.
- [17] Bajet, C.M., Kumar, A., Calingacion, M.N., Narvacan, T.C. (2012): Toxicological assessment of pesticides used in the Pagsanjan-Lumban catchment to selected non-target aquatic organisms in Laguna Lake, Philippines. *Agric. Water Manage.* 106, 42-49.
- [18] Benstead, R.S., Baynes, A., Casey, D., Routledge, E.J., Jobling, S. (2011): 17 beta-Oestradiol may prolong reproduction in seasonally breeding freshwater gastropod mollusks. *Aquat. Toxicol.* 101, 326-334.
- [19] Boltes, K., Rosal, R., Garcia-Calvo, E. (2012): Toxicity of mixtures of perfluorooctane sulphonic acid with chlorinated chemicals and lipid regulators. *Chemosphere* 86, 24-29.
- [20] Boonstra, H., Reichman, E.P., van den Brink, P.J. (2011): Effects of the Veterinary Pharmaceutical Ivermectin in Indoor Aquatic Microcosms. *Arch. Environ. Contam. Toxicol.* 60, 77-89.
- [21] Chandra, K., Bosker, T., Hogan, N., Lister, A., MacLatchy, D., Currie, S. (2012): Sustained high temperature increases the vitellogenin response to 17 alpha-ethynylestradiol in mummichog (*Fundulus heteroclitus*). *Aquat. Toxicol.* 118, 130-140.
- [22] Chen, X., Zhou, Y., Yang, D., Zhao, H., Wang, L., Yuan, X. (2012): CYP4 mRNA expression in marine polychaete Perinereis aibuhitensis in response to petroleum hydrocarbon and deltamethrin. *Marine Pollution Bulletin* 64, 1782-1788.
- [23] Claessens, M., Vanhaecke, L., Wille, K., Janssen, C.R. (2013): Emerging contaminants in Belgian marine waters: Single toxicant and mixture risks of pharmaceuticals. *Marine pollution bulletin* 71, 41-50.
- [24] Contardo-Jara, V., Lorenz, C., Pflugmacher, S., Nuetzmann, G., Kloas, W., Wiegand, C. (2011): Exposure to human pharmaceuticals Carbamazepine, Ibuprofen and Bezafibrate causes molecular effects in Dreissena polymorpha. *Aquat. Toxicol.* 105, 428-437.

- [25] Doyle, M.A., Bosker, T., Martyniuk, C.J., Maclatchy, D.L., Munkittrick, K.R. (2013): The effects of 17-alpha-ethinylestradiol (EE2) on molecular signalling cascades in mummichog (*Fundulus heteroclitus*). *Aquatic toxicology (Amsterdam, Netherlands)* 134-135, 34-46.
- [26] Feito, R., Valcarcel, Y., Catala, M. (2012): Biomarker assessment of toxicity with miniaturised bioassays: diclofenac as a case study. *Ecotoxicology* 21, 289-296.
- [27] Feito, R., Valcarcel, Y., Catala, M. (2013): Preliminary data suggest that venlafaxine environmental concentrations could be toxic to plants. *Chemosphere* 90, 2065-2069.
- [28] Finn, J., Hui, M., Li, V., Lorenzi, V., de la Paz, N., Cheng, S.H., Lai-Chan, L., Schlenk, D. (2012): Effects of propranolol on heart rate and development in Japanese medaka (*Oryzias latipes*) and zebrafish (*Danio rerio*). *Aquat. Toxicol.* 122, 214-221.
- [29] Fong, P.P., Hoy, C.M. (2012): Antidepressants (venlafaxine and citalopram) cause foot detachment from the substrate in freshwater snails at environmentally relevant concentrations. *Marine and Freshwater Behaviour and Physiology* 45, 145-153.
- [30] Gagne, F., Andre, C., Fortier, M., Fournier, M. (2012): Immunotoxic potential of aeration lagoon effluents for the treatment of domestic and hospital wastewaters in the freshwater mussel Elliptio complanata. *J. Environ. Sci.* 24, 781-789.
- [31] Gonzalez-Rey, M., Bebianno, M.J. (2011): Non-steroidal anti-inflammatory drug (NSAID) ibuprofen distresses antioxidant defense system in mussel Mytilus galloprovincialis gills. *Aquat. Toxicol.* 105, 264-269.
- [32] Gust, M., Gelinas, M., Fortier, M., Fournier, M., Gagne, F. (2012): In vitro immunotoxicity of environmentally representative antibiotics to the freshwater mussel Elliptio complanata. *Environ. Pollut.* 169, 50-58.
- [33] Gust, M., Fortier, M., Garric, J., Fournier, M., Gagne, F. (2013): Effects of short-term exposure to environmentally relevant concentrations of different pharmaceutical mixtures on the immune response of the pond snail Lymnaea stagnalis. *Sci. Total Environ.* 445, 210-218.
- [34] Hallgren, S., Volkova, K., Reyhanian, N., Olsen, K.H., Hallstrom, I.P. (2011): Anxiogenic behaviour induced by 17 alpha-ethynylestradiol in male guppies (*Poecilia reticulata*). *Fish Physiol. Biochem.* 37, 911-918.
- [35] Hallgren, P., Sorita, Z., Berglund, O., Persson, A. (2012): Effects of 17 alpha-ethinylestradiol on individual life-history parameters and estimated population growth rates of the freshwater gastropods Radix balthica and Bithynia tentaculata. *Ecotoxicology* 21, 803-810.
- [36] Hillis, D.G., Fletcher, J., Solomon, K.R., Sibley, P.K. (2011): Effects of ten antibiotics on seed germination and root elongation in three plant species. *Arch. Environ. Contam. Toxicol.* 60, 220-232.
- [37] Hoffmann, F., Kloas, W. (2012): Estrogens Can Disrupt Amphibian Mating Behavior. *PloS* one 7.
- [38] Huynh Thi, T., Silvestre, F., De Meulder, B., Thome, J.-P., Nguyen Thanh, P., Kestemont, P. (2012): Combined effects of deltamethrin, temperature and salinity on oxidative stress biomarkers and acetylcholinesterase activity in the black tiger shrimp (Penaeus monodon). *Chemosphere* 86, 83-91.
- [39] Ings, J.S., George, N., Peter, M.C.S., Servos, M.R., Vijayan, M.M. (2012): Venlafaxine and atenolol disrupt epinephrine-stimulated glucose production in rainbow trout hepatocytes. *Aquat. Toxicol.* 106, 48-55.

- [40] Johns, S.M., Denslow, N.D., Kane, M.D., Watanabe, K.H., Orlando, E.F., Sepulveda, M.S. (2011): Effects of estrogens and antiestrogens on gene expression of fathead minnow (*Pimephales promelas*) early life stages. *Environ. Toxicol.* 26, 195-206.
- [41] Kaptaner, B., Unal, G. (2011): Effects of 17 alpha-ethynylestradiol and nonylphenol on liver and gonadal apoptosis and histopathology in *Chalcalburnus tarichi. Environ. Toxicol.* 26, 610-622.
- [42] Lange, A., Katsu, Y., Miyagawa, S., Ogino, Y., Urushitani, H., Kobayashi, T., Hirai, T., Shears, J.A., Nagae, M., Yamamoto, J., Ohnishi, Y., Oka, T., Tatarazako, N., Ohta, Y., Tyler, C.R., Iguchi, T. (2012): Comparative responsiveness to natural and synthetic estrogens of fish species commonly used in the laboratory and field monitoring. *Aquat. Toxicol.* 109, 250-258.
- [43] Lawrence, J.R., Zhu, B., Swerhone, G.D.W., Roy, J., Tumber, V., Waiser, M.J., Topp, E., Korber, D.R. (2012): Molecular and microscopic assessment of the effects of caffeine, acetaminophen, diclofenac, and their mixtures on river biofilm communities. *Environ. Toxicol. Chem.* 31, 508-517.
- [44] Lei, B., Kang, J., Yu, Y., Zha, J., Li, W., Wang, Z. (2013): Beta-estradiol 17-valerate affects embryonic development and sexual differentiation in Japanese medaka (*Oryzias latipes*). *Aquatic toxicology (Amsterdam, Netherlands)* 134-135, 128-134.
- [45] Liu, B.Y., Liu, W.Q., Nie, X.P., Guan, C., Yang, Y.F., Wang, Z.H., Liao, W. (2011): Growth response and toxic effects of three antibiotics on *Selenastrum capricornutum* evaluated by photosynthetic rate and chlorophyll biosynthesis. *J. Environ. Sci.* 23, 1558-1563.
- [46] Liu, B.Y., Nie, X.P., Liu, W.Q., Snoeijs, P., Guan, C., Tsui, M.T.K. (2011): Toxic effects of erythromycin, ciprofloxacin and sulfamethoxazole on photosynthetic apparatus in *Selenastrum capricornutum. Ecotoxicology and Environmental Safety* 74, 1027-1035.
- [47]Madureira, T.V., Cruzeiro, C., Rocha, M.J., Rocha, E. (2011): The toxicity potential of pharmaceuticals found in the Douro River estuary (Portugal)-Experimental assessment using a zebrafish embryo test. *Environmental Toxicology and Pharmacology* 32, 212-217.
- [48] Meina, E.G., Lister, A., Bosker, T., Servos, M., Munkittrick, K., MacLatchy, D. (2013): Effects of 17 alpha-ethinylestradiol (EE2) on reproductive endocrine status in mummichog (*Fundulus heteroclitus*) under differing salinity and temperature conditions. *Aquat. Toxicol.* 134, 92-103.
- [49] Notch, E.G., Mayer, G.D. (2013): Impact of environmental estrogens on nucleotide excision repair gene expression in embryonic zebrafish. *Comparative biochemistry and physiology. Toxicology & pharmacology : CBP* 157, 361-365.
- [50] Oliveira, C., Almeida, J., Guilhermino, L., Soares, A.M.V.M., Gravato, C. (2012): Acute effects of deltamethrin on swimming velocity and biomarkers of the common prawn Palaemon serratus. *Aquat. Toxicol.* 124, 209-216.
- [51] Ozdemir, A., Duran, M., Sen, A. (2011): Potential Use of the Oligochaete Limnodrilus profundicola V., as a Bioindicator of Contaminant Exposure. *Environ. Toxicol.* 26, 37-44.
- [52] Parolini, M., Binelli, A., Provini, A. (2011a): Chronic effects induced by ibuprofen on the freshwater bivalve Dreissena polymorpha. *Ecotoxicology and Environmental Safety* 74, 1586-1594.
- [53] Parolini, M., Quinn, B., Binelli, A., Provini, A. (2011b): Cytotoxicity assessment of four pharmaceutical compounds on the zebra mussel (Dreissena polymorpha) haemocytes, gill and digestive gland primary cell cultures. *Chemosphere* 84, 91-100.

- [54] Ragugnetti, M., Adams, M.L., Guimaraes, A.T.B., Sponchiado, G., de Vasconcelos, E.C., de Oliveira, C.M.R. (2011): Ibuprofen genotoxicity in aquatic environment: An experimental model using *Oreochromis niloticus. Water Air Soil Pollut.* 218, 361-364.
- [55] Reyhanian, N., Volkova, K., Hallgren, S., Bollner, T., Olsson, P.E., Olsen, H., Hallstrom, I.P.
 (2011): 17 alpha-Ethinyl estradiol affects anxiety and shoaling behavior in adult male zebra fish (*Danio rerio*). *Aquat. Toxicol.* 105, 41-48.
- [56] Ribeiro, C., Urbatzka, R., Castro, L.F.C., Carrola, J., Fontainhas-Fernandes, A., Monteiro, R.A.F., Rocha, E., Rocha, M.J. (2012): In vitro exposure of Nile tilapia (*Oreochromis niloticus*) testis to estrogenic endocrine disrupting chemicals: mRNA expression of genes encoding steroidogenic enzymes. *Toxicol. Mech. Methods* 22, 47-53.
- [57] Rocco, L., Frenzilli, G., Zito, G., Archimandritis, A., Peluso, C., Stingo, V. (2012a): Genotoxic effects in fish induced by pharmacological agents present in the sewage of some Italian water-treatment plants. *Environ. Toxicol.* 27, 18-25.
- [58] Rocco, L., Peluso, C., Stingo, V. (2012b): Micronucleus test and comet assay for the evaluation of zebrafish genomic damage induced by erythromycin and lincomycin. *Environ. Toxicol.* 27, 598-604.
- [59] Saravanan, M., Karthika, S., Malarvizhi, A., Ramesh, M. (2011): Ecotoxicological impacts of clofibric acid and diclofenac in common carp (*Cyprinus carpio*) fingerlings: Hematological, biochemical, ionoregulatory and enzymological responses. *J. Hazard. Mater.* 195, 188-194.
- [60] Sarria, M.P., Santos, M.M., Reis-Henriques, M.A., Vieira, N.M., Monteiro, N.M. (2011): The unpredictable effects of mixtures of androgenic and estrogenic chemicals on fish early life. *Environ. Int.* 37, 418-424.
- [61] Shen, M.-F., Kumar, A., Ding, S.-Y., Grocke, S. (2012): Comparative study on the toxicity of pyrethroids, alpha-cypermethrin and deltamethrin to Ceriodaphnia dubia. *Ecotoxicology and Environmental Safety* 78, 9-13.
- [62] Shi, H.H., Sun, Z., Liu, Z., Xue, Y.G. (2012): Effects of clotrimazole and amiodarone on early development of amphibian (*Xenopus tropicalis*). *Toxicol Environ. Chem.* 94, 128-135.
- [63] Silva, P., Rocha, M.J., Cruzeiro, C., Malhao, F., Reis, B., Urbatzka, R., Monteiro, R.A.F., Rocha, E. (2012): Testing the effects of ethinylestradiol and of an environmentally relevant mixture of xenoestrogens as found in the Douro River (Portugal) on the maturation of fish gonads-A stereological study using the zebrafish (*Danio rerio*) as model. *Aquat. Toxicol.* 124, 1-10.
- [64] Skolness, S.Y., Durhan, E.J., Jensen, K.M., Kahl, M.D., Makynen, E.A., Villeneuve, D.L., Ankley, G.T. (2012): Effects of gemfibrozil on lipid metabolism, steroidogenesis, and reproduction in the fathead minnow (*Pimephales promelas*). *Environ. Toxicol. Chem.* 31, 2615-2624.
- [65] Sponchiado, G., Reynaldo, E., de Andrade, A.C.B., de Vasconcelos, E.C., Adam, M.L., de Oliveira, C.M.R. (2011): Genotoxic effects in erythrocytes of *Oreochromis niloticus* exposed to nanograms-per-liter concentration of 17 beta-estradiol (E-2): An assessment using micronucleus test and comet assay. *Water Air Soil Pollut.* 218, 353-360.
- [66] Stange, D., Oehlmann, J. (2012a): Identification of oestrogen-responsive transcripts in Potamopyrgus antipodarum. *Journal of Molluscan Studies* 78, 337-342.
- [67] Stange, D., Sieratowicz, A., Horres, R., Oehlmann, J. (2012b): Freshwater mudsnail (Potamopyrgus antipodarum) estrogen receptor: Identification and expression analysis under exposure to (xeno-)hormones. *Ecotoxicology and Environmental Safety* 75, 94-101.

- [68] Thomas, M.A., Joshi, P.P., Klaper, R.D. (2012): Gene-class analysis of expression patterns induced by psychoactive pharmaceutical exposure in fathead minnow (*Pimephales promelas*) indicates induction of neuronal systems. *Comp. Biochem. Physiol. C-Toxicol. Pharmacol.* 155, 109-120.
- [69] Toumi, H., Boumaiza, M., Millet, M., Radetski, C.M., Felten, V., Fouque, C., Ferard, J.F. (2013): Effects of deltamethrin (pyrethroid insecticide) on growth, reproduction, embryonic development and sex differentiation in two strains of Daphnia magna (Crustacea, Cladocera). *Sci. Total Environ.* 458, 47-53.
- [70] van Leeuwen, J.S., Unlu, B., Vermeulen, N.P.E., Vos, J.C. (2012): Differential involvement of mitochondrial dysfunction, cytochrome P450 activity, and active transport in the toxicity of structurally related NSAIDs. *Toxicol. Vitro* 26, 197-205.
- [71] Veach, A., Bernot, M.J., Mitchell, J.K. (2012): The influence of six pharmaceuticals on freshwater sediment microbial growth incubated at different temperatures and UV exposures. *Biodegradation* 23, 497-507.
- [72] Wang, S.Y., Gunsch, C.K. (2011): Effects of selected pharmaceutically active compounds on the ammonia oxidizing bacterium *Nitrosomonas europaea. Chemosphere* 82, 565-572.
- [73] Wang, H.P., Wang, J.J., Wu, T.T., Qin, F., Hu, X.Q., Wang, L.H., Wang, Z.Z. (2011): Molecular characterization of estrogen receptor genes in *Gobiocypris rarus* and their expression upon endocrine disrupting chemicals exposure in juveniles. *Aquat. Toxicol.* 101, 276-287.
- [74] Wu, N., Wei, H., Shen, H., Wu, T.T., Guo, M. (2012): Acute toxic effects of deltamethrinon red swampcrayfish, procambarus clarkia (decapoda, cambaridae). *Crustaceana* 85, 993-1005.
- [75] Yan, Z.H., Lu, G.H., Yang, X.F. (2013): Single and combined effects of estrone and 17 betaestradiol on male goldfish. *Biomed. Environ. Sci.* 26, 176-184.
- [76] Yergeau, E., Sanschagrin, S., Waiser, M.J., Lawrence, J.R., Greer, C.W. (2012): Sub-inhibitory concentrations of different pharmaceutical products affect the meta-transcriptome of river biofilm communities cultivated in rotating annular reactors. *Environ. Microbiol. Rep.* 4, 350-359.
- [77] Yonar, S.M., Sakin, F., Yonar, M.E., Ispir, U., Kirici, M. (2011): Oxidative stress biomarkers of exposure to deltamethrin in rainbow trout fry (*Oncorhynchus mykiss*). *Fresenius Environ. Bull.* 20, 1931-1935.
- [78] Zhang, X.Y., Gong, Z.Y. (2013): Fluorescent transgenic zebrafish Tg(nkx2.2a:mEGFP) provides a highly sensitive monitoring tool for neurotoxins. *PloS one* 8.
- [79] Zhang, Y.Y., Wang, H.P., Qin, F., Liu, S.Z., Wu, T.T., Li, M., Xu, P., Zhang, X., Wang, X.Q., Hu, G.J., Yuan, C., Wang, Z.Z. (2012c): Molecular characterization of estrogen receptor genes in loach *Paramisgurnus dabryanus* and their expression upon 17 alpha-ethinylestradiol exposure in juveniles. *Gen. Comp. Endocrinol.* 178, 194-205.
- [80] Zhang, W., Zhang, M., Lin, K.F., Sun, W.F., Xiong, B., Guo, M.J., Cui, X.H., Fu, R.B. (2012a): Eco-toxicological effect of carbamazepine on *Scenedesmus obliquus* and *Chlorella pyrenoidosa. Environmental Toxicology and Pharmacology* 33, 344-352.

Part 2

- [1] Burns, S., Travers, J., Collins, I., et al. (2006): Identification of small-molecule inhibitors of protein kinase B (PKB/AKT) in an AlphaScreenTM high-throughput screen. Journal of Biomolecular Screening, 11(7), 822–827.
- [2] Sudo, K., Yamaji, K., Kawamura, K., et al. (2005): Highthroughput screening of low molecular weight NS3-NS4A protease inhibitors using a fluorescence resonance energy transfer substrate. Antiviral Chemistry & Chemotherapy, 16(6), 385–392.
- [3] Swaney, S., McCroskey, M., Shinabarger, D., et al. (2006): Characterization of a high-throughput screening assay for inhibitors of elongation factor P and ribosomal peptidyl transferase activity. Journal of Biomolecular Screening, 11(7), 736–742.
- [4] Allen, M., Reeves, J., & Mellor, G. (2000): High throughput fluorescence polarization: A homogeneous alternative to radioligand binding for cell surface receptors. Journal of Biomolecular Screening, 5(2), 63–69.
- [5] Parker, G. J., Law, T. L., Lenoch, F. J., et al. (2000): Development of high throughput screening assays using fluorescence polarization: Nuclear receptor-ligand-binding and kinase/phosphatase assays. Journal of Biomolecular Screening, 5(2), 77–88.
- [6] Kenny, C. H., Ding, W., Kelleher, K., et al. (2003): Development of a fluorescence polarization assay to screen for inhibitors of the FtsZ/ZipA interaction. Analytical Biochemistry, 323(2), 224–233.
- [7] Lohse MJ, Bünemann M, Hoffmann C, Vilardaga JP, Nikolaev VO. (2007): Monitoring receptor signaling by intramolecular FRET. Curr Opin Pharmacol. Oct;7(5):547-53.
- [8]Kariv, I., Stevens, M. E., Behrens, D. L., et al. (1999): High throughput quantitation of cAMP production mediated by activation of seven transmembrane domain receptors. Journal of Biomolecular Screening, 4(1), 27–32.
- [9] Li, X., Shen, F., Zhang, Y., et al. (2007): Functional characterization of cell lines for highthroughput screening of human neuromedin U receptor subtype 2 specific agonists using a luciferase reporter gene assay. European Journal of Pharmaceutics and Biopharmaceutics.
- [10] Beck, V., Pfitscher, A., & Jungbauer, A. (2005): GFP-reporter for a high throughput assay to monitor estrogenic compounds. Journal of Biochemical and Biophysical Methods, 64(1), 19– 37.
- [11] Wu S, Liu B. (2005): Application of scintillation proximity assay in drug discovery. BioDrugs. 19(6):383-92.
- [12] McDonald OB, Chen WJ, Ellis B, Hoffman C, Overton L, Rink M, Smith A, Marshall CJ, Wood ER (1999): A scintillation proximity-assay for the Raf/MEK/ERK kinase cascade: highthroughput screening and identification of selective enzyme inhibitors, Anal Biochem, 268:318-329.
- [13] Park YW, Cummings RT, Wu L, Zheng S, Cameron PM, Woods A, Zaller DM, Marcy AI, Hermes JD (1999): Homogeneous proximity tyrosine kinase assays: scintillation proximity assay versus homogeneous time-resolved fluorescence. Anal Biochem, 269:94-104.
- [14] Nare B, Allocco JJ, Kuningas R, Galuska S, Myers RW, Bednarek MA, Schmatz DM (1999): Development of a scintillation proximity assay for histone deacetylase using a biotinylated peptide derived from histone H4, Anal Biochem, 267:390-396.

- [15] Kahl SD, Hubbard FR, Sittampalam GS, Zock JM (1997): Validation of a high throughput scintillation proximity assay for 5-hydroxytryptamine(1E) receptor binding activity. J Biomol Screen, 2:33-40.
- [16] Kariv I, Stevens ME, Behrens DL, Oldenburg KR (1999): High throughput quantitation of cAMP production mediated by activation of seven transmembrane domain receptors. J Biomol Screen, 4:27-32.
- [17] Bosse R, Garlick R, Brown B, Menard L (1998): Development of nonseparation binding and functional assays for G protein-coupled receptors for high throughput screening: pharmacological characterization of the immobilized CCR5 receptor on FlashPlate[™]. J Biomol Screen, 3:285-292
- [18] Owicki, J.C. (2000): Fluorescence polarization and anisotropy in high throughput screening: perspectives and primer. J. Biomol. Screen. 5, 297–306
- [19] Jameson DM, Ross JA. (2010): Fluorescence polarization/anisotropy in diagnostics and imaging, Chem Rev. May 12;110(5):2685-708.
- [20] Lakowicz, J. (2006): Principles of Fluorescence Spectroscopy (Third Edition) Springer Science+Business Media
- [21] Morita I, Smith WL, DeWitt DL, Schindler M. (1995): Expression-activity profiles of cells transfected with prostaglandin endoperoxide H synthase measured by quantitative fluorescence microscopy, Biochemistry. May 30;34(21):7194-9.
- [22] Medintz IL, Anderson GP, Lassman ME, Goldman ER, Bettencourt LA, Mauro JM. (2004): General strategy for biosensor design and construction employing multifunctional surfacetethered components, Anal Chem 76:5620–5629.
- [23] Pfleger, K. D.; Eidne, K. A. (2006): Illuminating insights into protein-protein interactions using bioluminescence resonance energy transfer (BRET), Nat. Methods, 3, 165.
- [24] Bacart, J.; Corbel, C.; Jockers, R.; Bach, S.; Couturier, C. (2008): The BRET technology and its application to screening assays, Biotechnol. J., 3, 311.
- [25] Hoffmann, C.; Gaietta, G.; Bunemann, M.; Adams, S. R.; Oberdorff-Maass, S.; Behr, B.; Vilardaga, J. P.; Tsien, R. Y.; Ellisman (2005): A FlAsH-based FRET approach to determine G protein–coupled receptor activation in living cells, J. Nat. Methods 2005, 2, 171.
- [26] Balla, T. (2009): Green light to illuminate signal transduction events, Trends Cell Biol., 19, 575.
- [27] Chalfie, M.; Tu, Y.; Euskirchen, G.; Ward, W. W.; Prasher, D. C. (1994): Green Fluorescent Protein as a Marker for Gene Expression, Science, 263, 802
- [28] Verkhusha, V. V.; Kuznetsova, I. M.; Stepanenko, O. V.; Zaraisky, A. G.; Shavlovsky, M. M.; Turoverov, K. K.; Uversky, V. N. (2003): High Stability of Discosoma DsRed As Compared to Aequorea EGFP, Biochemistry , 42, 7879.
- [29] Li, X.; Zhao, X.; Fang, Y.; Jiang, X.; Duong, T.; Fan, C.; Huang, C. C.; Kain, S. R. (1998):Generation of Destabilized Green Fluorescent Protein as a Transcription Reporter, J. Biol. Chem., 273, 34970.
- [30] Jayaraman, S.; Haggie, P.; Wachter, R. M.; Remington, S. J.; Verkman, A. S. (2000): Mechanism and Cellular Applications of a Green Fluorescent Protein-based Halide Sensor J. Biol. Chem., 275, 6047.
- [31] Llopis, J.; McCaffery, J. M.; Miyawaki, A.; Farquhar, M. G.; Tsien, R. Y. (1998): Measurement of cytosolic, mitochondrial, and Golgi pH in single living cells with green fluorescent proteins Proc. Natl. Acad. Sci. U.S.A., 95, 6803.
- [32] Cannon MB, Remington SJ. (2006): Re-engineering redox-sensitive green fluorescent protein for improved response rate Protein Sci. 2006 Jan;15(1):45-57.
- [33] Hanson, G. T.; Aggeler, R.; Oglesbee, D.; Cannon, M.; Capaldi, R. A.; Tsien, R. Y.; Remington, S.J. (2004): Investigating Mitochondrial Redox Potential with Redox-sensitive Green Fluorescent Protein Indicators J. Biol. Chem., 279, 13044.
- [34] Ostergaard, H.; Henriksen, A.; Hansen, F. G.; Winther, J. R. (2001): Shedding light on disulfide bond formation: engineering a redox switch in green fluorescent protein. EMBO J., 20, 5853.
- [35] Tian, L.; Hires, S. A.; Mao, T.; Huber, D.; Chiappe, M. E.; Chalasani, S. H.; Petreanu, L.; Akerboom, J.; McKinney, S. A.; Schreiter, L. E. R.; Bargmann, C. I.; Jayaraman, V.; Svoboda, K.; Looger, L. (2009): Imaging neural activity in worms, flies and mice with improved GCaMP calcium indicators Nat. Methods, 6, 875.
- [36] Nausch, L. W.; Ledoux, J.; Bonev, A. D.; Nelson, M. T.; Dostmann, W. R. (2008): Differential patterning of cGMP in vascular smooth muscle cells revealed by single GFP-linked biosensors Proc. Natl. Acad. Sci. U.S.A., 105, 365.
- [37] Belousov, V. V.; Fradkov, A. F.; Lukyanov, K. A.; Staroverov, D. B.; Shakhbazov, K. S.; Terskikh, A. V.; Lukyanov, S. (2006): Genetically encoded fluorescent indicator for intracellular hydrogen peroxide Nat. Methods, 3, 281.
- [38] Wang, W.; Fang, H.; Groom, L.; Cheng, A.; Zhang, W.; Liu, J.; Wang, X.; Li, K.; Han, P.; Zheng, M.; Yin, J.; Wang, W.; Mattson, M. P.; Kao, J. P.; Lakatta, E. G.; Sheu, S. S.; Ouyang, K.; Chen, J.; Dirksen, R. T.; Cheng, H. (2008): Superoxide Flashes in Single Mitochondria Cell, 134, 279.
- [39] Berg, J.; Hung, Y. P.; Yellen, G. (2009): A genetically encoded fluorescent reporter of ATP:ADP ratio Nat. Methods, 6, 161.
- [40] Baker, B. J.; Mutoh, H.; Dimitrov, D.; Akemann, W.; Perron, A.; Iwamoto, Y.; Jin, L.; Cohen, L. B.; Isacoff, E. Y.; Pieribone, V. A.; Hughes, T.; Knopfel, T. (2008): Genetically encoded fluorescent sensors of membrane potential Brain Cell Biol., 36, 53.
- [41] Lundby, A.; Mutoh, H.; Dimitrov, D.; Akemann, W.; Knopfel (2008): Engineering of a Genetically Encodable Fluorescent Voltage Sensor Exploiting Fast Ci-VSP Voltage-Sensing Movements T. PLoS One, 3, e2514.
- [42] Nikolaev, V. O.; Bunemann, M.; Hein, L.; Hannawacker, A.; Lohse, M. J. (2004): Novel Single Chain cAMP Sensors for Receptor-induced Signal Propagation J. Biol. Chem., 279, 37215.
- [43] Violin, J. D.; DiPilato, L. M.; Yildirim, N.; Elston, T. C.; Zhang, J.; Lefkowitz, R. J. (2008): β2-Adrenergic Receptor Signaling and Desensitization Elucidated by Quantitative Modeling of Real Time cAMP Dynamics J. Biol. Chem., 283, 2949.
- [44] DiPilato, L. M.; Cheng, X.; Zhang, J. (2004): Fluorescent indicators of cAMP and Epac activation reveal differential dynamics of cAMP signaling within discrete subcellular compartments Proc. Natl. Acad. Sci. U.S.A., 101, 16513.
- [45] Ponsioen, B.; Zhao, J.; Riedl, J.; Zwartkruis, F.; van der, K. G.;Zaccolo, M.; Moolenaar, W. H.; Bos, J. L.; Jalink, K. (2004): Detecting cAMP-induced Epac activation by fluorescence resonance energy transfer: Epac as a novel cAMP indicator EMBO Rep., 5, 1176.

- [46] Jiang, L. I.; Collins, J.; Davis, R.; Lin, K. M.; DeCamp, D.; Roach, T.; Hsueh, R.; Rebres, R. A.; Ross, E. M.; Taussig, R.; Fraser, I.; Sternweis, P. C. (2007): Use of a cAMP BRET Sensor to Characterize a Novel Regulation of cAMP by the Sphingosine 1-Phosphate/G13 Pathway J. Biol. Chem., 282, 10576
- [47] Masri, B.; Salahpour, A.; Didriksen, M.; Ghisi, V.; Beaulieu, J. M.; Gainetdinov, R. R.; Caron, M. G. (2008): Antagonism of dopamine D2 receptor/β-arrestin 2 interaction is a common property of clinically effective antipsychoticsProc. Natl. Acad. Sci. U.S.A., 105, 13656.
- [48] Lohse, M. J.; Nikolaev, V. O.; Hein, P.; Hoffmann, C.; Vilardaga, J. P.; Bunemann, M. (2008):Optical techniques to analyze real-time activation and signalling of G-protein-coupled receptors Trends Pharmacol. Sci., 29, 159.
- [49] Chachisvilis, M.; Zhang, Y. L.; Frangos, J. A. (2006): G protein-coupled receptors sense fluid shearstress in endothelial cells Proc. Natl. Acad. Sci. U.S.A. 2006, 103, 15463.
- [50] Vilardaga, J. P.; Steinmeyer, R.; Harms, G. S.; Lohse, M. (2005): Molecular basis of inverse agonism in a G protein–coupled receptor J. Nat. Chem. Biol., 1, 25.
- [51] Rochais F, Vilardaga JP, Nikolaev VO, Bünemann M, Lohse MJ, Engelhardt S. (2007): Realtime optical recording of beta1-adrenergic receptor activation reveals supersensitivity of the Arg389 variant to carvedilol. J Clin Invest. Jan;117(1):229-35.
- [52] Zhang, J.; Hupfeld, C. J.; Taylor, S. S.; Olefsky, J. M.; Tsien, R. Y. (2005): Insulin disrupts βadrenergic signalling to protein kinase A in adipocytes Nature, 437, 569.
- [53] Sato, M.; Nakajima, T.; Goto, M.; Umezawa, Y. (2006): Cell-Based Indicator to Visualize PicomolarDynamics of Nitric Oxide Release from Living Cells Anal. Chem., 78, 8175.
- [54] MacDonald ML, Lamerdin J, Owens S, Keon BH, Bilter GK, Shang Z, Huang Z, Yu H, Dias J, Minami T, et al. (2006): Identifying off-target effects and hidden phenotypes of drugs in human cells Nat Chem Biol 2:329–337.
- [55] Fan F, Wood KV. (2007): Bioluminescent assays for high-throughput screening. Assay Drug Dev Technol; 5: 127–36.
- [56] Brodde, O.E., and Michel, M.C. (1999): Adrenergic and muscarinic receptors in the human heart. Pharmacol. Rev. 51:651–690.
- [57] Bristow, M. (2003): Antiadrenergic therapy of chronic heart failure: surprises and new opportunities.Circulation. 107:1100–1102.
- [58] Binkowski BF, Butler BL, Stecha PF, Eggers CT, Otto P, Zimmerman K, Vidugiris G, Wood MG, Encell LP, Fan F, Wood KV. (2011): A luminescent biosensor with increased dynamic range for intracellular cAMP. ACS Chem Biol. Nov 18;6(11):1193-7.
- [59] Shan Q, Storm DR. (2010): Optimization of a cAMP response element signal pathway reporter system J Neurosci Methods Aug 15;191(1):21-5.
- [60] Rich TC, Tse TE, Rohan JG, Schaack J, Karpen JW. (2001): *In vivo* assessment of local phosphodiesterase activity using tailored cyclic nucleotide-gated channels as cAMP sensors. J Gen Physiol. 118:63–77.
- [61] Nikolaev, V.O.; Bunemann, M.; Schmitteckert, E.; Lohse, M.J.; Engelhardt, S. (2006): Cyclic AMP imaging in adult cardiac myocytes reveals far-reaching beta1-adrenergic but locally confined beta2-adrenergic receptor-mediated signaling. Circ. Res. 99, 1084–1091

- [62] Mitchell JA, Akarasereenont P, Thiemermann C, Flower RJ and Vane JR (1993): Selectivity of nonsteroidal anti-inflammatory drugs as inhibitors of constitutive and inducible cyclooxygenase. Proc Natl Acad Sci USA 90: 11693-11697
- [63] Lee CR, Bottone FG Jr, Krahn JM, Li L, Mohrenweiser HW, Cook ME, Petrovich RM, Bell DA, Eling TE, Zeldin DC. (2007): Identification and functional characterization of polymorphisms in human cyclooxygenase-1 (PTGS1). Pharmacogenet Genomics. Feb;17(2):145-60.
- [64] Liu W, Poole EM, Ulrich CM, Kulmacz RJ. (2012): Decreased cyclooxygenase inhibition by aspirin in polymorphic variants of human prostaglandin H synthase-1. Pharmacogenet Genomics Jul;22(7):525-37.
- [65] Kargman S, Wong E, Greig GM, Falgueyret JP, Cromlish W, Ethier D, Yergey JA, Riendeau D, Evans JF, Kennedy B, Tagari P, Francis DA, O'Neill GP. (1996): Mechanism of selective inhibition of human Prostaglandin G/H Synthase-1 and -2 in intact cells. Biochem Pharmacol. Oct 11;52(7): 1113-1125
- [66] Scholz S, Renner P, Belanger SE, Busquet F, Davi R, Demeneix BA, Denny JS, Léonard M, McMaster ME, Villeneuve DL, Embry MR. (2013): Alternatives to *in vivo* tests to detect endocrine disrupting chemicals (EDCs) in fish and amphibians–screening for estrogen, androgen and thyroid hormone disruption. Crit Rev Toxicol. Jan;43(1):45-72
- [67] Kulmacz RJ, Lands WEM (1983): Requirements for hydroperoxides by the cyclooxygenase and peroxidase activities of prostaglandin H synthase. Prostaglandins, 25:531-540
- [68] Björnstedt M, Hamberg M, Kumar S, Xue J, Holmgren A. (1995): Human thioredoxin reductase directly reduces lipid hydroperoxides by NADPH and selenocystine strongly stimulates the reaction via catalytically generated selenols. J Biol Chem. May 19;270(20):11761-4
- [69] Zaccolo, M.; De Giorgi, F.; Cho, C.Y.; Feng, L.; Knapp, T.; Negulescu, P.A.; Taylor, S.S.; Tsien, R.Y.; Pozzan, T., (2000): A genetically encoded, fluorescent indicator for cyclic AMP in living cells. Nat. Cell. Biol. 2, 25–29.
- [70] Norris, R.P.; Ratzan, W.J.; Freudzon, M.; Mehlmann, L.M.; Krall, J.; Movsesian, M.A.; Wang, H.; Ke, H.; Nikolaev, V.O.; Jaffe, L.A. (2009): Cyclic GMP from the surrounding somatic cells regulates cyclic AMP and meiosis in the mouse oocyte. Development 136, 1869–1878

Reviewed literature

- Agerstrand, M., M. Breitholtz and C. Ruden (2011). "Comparison of four different methods for reliability evaluation of ecotoxicity data: a case study of non-standard test data used in environmental risk assessments of pharmaceutical substances." *Environmental Sciences Europe* 23(1): 17.
- Aguirre-Martinez, G. V., S. Buratti, E. Fabbri, T. A. Del Valls and M. L. Martin-Diaz (2013). "Stability of lysosomal membrane in *Carcinus maenas* acts as a biomarker of exposure to pharmaceuticals." *Environmental Monitoring and Assessment* 185(5): 3783-3793.
- Aguirre-Martinez, G. V., T. A. Del Valls and M. Martin-Diaz (2013). "Identification of biomarkers responsive to chronic exposure to pharmaceuticals in target tissues of *Carcinus maenas*." *Marine Environmental Research* 87-88: 1-11.

- Aguirre-Martinez, G. V., T. A. DelValls and M. L. Martin-Diaz (2012). "Evaluation of adverse effects induced by carbamazepine and novobiocin drugs in *Carcinus maenas*." *Comparative Biochemistry and Physiology a-Molecular & Integrative Physiology* 163(1): \$10-\$11.
- Al Aukidy, M., P. Verlicchi, A. Jelic, M. Petrovic and D. Barcelo (2012). "Monitoring release of pharmaceutical compounds: Occurrence and environmental risk assessment of two WWTP effluents and their receiving bodies in the Po Valley, Italy." *Science of the Total Environment* 438: 15-25.
- Al-Ghanbousi, R., T. Ba-Omar and R. Victor (2012). "Effect of deltamethrin on the gills of *Aphanius dispar*: A microscopic study." *Tissue & Cell* 44(1): 7-14.
- Al-Odaini, N. A., M. P. Zakaria, M. I. Yaziz, S. Surif and M. Abdulghani (2013). "The occurrence of human pharmaceuticals in wastewater effluents and surface water of Langat River and its tributaries, Malaysia." *International Journal of Environmental Analytical Chemistry* 93(3): 245-264.
- Ambili, T. R., M. Saravanan, M. Ramesh, D. B. Abhijith and R. K. Poopal (2013). "Toxicological Effects of the Antibiotic Oxytetracycline to an Indian Major Carp *Labeo rohita*." *Archives of Environmental Contamination and Toxicology* 64(3): 494-503.
- Amin, K. A. and K. S. Hashem (2012). "Deltamethrin-induced oxidative stress and biochemical changes in tissues and blood of catfish (*Clarias gariepinus*): antioxidant defense and role of alpha-tocopherol." *Bmc Veterinary Research* 8.
- An, J. Y., H. N. Lee, K. I. Park, J. Y. Kim, J. Y. Lee and K. H. Park (2012). "Effects of Nacetylcysteine (NAC) on non-specific immune parameters, respiratory burst and lysozyme activities, in different fishes." *Journal of Fish Pathology* 25(1): 1-10.
- Ansari, R. A., S. Rahman, M. Kaur, S. Anjum and S. Raisuddin (2011). "*In vivo* cytogenetic and oxidative stress-inducing effects of cypermethrin in freshwater fish, *Channa punctata* Bloch." *Ecotoxicology and Environmental Safety* 74(1): 150-156.
- Anselmo, H. M. R., L. Koerting, S. Devito, J. H. J. van den Berg, M. Dubbeldam, C. Kwadijk and A. J. Murk (2011). "Early life developmental effects of marine persistent organic pollutants on the sea urchin *Psammechinus miliaris*." *Ecotoxicology and Environmental Safety* 74(8): 2182-2192.
- Antunes, S. C., R. Freitas, E. Figueira, F. Goncalves and B. Nunes (2013). "Biochemical effects of acetaminophen in aquatic species: edible clams *Venerupis decussata* and *Venerupis philippinarum*." *Environmental Science and Pollution Research* 20(9): 6658-6666.
- Arnold, K. E., A. B. A. Boxall, A. R. Brown, R. J. Cuthbert, S. Gaw, T. H. Hutchinson, S. Jobling, J. C. Madden, C. D. Metcalfe, V. Naidoo, R. F. Shore, J. E. Smits, M. A. Taggart and H. M. Thompson (2013). "Assessing the exposure risk and impacts of pharmaceuticals in the environment on individuals and ecosystems." *Biology letters* 9(4): 20130492-20130492.
- Aufartova, J., C. Mahugo-Santana, Z. Sosa-Ferrera, J. J. Santana-Rodriguez, L. Novakova and P. Solich (2011). "Determination of steroid hormones in biological and environmental samples using green microextraction techniques: An overview." *Analytica Chimica Acta* 704(1-2): 33-46.
- Aydin, E. and I. Talinli (2013). "Analysis, occurrence and fate of commonly used pharmaceuticals and hormones in the Buyukcekmece Watershed, Turkey." *Chemosphere* 90(6): 2004-2012.

- Aydin-Sinan, H., A. Gungordu and M. Ozmen (2012). "Toxic effects of deltamethrin and lambdacyhalothrin on *Xenopus laevis* tadpoles." *Journal of Environmental Science and Health Part B-Pesticides Food Contaminants and Agricultural Wastes* 47(5): 397-402.
- Aygun, D., S. Kaplan, E. Odaci, M. E. Onger and M. E. Altunkaynak (2012). "Toxicity of nonsteroidal anti-inflammatory drugs: a review of melatonin and diclofenac sodium association." *Histology and Histopathology* 27(4): 417-436.
- Backhaus, T., T. Porsbring, A. Arrhenius, S. Brosche, P. Johansson and H. Blanck (2011). "Singlesubstance and mixture toxicity of five pharmaceuticals and personal care products to marine periphyton communities." *Environmental Toxicology and Chemistry* 30(9): 2030-2040.
- Bajet, C. M., A. Kumar, M. N. Calingacion and T. C. Narvacan (2012). "Toxicological assessment of pesticides used in the Pagsanjan-Lumban catchment to selected non-target aquatic organisms in Laguna Lake, Philippines." *Agricultural Water Management* 106: 42-49.
- Bakry, F. A., W. S. Hasheesh and S. A. H. Hamdi (2011). "Biological, biochemical, and molecular parameters of *Helisoma duryi* snails exposed to the pesticides Malathion and Deltamethrin." *Pesticide Biochemistry and Physiology* 101(2): 86-92.
- Balcioglu, I. C., U. Z. Ok, Y. Ozbel, N. Girginkardesler and A. Ozbilgin (2012). "The in vitro Effects of Azithromycin and Clarithromycin on Promastigotes and Amastigotes of *Leishmania tropica*." *Kafkas Universitesi Veteriner Fakultesi Dergisi* 18: A115-A120.
- Baran, W., E. Adamek, J. Ziemianska and A. Sobczak (2011). "Effects of the presence of sulfonamides in the environment and their influence on human health." *Journal of Hazardous Materials* 196: 1-15.
- Barber, L. B., G. K. Brown, T. G. Nettesheim, E. W. Murphy, S. E. Bartell and H. L. Schoenfuss (2011). "Effects of biologically-active chemical mixtures on fish in a wastewater-impacted urban stream." *Science of the Total Environment* 409(22): 4720-4728.
- Barber, L. B., S. H. Keefe, G. K. Brown, E. T. Furlong, J. L. Gray, D. W. Kolpin, M. T. Meyer, M. W. Sandstrom and S. D. Zaugg (2013). "Persistence and potential effects of complex organic contaminant mixtures in wastewater-impacted streams." *Environmental Science & Technology* 47(5): 2177-2188.
- Bartram, A. E., M. J. Winter, D. B. Huggett, P. McCormack, L. A. Constantine, M. J. Hetheridge, T. H. Hutchinson, L. B. Kinter, J. F. Ericson, J. P. Sumpter and S. F. Owen (2012). "*In vivo* and *in vitro* liver and gill EROD activity in rainbow trout (*Oncorhynchus mykiss*) exposed to the beta-blocker propranolol." *Environmental Toxicology* 27(10): 573-582.
- Baudiffier, D., N. Hinfray, C. Ravaud, N. Creusot, E. Chadili, J. M. Porcher, R. W. Schulz and F.
 Brion (2013). "Effect of *in vivo* chronic exposure to clotrimazole on zebrafish testis function." *Environmental Science and Pollution Research* 20(5): 2747-2760.
- Baudiffier, D., N. Hinfray, M. Vosges, N. Creusot, E. Chadili, J. M. Porcher, R. W. Schulz and F. Brion (2012). "A critical role of follicle-stimulating hormone (Fsh) in mediating the effect of clotrimazole on testicular steroidogenesis in adult zebrafish." *Toxicology* 298(1-3): 30-39.
- Bayen, S. (2012). "Occurrence, bioavailability and toxic effects of trace metals and organic contaminants in mangrove ecosystems: A review." *Environment International* 48: 84-101.
- Bechi, N., G. Sorda, A. Spagnoletti, J. Bhattacharjee, E. A. V. Ferro, B. D. Barbosa, M. Frosini, M. Valoti, G. Sgaragli, L. Paulesu and F. Ietta (2013). "Toxicity assessment on trophoblast cells

for some environment polluting chemicals and 17 beta-estradiol." *Toxicology in Vitro* 27(3): 995-1000.

- Bedoux, G., B. Roig, O. Thomas, V. Dupont and B. Le Bot (2012). "Occurrence and toxicity of antimicrobial triclosan and by-products in the environment." *Environmental Science and Pollution Research* 19(4): 1044-1065.
- Benstead, R. S., A. Baynes, D. Casey, E. J. Routledge and S. Jobling (2011). "17 beta-Oestradiol may prolong reproduction in seasonally breeding freshwater gastropod molluscs." *Aquatic Toxicology* 101(2): 326-334.
- Bereswill, R., M. Streloke and R. Schulz (2013). "Current-use pesticides in stream water and suspended particles following runoff: Exposure, effects, and mitigation requirements." *Environmental Toxicology and Chemistry* 32(6): 1254-1263.
- Bergheim, M., R. Giere and K. Kummerer (2012). "Biodegradability and ecotoxicity of tramadol, ranitidine, and their photoderivatives in the aquatic environment." *Environmental Science and Pollution Research* 19(1): 72-85.
- Bergstrom, R. (2011). "The role of the pharmaceutical industry in meeting the public health threat of antibacterial resistance." *Drug Resistance Updates* 14(2): 77-78.
- Bhandari, K. and B. Venables (2011). "Ibuprofen bioconcentration and prostaglandin E2 levels in the bluntnose minnow *Pimephales notatus*." *Comparative Biochemistry and Physiology C-Toxicology & Pharmacology* 153(2): 251-257.
- Bilandzic, N., S. Tankovic, I. Varenina, B. S. Kolanovic and M. Smajlovic (2012).
 "Chloramphenicol residues in muscle of rainbow trout following two different dose treatments." *Bulletin of Environmental Contamination and Toxicology* 89(3): 461-466.
- Blanckenhorn, W. U., N. Puniamoorthy, A. Scheffczyk and J. Roembke (2013). "Evaluation of eco-toxicological effects of the parasiticide moxidectin in comparison to ivermectin in 11 species of dung flies." *Ecotoxicology and Environmental Safety* 89: 15-20.
- Boelsterli, U. A., M. R. Redinbo and K. S. Saitta (2013). "Multiple NSAID-induced hits injure the small intestine: underlying mechanisms and novel strategies." *Toxicological Sciences* 131(2): 654-667.
- Boltes, K., R. Rosal and E. Garcia-Calvo (2012). "Toxicity of mixtures of perfluorooctane sulphonic acid with chlorinated chemicals and lipid regulators." *Chemosphere* 86(1): 24-29.
- Boonstra, H., E. P. Reichman and P. J. van den Brink (2011). "Effects of the veterinary pharmaceutical ivermectin in indoor aquatic microcosms." *Archives of Environmental Contamination and Toxicology* 60(1): 77-89.
- Boxall, A. B. A., M. A. Rudd, B. W. Brooks, D. J. Caldwell, K. Choi, S. Hickmann, E. Innes, K. Ostapyk, J. P. Staveley, T. Verslycke, G. T. Ankley, K. F. Beazley, S. E. Belanger, J. P. Berninger, P. Carriquiriborde, A. Coors, P. C. DeLeo, S. D. Dyer, J. F. Ericson, F. Gagne, J. P. Giesy, T. Gouin, L. Hallstrom, M. V. Karlsson, D. G. J. Larsson, J. M. Lazorchak, F. Mastrocco, A. McLaughlin, M. E. McMaster, R. D. Meyerhoff, R. Moore, J. L. Parrott, J. R. Snape, R. Murray-Smith, M. R. Servos, P. K. Sibley, J. O. Straub, N. D. Szabo, E. Topp, G. R. Tetreault, V. L. Trudeau and G. Van Der Kraak (2012). "Pharmaceuticals and personal care products in the environment: What are the big questions?" *Environmental Health Perspectives* 120(9): 1221-1229.
- Brausch, J. M., K. A. Connors, B. W. Brooks and G. M. Rand (2012). Human pharmaceuticals in the aquatic environment: A review of recent toxicological studies and considerations for

toxicity testing. *Reviews of Environmental Contamination and Toxicology, Vol 218.* D. M. Whitacre. New York, Springer. 218: 1-99.

- Brausch, J. M. and G. M. Rand (2011). "A review of personal care products in the aquatic environment: Environmental concentrations and toxicity." *Chemosphere* 82(11): 1518-1532.
- Brezovsek, P., T. Elersek and M. Filipic (2013). "The effects of selected cytostatics and their mixtures on aquatic primary producers." *Toxicology Letters* 221: S96-S96.
- Brooks, B. W., J. P. Berninger, L. A. Kristofco, A. J. Ramirez, J. K. Stanley and T. W. Valenti (2012). Pharmaceuticals in the environment: Lessons learned for reducing uncertainties in environmental risk assessment. *Toxicology and Human Environments*. E. Hodgson. San Diego, Elsevier Academic Press Inc. 112: 231-258.
- Brown, A. R., L. K. Bickley, G. Le Page, D. J. Hosken, G. C. Paull, P. B. Hamilton, S. F. Owen, J. Robinson, A. D. Sharpe and C. R. Tyler (2011). "Are toxicological responses in laboratory (inbred) zebrafish representative of those in outbred (wild) populations? A case study with an endocrine disrupting chemical." *Environmental Science & Technology* 45(9): 4166-4172.
- Brown, J., M. J. Bernot and R. J. Bernot (2012). "The influence of TCS on the growth and behavior of the freshwater snail, *Physa acuta*." *Journal of Environmental Science and Health Part a-Toxic/Hazardous Substances & Environmental Engineering* 47(11): 1626-1630.
- Brozinski, J. M., M. Lahti, A. Meierjohann, A. Oikari and L. Kronberg (2013). "The antiinflammatory drugs diclofenac, naproxen and ibuprofen are found in the bile of wild fish caught downstream of a wastewater treatment plant." *Environmental Science & Technology* 47(1): 342-348.
- Bruchmann, J., S. Kirchen and T. Schwartz (2013). "Sub-inhibitory concentrations of antibiotics and wastewater influencing biofilm formation and gene expression of multi-resistant *Pseudomonas aeruginosa* wastewater isolates." *Environmental Science and Pollution Research* 20(6): 3539-3549.
- Buchinger, S., P. Heininger, M. Schlusener, G. Reifferscheid and E. Claus (2013). "Estrogenic effects along the river saale." *Environmental Toxicology and Chemistry* 32(3): 526-534.
- Bueno, M. J. M., M. J. Gomez, S. Herrera, M. D. Hernando, A. Agueera and A. R. Fernandez-Alba (2012). "Occurrence and persistence of organic emerging contaminants and priority pollutants in five sewage treatment plants of Spain: Two years pilot survey monitoring." *Environmental Pollution* 164: 267-273.
- Bunch, A. R. and M. J. Bernot (2011). "Distribution of nonprescription pharmaceuticals in central Indiana streams and effects on sediment microbial activity." *Ecotoxicology* 20(1): 97-109.
- Buono, S., S. Manzo, G. Maria and G. Sansone (2012). "Toxic effects of pentachlorophenol, azinphos-methyl and chlorpyrifos on the development of *Paracentrotus lividus* embryos." *Ecotoxicology* 21(3): 688-697.
- Buratti, S., S. Franzellitti, P. Valbonesi and E. Fabbri (2012). "Biomarkers and antioxidant responses in mussels exposed to fluoxetine, propranolol and their mixture." *Comparative Biochemistry and Physiology a-Molecular & Integrative Physiology* 163(1): S11-S11.
- Calderon-Preciado, D., V. Matamoros and J. M. Bayona (2011). "Occurrence and potential crop uptake of emerging contaminants and related compounds in an agricultural irrigation network." *Science of the Total Environment* 412: 14-19.

- Caldwell, D. J., F. Mastrocco, P. D. Anderson, R. Lange and J. P. Sumpter (2012). "Predicted-noeffect concentrations for the steroid estrogens estrone, 17 beta-estradiol, estriol, and 17 alpha-ethinylestradiol." *Environmental Toxicology and Chemistry* 31(6): 1396-1406.
- Calza, P., C. Medana, E. Padovano, V. Giancotti and C. Minero (2013). "Fate of selected pharmaceuticals in river waters." *Environmental Science and Pollution Research* 20(4): 2262-2270.
- Carballeira, C., M. R. De Orte, I. G. Viana, T. A. DelValls and A. Carballeira (2012). "Assessing the toxicity of chemical compounds associated with land-based marine fish farms: The sea urchin embryo bioassay with *Paracentrotus lividus* and *Arbacia lixula*." *Archives of Environmental Contamination and Toxicology* 63(2): 249-261.
- Carbonell-Martin, G., J. Pro-Gonzalez, P. Aragonese-Grunert, M. M. Babin-Vich, C. Fernandez-Torija and J. V. Tarazona-Lafarga (2011). "Targeting the environmental assessment of veterinary drugs with the multi-species-soil system (MS.3) agricultural soil microcosms: the ivermectin case study." *Spanish Journal of Agricultural Research* 9(2): 433-443.
- Carlos, L., D. O. Martire, M. C. Gonzalez, J. Gomis, A. Bernabeu, A. M. Amat and A. Arques (2012). "Photochemical fate of a mixture of emerging pollutants in the presence of humic substances." *Water Research* 46(15): 4732-4740.
- Carlsson, G., J. Patring, J. Kreuger, L. Norrgren and A. Oskarsson (2013). "Toxicity of 15 veterinary pharmaceuticals in zebrafish (*Danio rerio*) embryos." *Aquatic Toxicology* 126: 30-41.
- Casanova-Nakayama, A., M. Wenger, R. Burki, E. Eppler, A. Krasnov and H. Segner (2011). "Endocrine disrupting compounds: Can they target the immune system of fish?" *Marine pollution bulletin* 63(5-12): 412-416.
- Celebi, H. and D. T. Sponza (2012). "Comparison of the sensitivities of fish, Microtox and *Daphnia magna* bioassays to amoxycillin in anaerobic/aerobic sequential reactor systems." *Water Science and Technology* 66(5): 1117-1131.
- Cengiz, E. I., Y. Kan, V. Kizmaz, M. Bashan and M. Yanar (2012). "The protective role of vitamin E on the fatty acid composition of phospholipid structure in gill and liver tissues of *Oreochromis niloticus* exposed to deltamethrin." *Ecotoxicology and Environmental Safety* 80: 381-385.
- Ceyhun, S. B., E. Aksakal, B. Kirim, K. Atabeyoglu and O. Erdogan (2012). "Chronic toxicity of pesticides to the mRNA expression levels of metallothioneins and cytochrome P450 1A genes in rainbow trout." *Toxicology and Industrial Health* 28(2): 162-169.
- Chandra, K., T. Bosker, N. Hogan, A. Lister, D. MacLatchy and S. Currie (2012). "Sustained high temperature increases the vitellogenin response to 17 alpha-ethynylestradiol in mummichog (*Fundulus heteroclitus*)." *Aquatic Toxicology* 118: 130-140.
- Chen, J. L., S. Ravindran, S. Swift, L. J. Wright and N. Singhal (2012). "Catalytic oxidative degradation of 17 alpha-ethinylestradiol by Fe-III-TAML/H2O2: Estrogenicities of the products of partial, and extensive oxidation." *Water Research* 46(19): 6309-6318.
- Chen, X., Y. Zhou, D. Yang, H. Zhao, L. Wang and X. Yuan (2012). "CYP4 mRNA expression in marine polychaete *Perinereis aibuhitensis* in response to petroleum hydrocarbon and deltamethrin." *Marine Pollution Bulletin* 64(9): 1782-1788.

- Chen, Y., H. Li, Z. P. Wang, T. Tao, D. B. Wei and C. Hu (2012). "Photolysis of chlortetracycline in aqueous solution: Kinetics, toxicity and products." *Journal of Environmental Sciences-China* 24(2): 254-260.
- Claessens, M., L. Vanhaecke, K. Wille and C. R. Janssen (2013). "Emerging contaminants in Belgian marine waters: Single toxicant and mixture risks of pharmaceuticals." *Marine pollution bulletin* 71(1-2): 41-50.
- Clarke, B. O. and S. R. Smith (2011). "Review of 'emerging' organic contaminants in biosolids and assessment of international research priorities for the agricultural use of biosolids." *Environment International* 37(1): 226-247.
- Claudio Cacciatore, L., N. Verrengia Guerrero and A. Cristina Cochon (2013). "Cholinesterase and carboxylesterase inhibition in *Planorbarius corneus* exposed to binary mixtures of azinphos-methyl and chlorpyrifos." *Aquatic Toxicology* 128: 124-134.
- Conkle, J. L. and J. R. White (2012). "An initial screening of antibiotic effects on microbial respiration in wetland soils." *Journal of Environmental Science and Health Part a-Toxic/Hazardous Substances & Environmental Engineering* 47(10): 1381-1390.
- Contardo-Jara, V., C. Lorenz, S. Pflugmacher, G. Nuetzmann, W. Kloas and C. Wiegand (2011). "Exposure to human pharmaceuticals carbamazepine, ibuprofen and bezafibrate causes molecular effects in *Dreissena polymorpha*." *Aquatic Toxicology* 105(3-4): 428-437.
- Cortes, J. M., E. Larsson and J. A. Jonsson (2013). "Study of the uptake of non-steroid antiinflammatory drugs in wheat and soybean after application of sewage sludge as a fertilizer." *Science of the Total Environment* 449: 385-389.
- Cortez, F. S., C. D. Seabra Pereira, A. R. Santos, A. Cesar, R. B. Choueri, G. d. A. Martini and M. B. Bohrer-Morel (2012). "Biological effects of environmentally relevant concentrations of the pharmaceutical Triclosan in the marine mussel *Perna perna* (Linnaeus, 1758)." *Environmental Pollution* 168: 145-150.
- Coutu, S., L. Rossi, D. A. Barry, S. Rudaz and N. Vernaz (2013). "Temporal variability of antibiotics fluxes in wastewater and contribution from hospitals." *PloS one* 8(1): e53592.
- Cowden, J., B. Padnos, D. Hunter, R. MacPhail, K. Jensen and S. Padilla (2012). "Developmental exposure to valproate and ethanol alters locomotor activity and retino-tectal projection area in zebrafish embryos." *Reproductive Toxicology* 33(2): 165-173.
- Cuklev, F., J. Fick, M. Cvijovic, E. Kristiansson, L. Forlin and D. G. J. Larsson (2012). "Does ketoprofen or diclofenac pose the lowest risk to fish?" *Journal of Hazardous Materials* 229: 100-106.
- Cuklev, F., E. Kristiansson, J. Fick, N. Asker, L. Forlin and D. G. J. Larsson (2011). "Diclofenac in fish: Blood plasma levels similar to human therapeutical levels affect global hepatic gene expression." *Environmental Toxicology and Chemistry* 30(9): 2126-2134.
- Dahms, H.-U., A. Hagiwara and J.-S. Lee (2011). "Ecotoxicology, ecophysiology, and mechanistic studies with rotifers." *Aquatic Toxicology* 101(1): 1-12.
- Dalzell, A. and R. Glass (2013). "Extrapolation of toxicology endpoint data in developing integrated human and environmental risk assessments." *Toxicology Letters* 221: S221-S221.
- Dann, A. B. and A. Hontela (2011). "Triclosan: environmental exposure, toxicity and mechanisms of action." *Journal of Applied Toxicology* 31(4): 285-311.

- Daughton, C. G. (2011). Illicit Drugs: Contaminants in the Environment and Utility in Forensic Epidemiology. *Reviews of Environmental Contamination and Toxicology, Vol 210.* D. M. Whitacre. 210: 59-110.
- Dave, G. and G. Herger (2012). "Determination of detoxification to *Daphnia magna* of four pharmaceuticals and seven surfactants by activated sludge." *Chemosphere* 88(4): 459-466.
- David, A., H. Fenet, A. Escande, D. Munaron, D. Rosain, E. Maillot-Marechal, S. Ait-Aissa, C. Casellas and E. Gomez (2012). "*In vitro* biomonitoring of contamination by estrogenic compounds in coastal environments: Comments on the use of *M. galloprovincialis*." *Environmental Toxicology* 27(2): 74-82.
- D'Costa, P. M. and A. C. Anil (2012). "The effect of antibiotics on diatom communities." *Current Science* 102(11): 1552-1559.
- De Felice, B., L. Copia and M. Guida (2012). "Gene expression profiling in zebrafish embryos exposed to diclofenac, an environmental toxicant." *Molecular Biology Reports* 39(3): 2119-2128.
- de Garcia, S. O., G. P. Pinto, P. G. Encina and R. I. Mata (2013). "Consumption and occurrence of pharmaceutical and personal care products in the aquatic environment in Spain." *Science of the Total Environment* 444: 451-465.
- de Jong, E., A. Doedee, M. A. Reis-Fernandes, H. Nau and A. H. Piersma (2011). "Potency ranking of valproic acid analogues as to inhibition of cardiac differentiation of embryonic stem cells in comparison to their *in vivo* embryotoxicity." *Reproductive Toxicology* 31(4): 375-382.
- de Mendoza, J. H., L. Maggi, L. Bonetto, B. R. Carmena, A. Lezana, F. A. Mocholi and M. Carmona (2012). "Validation of antibiotics in catfish by on-line solid phase extraction coupled to liquid chromatography tandem mass spectrometry." *Food Chemistry* 134(2): 1149-1155.
- Deblonde, T., C. Cossu-Leguille and P. Hartemann (2011). "Emerging pollutants in wastewater: A review of the literature." *International Journal of Hygiene and Environmental Health* 214(6): 442-448.
- Deblonde, T. and P. Hartemann (2013). "Environmental impact of medical prescriptions: assessing the risks and hazards of persistence, bioaccumulation and toxicity of pharmaceuticals." *Public Health* 127(4): 312-317.
- Dirany, A., S. E. Aaron, N. Oturan, I. Sires, M. A. Oturan and J. J. Aaron (2011). "Study of the toxicity of sulfamethoxazole and its degradation products in water by a bioluminescence method during application of the electro-Fenton treatment." *Analytical and Bioanalytical Chemistry* 400(2): 353-360.
- Donner, E., T. Kosjek, S. Qualmann, K. O. Kusk, E. Heath, D. M. Revitt, A. Ledin and H. R. Andersen (2013). "Ecotoxicity of carbamazepine and its UV photolysis transformation products." *Science of the Total Environment* 443: 870-876.
- Doyle, M. A., T. Bosker, C. J. Martyniuk, D. L. Maclatchy and K. R. Munkittrick (2013). "The effects of 17-alpha-ethinylestradiol (EE2) on molecular signaling cascades in mumnichog (*Fundulus heteroclitus*)." *Aquatic toxicology (Amsterdam, Netherlands)* 134-135: 34-46.
- Du, B., P. Perez-Hurtado, B. W. Brooks and C. K. Chambliss (2012). "Evaluation of an isotope dilution liquid chromatography tandem mass spectrometry method for pharmaceuticals in fish." *Journal of Chromatography A* 1253: 177-183.

- Duan, Y. P., X. Z. Meng, Z. H. Wen and L. Chen (2013). "Acidic pharmaceuticals in domestic wastewater and receiving water from hyper-urbanization city of China (Shanghai): environmental release and ecological risk." *Environmental Science and Pollution Research* 20(1): 108-116.
- Duan, Y. P., X. Z. Meng, Z. H. Wen, R. H. Ke and L. Chen (2013). "Multi-phase partitioning, ecological risk and fate of acidic pharmaceuticals in a wastewater receiving river: The role of colloids." *Science of the Total Environment* 447: 267-273.
- Ebert, I., J. Bachmann, U. Kuhnen, A. Kuster, C. Kussatz, D. Maletzki and C. Schluter (2011). "Toxicity of the fluoroquinolone antibiotics enrofloxacin and ciprofloxacin to photoautotrophic aquatic organisms." *Environmental Toxicology and Chemistry* 30(12): 2786-2792.
- Eggen, T., T. N. Asp, K. Grave and V. Hormazabal (2011). "Uptake and translocation of metformin, ciprofloxacin and narasin in forage- and crop plants." *Chemosphere* 85(1): 26-33.
- El-Bassat, R. A., H. E. Touliabah and G. I. Harisa (2012). "Toxicity of four pharmaceuticals from different classes to isolated plankton species." *African Journal of Aquatic Science* 37(1): 71-80.
- Ellis, L. D. and K. H. Soanes (2012). "A larval zebrafish model of bipolar disorder as a screening platform for neuro-therapeutics." *Behavioural Brain Research* 233(2): 450-457.
- Ensibi, C., M. Perez-Lopez, F. S. Rodriguez, M. P. Miguez-Santiyan, M. N. D. Yahya and D. Hernandez-Moreno (2013). "Effects of deltamethrin on biometric parameters and liver biomarkers in common carp (*Cyprinus carpio* L.)." *Environmental Toxicology and Pharmacology* 36(2): 384-391.
- Escher, B. I., R. Baumgartner, M. Koller, K. Treyer, J. Lienert and C. S. McArdell (2011). "Environmental toxicology and risk assessment of pharmaceuticals from hospital wastewater." *Water Research* 45(1): 75-92.
- Escher, B. I. and K. Fenner (2011). "Recent advances in environmental risk assessment of transformation products." *Environmental Science & Technology* 45(9): 3835-3847.
- Esteban, S., J. Fernandez Rodriguez, G. Diaz Lopez, M. Nunez, Y. Valcarcel and M. Catala (2013). "New microbioassays based on biomarkers are more sensitive to fluvial water micropollution than standard testing methods." *Ecotoxicology and Environmental Safety* 93: 52-59.
- Fairchild, W., K. Doe, P. Jackman, F. Akaishi, J. Arsenault, J. Aube, M. Losier and A. Cook (2011). "Acute and chronic toxicity of two formulations of the pyrethroid pesticide deltamethrin to amphipods, sand shrimp and lobster larvae (PO)." *Canadian Technical Report of Fisheries* and Aquatic Sciences 2949: 64-65.
- Fang, Y. X., G. G. Ying, J. L. Zhao, F. Chen, S. Liu, L. J. Zhang and B. Yang (2012). "Assessment of hormonal activities and genotoxicity of industrial effluents using *in vitro* bioassays combined with chemical analysis." *Environmental Toxicology and Chemistry* 31(6): 1273-1282.
- Farghaly, M. F. M., S. Zayed and S. M. Soliman (2013). "Deltamethrin degradation and effects on soil microbial activity." *Journal of Environmental Science and Health Part B-Pesticides Food Contaminants and Agricultural Wastes* 48(7): 575-581.

- Fatta-Kassinos, D., M. I. Vasquez and K. Kummerer (2011). "Transformation products of pharmaceuticals in surface waters and wastewater formed during photolysis and advanced oxidation processes Degradation, elucidation of byproducts and assessment of their biological potency." *Chemosphere* 85(5): 693-709.
- Feito, R., Y. Valcarcel and M. Catala (2012). "Biomarker assessment of toxicity with miniaturised bioassays: diclofenac as a case study." *Ecotoxicology* 21(1): 289-296.
- Feito, R., Y. Valcarcel and M. Catala (2013). "Preliminary data suggest that venlafaxine environmental concentrations could be toxic to plants." *Chemosphere* 90(7): 2065-2069.
- Ferguson, P. J., M. J. Bernot, J. C. Doll and T. E. Lauer (2013). "Detection of pharmaceuticals and personal care products (PPCPs) in near-shore habitats of southern Lake Michigan." *The Science of the total environment* 458-460: 187-196.
- Fernandes, D., S. Schnell and C. Porte (2011). "Can pharmaceuticals interfere with the synthesis of active androgens in male fish? An *in vitro* study." *Marine pollution bulletin* 62(10): 2250-2253.
- Ferrando-Climent, L., N. Collado, G. Buttiglieri, M. Gros, I. Rodriguez-Roda, S. Rodriguez-Mozaz and D. Barcelo (2012). "Comprehensive study of ibuprofen and its metabolites in activated sludge batch experiments and aquatic environment." *Science of the Total Environment* 438: 404-413.
- Ferrari, F., A. Gallipoli, M. Balderacchi, M. M. Ulaszewska, E. Capri and M. Trevisan (2011).
 "Exposure of the main Italian river basin to pharmaceuticals." *Journal of toxicology* 2011: 989270.
- Finn, J., M. Hui, V. Li, V. Lorenzi, N. de la Paz, S. H. Cheng, L. Lai-Chan and D. Schlenk (2012). "Effects of propranolol on heart rate and development in Japanese medaka (*Oryzias latipes*) and zebrafish (*Danio rerio*)." *Aquatic Toxicology* 122: 214-221.
- Fong, P. P. and C. M. Hoy (2012). "Antidepressants (venlafaxine and citalopram) cause foot detachment from the substrate in freshwater snails at environmentally relevant concentrations." *Marine and Freshwater Behaviour and Physiology* 45(2): 145-153.
- Franzellitti, S., S. Buratti, P. Valbonesi, A. Capuzzo and E. Fabbri (2011). "The beta-blocker propranolol affects cAMP-dependent signaling and induces the stress response in Mediterranean mussels, *Mytilus galloprovincialis*." *Aquatic Toxicology* 101(2): 299-308.
- Franzellitti, S., S. Buratti, P. Valbonesi and E. Fabbri (2013). "The mode of action (MOA) approach reveals interactive effects of environmental pharmaceuticals on *Mytilus galloprovincialis*." *Aquatic Toxicology* 140: 249-256.
- Franzellitti, S. and E. Fabbri (2013). "Cyclic-AMP Mediated Regulation of ABCB mRNA Expression in Mussel Haemocytes." *Plos One* 8(4).
- Fuzier, R., I. Serres, E. Guitton, M. Lapeyre-Mestre, J. L. Montastruc and P. French Network (2013). "Adverse drug reactions to gabapentin and pregabalin - A review of the French pharmacovigilance database." *Drug Safety* 36(1): 55-62.
- Gagne, F., C. Andre, M. Fortier and M. Fournier (2012). "Immunotoxic potential of aeration lagoon effluents for the treatment of domestic and hospital wastewaters in the freshwater mussel *Elliptio complanata*." *Journal of Environmental Sciences-China* 24(5): 781-789.
- Galus, M., J. Jeyaranjaan, E. Smith, H. Li, C. Metcalfe and J. Y. Wilson (2013). "Chronic effects of exposure to a pharmaceutical mixture and municipal wastewater in zebrafish." *Aquatic toxicology (Amsterdam, Netherlands)* 132-133: 212-222.

- Galus, M., N. Kirischian, S. Higgins, J. Purdy, J. Chow, S. Rangaranjan, H. X. Li, C. Metcalfe and J. Y. Wilson (2013). "Chronic, low concentration exposure to pharmaceuticals impacts multiple organ systems in zebrafish." *Aquatic Toxicology* 132: 200-211.
- Gao, L. H., Y. L. Shi, W. H. Li, J. M. Liu and Y. Q. Cai (2012). "Occurrence, distribution and bioaccumulation of antibiotics in the Haihe River in China." *Journal of Environmental Monitoring* 14(4): 1248-1255.
- Garcia, S. N., M. Foster, L. A. Constantine and D. B. Huggett (2012). "Field and laboratory fish tissue accumulation of the anti-convulsant drug carbamazepine." *Ecotoxicology and Environmental Safety* 84: 207-211.
- Garcia-Galan, M. J., M. S. Diaz-Cruz and D. Barcelo (2011). "Occurrence of sulfonamide residues along the Ebro river basin Removal in wastewater treatment plants and environmental impact assessment." *Environment International* 37(2): 462-473.
- Garcia-Kaufer, M., T. Haddad, M. Bergheim, R. Gminski, P. Gupta, N. Mathur, K. Kummerer and V. Mersch-Sundermann (2012). "Genotoxic effect of ciprofloxacin during photolytic decomposition monitored by the *in vitro* micronucleus test (MNvit) in HepG2 cells." *Environmental Science and Pollution Research* 19(5): 1719-1727.
- Gaume, B., N. Bourgougnon, S. Auzoux-Bordenave, B. Roig, B. Le Bot and G. Bedoux (2012). "*In vitro* effects of triclosan and methyl-triclosan on the marine gastropod *Haliotis tuberculata*." *Comparative Biochemistry and Physiology C-Toxicology & Pharmacology* 156(2): 87-94.
- Gelsleichter, J. and N. J. Szabo (2013). "Uptake of human pharmaceuticals in bull sharks (*Carcharhinus leucas*) inhabiting a wastewater-impacted river." *The Science of the total environment* 456-457: 196-201.
- Giandomenico, S., N. Cardellicchio, C. Annicchiarico, L. Lopez, M. Maggi, L. Spada and A. Di Leo (2011). "Simultaneous determination of pharmaceutical compounds in environmental samples by solid-phase extraction and gas chromatography-mass spectrometry." *Chemistry and Ecology* 27: 127-136.
- Gielen, G., P. W. Clinton, M. R. Van den Heuvel, M. O. Kimberley and L. G. Greenfield (2011)."Influence of sewage and pharmaceuticals on soil microbial function." *Environmental Toxicology and Chemistry* 30(5): 1086-1095.
- Gilroy, E. A. M., V. K. Balakrishnan, K. R. Solomon, E. Sverko and P. K. Sibley (2012). "Behaviour of pharmaceuticals in spiked lake sediments Effects and interactions with benthic invertebrates." *Chemosphere* 86(6): 578-584.
- Girardi, C., K. M. Nowak, O. Carranza-Diaz, B. Lewkow, A. Miltner, M. Gehre, A. Schaffer and M. Kastner (2013). "Microbial degradation of the pharmaceutical ibuprofen and the herbicide 2,4-D in water and soil Use and limits of data obtained from aqueous systems for predicting their fate in soil." *Science of the Total Environment* 444: 32-42.
- Gokce, B., N. Gencer, O. Arslan, S. A. Turkoglu, M. Alper and F. Kockar (2012). "Evaluation of *in vitro* effects of some analgesic drugs on erythrocyte and recombinant carbonic anhydrase I and II." *Journal of Enzyme Inhibition and Medicinal Chemistry* 27(1): 37-42.
- Gomez, E., M. Bachelot, C. Boillot, D. Munaron, S. Chiron, C. Casellas and H. Fenet (2012).
 "Bioconcentration of two pharmaceuticals (benzodiazepines) and two personal care products (UV filters) in marine mussels (*Mytilus galloprovincialis*) under controlled laboratory conditions." *Environmental Science and Pollution Research* 19(7): 2561-2569.

- Gomiero, A., A. Dagnino, S. Sforzini, A. Negri and A. Viarengo (2012). "Heat stress effects on toxicity of copper and oxytetracycline on the marine protozoa *Euplotes crassus* in a climate change perspective." *Comparative Biochemistry and Physiology a-Molecular & Integrative Physiology* 163(1): S4-S4.
- Gonzalez, S., R. Lopez-Roldan and J. L. Cortina (2012). "Presence and biological effects of emerging contaminants in Llobregat River basin: A review." *Environmental Pollution* 161: 83-92.
- Gonzalez-Pleiter, M., S. Gonzalo, I. Rodea-Palomares, F. Leganes, R. Rosal, K. Boltes, E. Marco and F. Fernandez-Pinas (2013). "Toxicity of five antibiotics and their mixtures towards photosynthetic aquatic organisms: Implications for environmental risk assessment." *Water Research* 47(6): 2050-2064.
- Gonzalez-Rey, M. and M. J. Bebianno (2011). "Non-steroidal anti-inflammatory drug (NSAID) ibuprofen distresses antioxidant defense system in mussel *Mytilus galloprovincialis* gills." *Aquatic Toxicology* 105(3-4): 264-269.
- Gonzalez-Rey, M. and M. J. Bebianno (2012). "Does non-steroidal anti-inflammatory (NSAID) ibuprofen induce antioxidant stress and endocrine disruption in mussel *Mytilus galloprovincialis*?" *Environmental Toxicology and Pharmacology* 33(2): 361-371.
- Gonzalez-Rey, M. and M. J. Bebianno (2012). "Non-steroidal anti-inflammatory drug (NSAID) diclofenac exposure effects in mussel *Mytilus galloprovincialis*." *Comparative Biochemistry and Physiology a-Molecular & Integrative Physiology* 163(1): S34-S34.
- Goulding, A. T., L. K. Shelley, P. S. Ross and C. J. Kennedy (2013). "Reduction in swimming performance in juvenile rainbow trout (*Oncorhynchus mykiss*) following sublethal exposure to pyrethroid insecticides." *Comparative Biochemistry and Physiology C-Toxicology & Pharmacology* 157(3): 280-286.
- Gozlan, I., A. Rotstein and D. Avisar (2013). "Amoxicillin-degradation products formed under controlled environmental conditions: Identification and determination in the aquatic environment." *Chemosphere* 91(7): 985-992.
- Grassi, M., L. Rizzo and A. Farina (2013). "Endocrine disruptors compounds, pharmaceuticals and personal care products in urban wastewater: implications for agricultural reuse and their removal by adsorption process." *Environmental Science and Pollution Research* 20(6): 3616-3628.
- Grundmann, H., K. P. Klugman, T. Walsh, P. Ramon-Pardo, B. Sigauque, W. Khan, R. Laxminarayan, A. Heddini and J. Stelling (2011). "A framework for global surveillance of antibiotic resistance." *Drug Resistance Updates* 14(2): 79-87.
- Guiloski, I. C., S. C. Rossi, C. A. da Silva and H. C. S. De Assis (2013). "Insecticides biomarker responses on a freshwater fish *Corydoras paleatus* (Pisces: Callichthyidae)." *Journal of Environmental Science and Health Part B-Pesticides Food Contaminants and Agricultural Wastes* 48(4): 272-277.
- Gunes, E. and S. V. Yerli (2011). "Effects of deltamethrin on lipase activity in guppies (*Poecilia reticulata*)." *Turkish Journal of Fisheries and Aquatic Sciences* 11(3): 473-476.
- Guo, R. X. and J. Q. Chen (2012). "Phytoplankton toxicity of the antibiotic chlortetracycline and its UV light degradation products." *Chemosphere* 87(11): 1254-1259.
- Gust, M., M. Fortier, J. Garric, M. Fournier and F. Gagne (2013). "Effects of short-term exposure to environmentally relevant concentrations of different pharmaceutical mixtures on the

immune response of the pond snail *Lymnaea stagnalis*." *Science of the Total Environment* 445: 210-218.

- Gust, M., M. Gelinas, M. Fortier, M. Fournier and F. Gagne (2012). "In vitro immunotoxicity of environmentally representative antibiotics to the freshwater mussel *Elliptio complanata*." *Environmental Pollution* 169: 50-58.
- Haack, S. K., D. W. Metge, L. R. Fogarty, M. T. Meyer, L. B. Barber, R. W. Harvey, D. R. LeBlanc and D. W. Kolpin (2012). "Effects on groundwater microbial communities of an engineered 30-day *in situ* exposure to the antibiotic sulfamethoxazole." *Environmental Science & Technology* 46(14): 7478-7486.
- Hagenbuch, I. M. and J. L. Pinckney (2012). "Toxic effect of the combined antibiotics ciprofloxacin, lincomycin, and tylosin on two species of marine diatoms." *Water Research* 46(16): 5028-5036.
- Hallgren, P., Z. Sorita, O. Berglund and A. Persson (2012). "Effects of 17 alpha-ethinylestradiol on individual life-history parameters and estimated population growth rates of the freshwater gastropods *Radix balthica* and *Bithynia tentaculata*." *Ecotoxicology* 21(3): 803-810.
- Hallgren, S., K. Volkova, N. Reyhanian, K. H. Olsen and I. P. Hallstrom (2011). "Anxiogenic behaviour induced by 17 alpha-ethynylestradiol in male guppies (*Poecilia reticulata*)." *Fish Physiology and Biochemistry* 37(4): 911-918.
- Hampel, M., E. Alonso, I. Aparicio, J. E. Bron, J. L. Santos, J. B. Taggart and M. J. Leaver (2012). "Atlantic salmon liver transcriptome response to exposure to three human pharmaceuticals." *ArrayExpress Archive*.
- Hapeshi, E., A. Lambrianides, P. Koutsoftas, E. Kastanos, C. Michael and D. Fatta-Kassinos (2013).
 "Investigating the fate of iodinated X-ray contrast media iohexol and diatrizoate during microbial degradation in an MBBR system treating urban wastewater." *Environmental Science and Pollution Research* 20(6): 3592-3606.
- Harmon, S. M. and F. E. Wiley (2011). "Effects of pollution on freshwater organisms." *Water Environment Research* 83(10): 1733-1788.
- He, J. H., S. Y. Guo, F. Zhu, J. J. Zhu, Y. X. Chen, C. J. Huang, J. M. Gao, Q. X. Dong, Y. X. Xuan and C. Q. Li (2013). "A zebrafish phenotypic assay for assessing drug-induced hepatotoxicity." *Journal of Pharmacological and Toxicological Methods* 67(1): 25-32.
- He, X. T., X. P. Nie, Z. H. Wang, Z. Cheng, K. B. Li, G. Y. Li, M. H. Wong, X. M. Liang and M. T. K. Tsui (2011). "Assessment of typical pollutants in waterborne by combining active biomonitoring and integrated biomarkers response." *Chemosphere* 84(10): 1422-1431.
- He, X. T., X. P. Nie, Y. F. Yang, X. Y. Liu, D. B. Pan, Z. Cheng and X. M. Liang (2012). "Multibiomarker responses in fishes from two typical marine aquaculture regions of South China." *Marine pollution bulletin* 64(11): 2317-2324.
- Helbing, C. C., C. R. Propper and N. Veldhoen (2011). "Triclosan affects the thyroid axis of amphibians." *Toxicological Sciences* 123(2): 601-602.
- Hernandez-Moreno, D., L. Guilhermino, C. Gravato, F. Soler and M. Perez-Lopez (2011). "Acute toxicity of the pesticides carbofuran and deltamethrin on sea bass (*Dicentrarchus labrax*)." *Toxicology Letters* 205: S230-S230.

- Hillis, D. G., J. Fletcher, K. R. Solomon and P. K. Sibley (2011). "Effects of ten antibiotics on seed germination and root elongation in three plant species." *Archives of Environmental Contamination and Toxicology* 60(2): 220-232.
- Hinfray, N., D. Baudiffier, M. C. Leal, J. M. Porcher, S. Ait-Aissa, F. Le Gac, R. W. Schulz and F.
 Brion (2011). "Characterization of testicular expression of P450 17 alpha-hydroxylase, 17,20-lyase in zebrafish and its perturbation by the pharmaceutical fungicide clotrimazole." *General and Comparative Endocrinology* 174(3): 309-317.
- Hirakawa, I., S. Miyagawa, Y. Katsu, Y. Kagami, N. Tatarazako, T. Kobayashi, T. Kusano, T. Mizutani, Y. Ogino, T. Takeuchi, Y. Ohta and T. Iguchi (2012). "Gene expression profiles in the testis associated with testis-ova in adult Japanese medaka (*Oryzias latipes*) exposed to 17 alpha-ethinylestradiol." *Chemosphere* 87(7): 668-674.
- Ho, K. T., A. A. Chariton, L. M. Portis, D. Proestou, M. G. Cantwell, J. G. Baguley, R. M. Burgess, S. Simpson, M. C. Pelletier, M. M. Perron, C. K. Gunsch, H. M. Bik, D. Katz and A. Kamikawa (2013). "Use of a novel sediment exposure to determine the effects of triclosan on estuarine benthic communities." *Environmental Toxicology and Chemistry* 32(2): 384-392.
- Hoffmann, F. and W. Kloas (2012). "Estrogens can disrupt amphibian mating behavior." *PloS* one 7(2).
- Homem, V. and L. Santos (2011). "Degradation and removal methods of antibiotics from aqueous matrices A review." *Journal of Environmental Management* 92(10): 2304-2347.
- Huerta, B., A. Jakimska, M. Gros, S. Rodriguez-Mozaz and D. Barcelo (2013). "Analysis of multiclass pharmaceuticals in fish tissues by ultra-high-performance liquid chromatography tandem mass spectrometry." *Journal of Chromatography A* 1288: 63-72.
- Huerta, B., S. Rodriguez-Mozaz and D. Barcelo (2012). "Pharmaceuticals in biota in the aquatic environment: analytical methods and environmental implications." *Analytical and Bioanalytical Chemistry* 404(9): 2611-2624.
- Hughes, S. R., P. Kay and L. E. Brown (2013). "Global synthesis and critical evaluation of pharmaceutical data sets collected from river systems." *Environmental Science & Technology* 47(2): 661-677.
- Huynh Thi, T., F. Silvestre, B. De Meulder, J.-P. Thome, P. Nguyen Thanh and P. Kestemont (2012). "Combined effects of deltamethrin, temperature and salinity on oxidative stress biomarkers and acetylcholinesterase activity in the black tiger shrimp (*Penaeus monodon*)." *Chemosphere* 86(1): 83-91.
- Igbinosa, E. O., E. E. Odjadjare, V. N. Chigor, I. H. Igbinosa, A. O. Emoghene, F. O. Ekhaise, N. O. Igiehon and O. G. Idemudia (2013). "Toxicological profile of chlorophenols and their derivatives in the environment: The public health perspective." *Scientific World Journal*.
- Igbinosa, I. H. and A. I. Okoh (2012). "Antibiotic susceptibility profile of *Aeromonas* species isolated from wastewater treatment plant." *Scientific World Journal*.
- Ings, J. S., N. George, M. C. S. Peter, M. R. Servos and M. M. Vijayan (2012). "Venlafaxine and atenolol disrupt epinephrine-stimulated glucose production in rainbow trout hepatocytes." *Aquatic Toxicology* 106: 48-55.
- Islas-Flores, H., L. M. Gomez-Olivan, M. Galar-Martinez, A. Colin-Cruz, N. Neri-Cruz and S. Garcia-Medina (2013). "Diclofenac-induced oxidative stress in brain, liver, gill and blood of common carp (*Cyprinus carpio*)." *Ecotoxicology and Environmental Safety* 92: 32-38.

- Jawecki, G. N., L. Szczesny, D. Solecka and J. Sawicki (2011). "Short ingestion tests as alternative proposal for conventional range finding assays with *Thamnocephalus platyurus* and *Brachionus calyciflorus*." *International Journal of Environmental Science and Technology* 8(4): 687-694.
- Jelic, A., C. Cruz-Morato, E. Marco-Urrea, M. Sarra, S. Perez, T. Vicent, M. Petrovic and D. Barcelo (2012). "Degradation of carbamazepine by *Trametes versicolor* in an air pulsed fluidized bed bioreactor and identification of intermediates." *Water Research* 46(4): 955-964.
- Jensen, J. and J. J. Scott-Fordsmand (2012). "Ecotoxicity of the veterinary pharmaceutical ivermectin tested in a soil multi-species (SMS) system." *Environmental Pollution* 171: 133-139.
- Jeon, J., D. Kurth, R. Ashauer and J. Hollender (2013). "Comparative toxicokinetics of organic micropollutants in freshwater crustaceans." *Environmental Science & Technology* 47(15): 8809-8817.
- Jerbi, M. A., Z. Ouanes, R. Besbes, L. Achour and A. Kacem (2011). "Single and combined genotoxic and cytotoxic effects of two xenobiotics widely used in intensive aquaculture." *Mutation Research-Genetic Toxicology and Environmental Mutagenesis* 724(1-2): 22-27.
- Jerbi, M. A., Z. Ouanes, R. Besbes, Z. Haouas, L. Achour and A. Kacem (2011). "Single and combined effects associated with two xenobiotics widely used in intensive aquaculture on European sea bass (*Dicentrarchus labrax*)." *Toxicology Letters* 205: S119-S119.
- Jesus Garcia-Galan, M., S. Gonzalez Blanco, R. Lopez Roldan, S. Diaz-Cruz and D. Barcelo (2012). "Ecotoxicity evaluation and removal of sulfonamides and their acetylated metabolites during conventional wastewater treatment." *Science of the Total Environment* 437: 403-412.
- Ji, J.-Y., Y.-J. Xing, Z.-T. Ma, M. Zhang and P. Zheng (2013). "Acute toxicity of pharmaceutical wastewaters containing antibiotics to anaerobic digestion treatment." *Chemosphere* 91(8): 1094-1098.
- Ji, J.-y., Y.-j. Xing and P. Zheng (2012). "Acute toxicity of antibiotics and anaerobic digestion intermediates in pharmaceutical wastewaters." *Huanjing Kexue* 33(12): 4367-4372.
- Ji, K., S. Kim, S. Han, J. Seo, S. Lee, Y. Park, K. Choi, Y. L. Kho, P. G. Kim and J. Park (2012). "Risk assessment of chlortetracycline, oxytetracycline, sulfamethazine, sulfathiazole, and erythromycin in aquatic environment: are the current environmental concentrations safe?" *Ecotoxicology* 21(7): 2031-2050.
- Ji, Y., C. Zeng, C. Ferronato, J.-M. Chovelon and X. Yang (2012). "Nitrate-induced photodegradation of atenolol in aqueous solution: Kinetics, toxicity and degradation pathways." *Chemosphere* 88(5): 644-649.
- Jo, Y., J.-H. Oh, S. Yoon, H. Bae, M.-C. Hong, M.-K. Shin and Y. Kim (2012). "The comparative analysis of in vivo and in vitro transcriptome data based on systems biology." *Biochip Journal* 6(3): 280-292.
- Johns, S. M., N. D. Denslow, M. D. Kane, K. H. Watanabe, E. F. Orlando and M. S. Sepulveda (2011). "Effects of estrogens and antiestrogens on gene expression of fathead minnow (*Pimephales promelas*) early life stages." *Environmental Toxicology* 26(2): 195-206.
- Jones, B., D. A. Smith and E. F. Schmid (2012). "Are predictive biomarkers of toxicity worth having? An economic model." *Xenobiotica* 42(1): 4-10.
- Jones, H. S., H. T. Trollope, T. H. Hutchinson, G. H. Panter and J. K. Chipman (2012). "Metabolism of ibuprofen in zebrafish larvae." *Xenobiotica* 42(11): 1069-1075.

- Jubeaux, G., R. Simon, A. Salvador, H. Queau, A. Chaumot and O. Geffard (2012). "Vitellogeninlike proteins in the freshwater amphipod *Gammarus fossarum* (Koch, 1835): Functional characterization throughout reproductive process, potential for use as an indicator of oocyte quality and endocrine disruption biomarker in males." *Aquatic Toxicology* 112: 72-82.
- Kaptaner, B. and G. Unal (2011). "Effects of 17 alpha-ethynylestradiol and nonylphenol on liver and gonadal apoptosis and histopathology in *Chalcalburnus tarichi*." *Environmental Toxicology* 26(6): 610-622.
- Kar, S. and K. Roy (2012). "Risk assessment for ecotoxicity of pharmaceuticals an emerging issue." *Expert Opinion on Drug Safety* 11(2): 235-274.
- Karaaslan, M. A., H. Parlak, O. C. Arslan and M. Boyacioglu (2012). "The embryotoxic effects of beta blocker atenolol on sea urchin *Arbacia lixula* embryos." *Fresenius Environmental Bulletin* 21(11B): 3362-3364.
- Karimullina, E., Y. Li, G. K. Ginjupalli and W. S. Baldwin (2012). "*Daphnia* HR96 is a promiscuous xenobiotic and endobiotic nuclear receptor." *Aquatic Toxicology* 116: 69-78.
- Kaur, K., B. Singh and A. K. Malik (2011). "Chemiluminescence and spectrofluorimetric methods for determination of fluoroquinolones: a review." *Analytical Letters* 44(9): 1602-1639.
- Kavitha, P., R. Ramesh, G. Bupesh, A. Stalin and P. Subramanian (2011). "Hepatoprotective activity of *Tribulus terrestris* extract against acetaminophen-induced toxicity in a freshwater fish (*Oreochromis mossambicus*)." *In Vitro Cellular & Developmental Biology-Animal* 47(10): 698-706.
- Kawabata, K., K. Sugihara, S. Sanoh, S. Kitamura and S. Ohta (2012). "Ultraviolet-photoproduct of acetaminophen: Structure determination and evaluation of ecotoxicological effect." *Journal of Photochemistry and Photobiology a-Chemistry* 249: 29-35.
- Khadra, A., E. Pinelli, M. Z. Lacroix, A. Bousquet-Melou, H. Hamdi, G. Merlina, M. Guiresse and M. Hafidi (2012). "Assessment of the genotoxicity of quinolone and fluoroquinolones contaminated soil with the *Vicia faba* micronucleus test." *Ecotoxicology and Environmental Safety* 76: 187-192.
- Khataee, A. R., M. Fathinia and S. W. Joo (2013). "Simultaneous monitoring of photocatalysis of three pharmaceuticals by immobilized TiO2 nanoparticles: Chemometric assessment, intermediates identification and ecotoxicological evaluation." *Spectrochimica Acta Part a-Molecular and Biomolecular Spectroscopy* 112: 33-45.
- Kim, H., Y. Hong, J. E. Park, V. K. Sharma and S. I. Cho (2013). "Sulfonamides and tetracyclines in livestock wastewater." *Chemosphere* 91(7): 888-894.
- Kim, H. Y., M. J. Lee, S. H. Yu and S. D. Kim (2012). "The individual and population effects of tetracycline on *Daphnia magna* in multigenerational exposure." *Ecotoxicology* 21(4): 993-1002.
- Kim, P., Y. Park, K. Ji, J. Seo, S. Lee, K. Choi, Y. Kho and J. Park (2012). "Effect of chronic exposure to acetaminophen and lincomycin on Japanese medaka (*Oryzias latipes*) and freshwater cladocerans *Daphnia magna* and *Moina macrocopa*, and potential mechanisms of endocrine disruption." *Chemosphere* 89(1): 10-18.
- Kim, T. H., S. D. Kim, H. Y. Kim, S. J. Lim, M. Lee and S. Yu (2012). "Degradation and toxicity assessment of sulfamethoxazole and chlortetracycline using electron beam, ozone and UV." *Journal of Hazardous Materials* 227: 237-242.

- Kolarova, J., V. Zlabek, R. Grabic, O. Golovko, K. Grabicova, V. Burkina and T. Randak (2013).
 "The effect of environmentally relevant concentration of selected PPCPs on fish cell lines." *Toxicology Letters* 221: S161-S161.
- Komori, K., Y. Suzuki, M. Minamiyama and A. Harada (2013). "Occurrence of selected pharmaceuticals in river water in Japan and assessment of their environmental risk." *Environmental Monitoring and Assessment* 185(6): 4529-4536.
- Kormos, J. L., M. Schulz and T. A. Ternes (2011). "Occurrence of iodinated X-ray contrast media and their biotransformation products in the urban water cycle." *Environmental Science & Technology* 45(20): 8723-8732.
- Kosjek, T. and E. Heath (2011). "Occurrence, fate and determination of cytostatic pharmaceuticals in the environment." *Trac-Trends in Analytical Chemistry* 30(7): 1065-1087.
- Kristiansson, E. (2012). "Diclofenac affects global hepatic gene expression in fish at plasma concentrations similar to human therapeutic levels." *ArrayExpress Archive*.
- Kristoff, G., L. C. Cacciatore, N. R. Verrengia Guerrero and A. C. Cochon (2011). "Effects of the organophosphate insecticide azinphos-methyl on the reproduction and cholinesterase activity of *Biomphalaria glabrata*." *Chemosphere* 84(5): 585-591.
- Kristoff, G., D. Chiny Barrionuevo, L. C. Cacciatore, N. R. Verrengia Guerrero and A. C. Cochon (2012). "*In vivo* studies on inhibition and recovery of B-esterase activities in *Biomphalaria* glabrata exposed to "azinphos-methyl: Analysis of enzyme, substrate and tissue dependence." *Aquatic Toxicology* 112: 19-26.
- Kunkel, U. and M. Radke (2012). "Fate of pharmaceuticals in rivers: Deriving a benchmark dataset at favorable attenuation conditions." *Water Research* 46(17): 5551-5565.
- Lahti, M., J. M. Brozinski, A. Jylha, L. Kronberg and A. Oikari (2011). "Uptake from water, biotransformation, and biliary excretion of pharmaceuticals by rainbow trout." *Environmental Toxicology and Chemistry* 30(6): 1403-1411.
- Lahti, M., J. M. Brozinski, H. Segner, L. Kronberg and A. Oikari (2012). "Bioavailability of pharmaceuticals in waters close to wastewater treatment plants: Use of fish bile for exposure assessment." *Environmental Toxicology and Chemistry* 31(8): 1831-1837.
- Lahti, M. and A. Oikari (2011). "Microbial transformation of pharmaceuticals naproxen, bisoprolol, and diclofenac in aerobic and anaerobic environments." *Archives of Environmental Contamination and Toxicology* 61(2): 202-210.
- Lamichhane, K., S. N. Garcia, D. B. Huggett, D. L. DeAngelis and T. W. La Point (2013). "Chronic effects of carbamazepine on life-history strategies of *Ceriodaphnia dubia* in three successive generations." *Archives of Environmental Contamination and Toxicology* 64(3): 427-438.
- Lang, J. and L. Kohidai (2012). "Effects of the aquatic contaminant human pharmaceuticals and their mixtures on the proliferation and migratory responses of the bioindicator freshwater ciliate *Tetrahymena*." *Chemosphere* 89(5): 592-601.
- Lange, A., Y. Katsu, S. Miyagawa, Y. Ogino, H. Urushitani, T. Kobayashi, T. Hirai, J. A. Shears, M. Nagae, J. Yamamoto, Y. Ohnishi, T. Oka, N. Tatarazako, Y. Ohta, C. R. Tyler and T. Iguchi (2012). "Comparative responsiveness to natural and synthetic estrogens of fish species commonly used in the laboratory and field monitoring." *Aquatic Toxicology* 109: 250-258.
- Langford, K. and K. V. Thomas (2011). "Input of selected human pharmaceutical metabolites into the Norwegian aquatic environment." *Journal of Environmental Monitoring* 13(2): 416-421.

- Lapworth, D. J., N. Baran, M. E. Stuart and R. S. Ward (2012). "Emerging organic contaminants in groundwater: A review of sources, fate and occurrence." *Environmental Pollution* 163: 287-303.
- Lawrence, J. R., B. Zhu, G. D. W. Swerhone, J. Roy, V. Tumber, M. J. Waiser, E. Topp and D. R. Korber (2012). "Molecular and microscopic assessment of the effects of caffeine, acetaminophen, diclofenac, and their mixtures on river biofilm communities." *Environmental Toxicology and Chemistry* 31(3): 508-517.
- Lazzara, R., M. Blazquez, C. Porte and C. Barata (2012). "Low environmental levels of fluoxetine induce spawning and changes in endogenous estradiol levels in the zebra mussel *Dreissena polymorpha*." *Aquatic Toxicology* 106: 123-130.
- Le, T.-H., E.-S. Lim, S.-K. Lee, J.-S. Park, Y.-H. Kim and J. Min (2011). "Toxicity Evaluation of Verapamil and Tramadol Based on Toxicity Assay and Expression Patterns of Dhb, Vtg, Arnt, CYP4, and CYP314 in *Daphnia magna*." *Environmental Toxicology* 26(5): 515-523.
- Lee, H. J., E. Lee, S. H. Yoon, H. R. Chang, K. Kim and J. H. Kwon (2012). "Enzymatic and microbial transformation assays for the evaluation of the environmental fate of diclofenac and its metabolites." *Chemosphere* 87(8): 969-974.
- Lee, J., K. Ji, Y. L. Kho, P. Kim and K. Choi (2011). "Chronic exposure to diclofenac on two freshwater cladocerans and Japanese medaka." *Ecotoxicology and Environmental Safety* 74(5): 1216-1225.
- Lee, S. H., H. R. Kim, R. X. Han, R. K. Oqani and D. I. Jin (2013). "Cardiovascular risk assessment of atypical antipsychotic drugs in a zebrafish model." *Journal of Applied Toxicology* 33(6): 466-470.
- Lei, B., J. Kang, Y. Yu, J. Zha, W. Li and Z. Wang (2013). "Beta-estradiol 17-valerate affects embryonic development and sexual differentiation in Japanese medaka (*Oryzias latipes*)." *Aquatic toxicology (Amsterdam, Netherlands)* 134-135: 128-134.
- Leitner, J. (2011). "Melting point of nanoparticles." Chemicke Listy 105(3): 174-185.
- Leston, S., M. Nunes, I. Viegas, F. Ramos and M. A. Pardal (2013). "The effects of chloramphenicol on *Ulva lactuca.*" *Chemosphere* 91(4): 552-557.
- Lewis, G., A. Juhasz and E. Smith (2012). "Environmental metabolites of fluoroquinolones: synthesis, fractionation and toxicological assessment of some biologically active metabolites of ciprofloxacin." *Environmental Science and Pollution Research* 19(7): 2697-2707.
- Li, W. Z., Y. Ding, Q. Sui, S. G. Lu, Z. F. Qiu and K. F. Lin (2012). "Identification and ecotoxicity assessment of intermediates generated during the degradation of clofibric acid by advanced oxidation processes." *Frontiers of Environmental Science & Engineering* 6(4): 445-454.
- Li, Z. H., G. H. Lu, X. F. Yang and C. Wang (2012). "Single and combined effects of selected pharmaceuticals at sublethal concentrations on multiple biomarkers in *Carassius auratus*." *Ecotoxicology* 21(2): 353-361.
- Li, Z. H., V. Zlabek, J. Velisek, R. Grabic, J. Machova, J. Kolarova, P. Li and T. Randak (2011). "Acute toxicity of carbamazepine to juvenile rainbow trout (*Oncorhynchus mykiss*): Effects on antioxidant responses, hematological parameters and hepatic EROD." *Ecotoxicology and Environmental Safety* 74(3): 319-327.
- Lim, S. J., E. Jang, S. H. Lee, B. H. Yoo, S. K. Kim and T. H. Kim (2013). "Antibiotic resistance in bacteria isolated from freshwater aquacultures and prediction of the persistence and

toxicity of antimicrobials in the aquatic environment." *Journal of Environmental Science and Health Part B-Pesticides Food Contaminants and Agricultural Wastes* 48(6): 495-504.

- Lim, S. J., C. K. Seo, T. H. Kim and S. W. Myung (2013). "Occurrence and ecological hazard assessment of selected veterinary medicines in livestock wastewater treatment plants." *Journal of Environmental Science and Health Part B-Pesticides Food Contaminants and Agricultural Wastes* 48(8): 658-670.
- Liu, B. Y., W. Q. Liu, X. P. Nie, C. Guan, Y. F. Yang, Z. H. Wang and W. Liao (2011). "Growth response and toxic effects of three antibiotics on *Selenastrum capricornutum* evaluated by photosynthetic rate and chlorophyll biosynthesis." *Journal of Environmental Sciences-China* 23(9): 1558-1563.
- Liu, B. Y., X. P. Nie, W. Q. Liu, P. Snoeijs, C. Guan and M. T. K. Tsui (2011). "Toxic effects of erythromycin, ciprofloxacin and sulfamethoxazole on photosynthetic apparatus in *Selenastrum capricornutum*." *Ecotoxicology and Environmental Safety* 74(4): 1027-1035.
- Liu, F., J. S. Wu, G. G. Ying, Z. X. Luo and H. Feng (2012). "Changes in functional diversity of soil microbial community with addition of antibiotics sulfamethoxazole and chlortetracycline." *Applied Microbiology and Biotechnology* 95(6): 1615-1623.
- Liu, J.-L. and M.-H. Wong (2013). "Pharmaceuticals and personal care products (PPCPs): A review on environmental contamination in China." *Environment International* 59: 208-224.
- Liu, L., Y. H. Liu, C. X. Liu, Z. Wang, J. Dong, G. F. Zhu and X. Huang (2013). "Potential effect and accumulation of veterinary antibiotics in *Phragmites australis* under hydroponic conditions." *Ecological Engineering* 53: 138-143.
- Liu, X., J. Lee, K. Ji, S. Takeda and K. Choi (2012). "Potentials and mechanisms of genotoxicity of six pharmaceuticals frequently detected in freshwater environment." *Toxicology Letters* 211(1): 70-76.
- Liu, Y., X. Wang, L. Qiu, X. Li and C. Wang (2011). "Acute toxicity of seven pyrethroid insecticides to brine shrimp *Artemia*." *Asian Journal of Ecotoxicology* 6(5): 557-560.
- Lopez-Doval, J. C., J. V. K. Kukkonen, P. Rodrigo and I. Munoz (2012). "Effects of indomethacin and propranolol on *Chironomus riparius* and *Physella* (*Costatella*) *acuta*." *Ecotoxicology and Environmental Safety* 78: 110-115.
- Lu, G. H., Z. H. Yan, Y. H. Wang and W. Chen (2011). "Assessment of estrogenic contamination and biological effects in Lake Taihu." *Ecotoxicology* 20(5): 974-981.
- Lumaret, J. P., F. Errouissi, K. Floate, J. Rombke and K. Wardhaugh (2012). "A review on the toxicity and non-target effects of macrocyclic lactones in terrestrial and aquatic environments." *Current Pharmaceutical Biotechnology* 13(6): 1004-1060.
- Luna-Acosta, A., T. Renault, H. Thomas-Guyon, N. Faury, D. Saulnier, H. Budzinski, K. Le Menach, P. Pardon, I. Fruitier-Arnaudin and P. Bustamante (2012). "Detection of early effects of a single herbicide (diuron) and a mix of herbicides and pharmaceuticals (diuron, isoproturon, ibuprofen) on immunological parameters of Pacific oyster (*Crassostrea gigas*) spat." *Chemosphere* 87(11): 1335-1340.
- Lyons, M., K. Mackeigan, D. Wong, W. Fairchild and L. Burridge (2011). "Effects of 4nonylphenol, cypermethrin, deltamethrin, glyphosate, imidacloprid and mancozeb on growth of Atlantic salmon (*Salmo salar* L.) (PO)." *Canadian Technical Report of Fisheries and Aquatic Sciences* 2949: 66.

- Ma, X. Y. and X. C. Wang (2013). "Ecotoxicity comparison of organic contaminants and heavy metals using *Vibrio-qinghaiensis* sp.-Q67." *Water Science and Technology* 67(10): 2221-2227.
- Madureira, T. V., C. Cruzeiro, M. J. Rocha and E. Rocha (2011). "The toxicity potential of pharmaceuticals found in the Douro River estuary (Portugal)-Experimental assessment using a zebrafish embryo test." *Environmental Toxicology and Pharmacology* 32(2): 212-217.
- Madureira, T. V., M. J. Rocha, C. Cruzeiro, M. H. Galante, R. A. F. Monteiro and E. Rocha (2011).
 "The toxicity potential of pharmaceuticals found in the Douro River estuary (Portugal): Assessing impacts on gonadal maturation with a histopathological and stereological study of zebrafish ovary and testis after sub-acute exposures." *Aquatic Toxicology* 105(3-4): 292-299.
- Maeng, S. K., S. K. Sharma, K. Lekkerkerker-Teunissen and G. L. Amy (2011). "Occurrence and fate of bulk organic matter and pharmaceutically active compounds in managed aquifer recharge: A review." *Water Research* 45(10): 3015-3033.
- Magdeburg, A., D. Stalter and J. Oehlmann (2012). "Whole effluent toxicity assessment at a wastewater treatment plant upgraded with a full-scale post-ozonation using aquatic key species." *Chemosphere* 88(8): 1008-1014.
- Malarvizhi, A., C. Kavitha, M. Saravanan and M. Ramesh (2012). "Carbamazepine (CBZ) induced enzymatic stress in gill, liver and muscle of a common carp, *Cyprinus carpio*." *Journal of King Saud University Science* 24(2): 179-186.
- Mankes, R. F. and C. D. Silver (2013). "Quantitative study of controlled substance bedside wasting, disposal and evaluation of potential ecologic effects." *Science of the Total Environment* 444: 298-310.
- Martin, J., D. Camacho-Munoz, J. L. Santos, I. Aparicio and E. Alonso (2011). "Monitoring of pharmaceutically active compounds on the Guadalquivir River basin (Spain): occurrence and risk assessment." *Journal of Environmental Monitoring* 13(7): 2042-2049.
- Martin, J., D. Camacho-Munoz, J. L. Santos, I. Aparicio and E. Alonso (2012). "Occurrence of pharmaceutical compounds in wastewater and sludge from wastewater treatment plants: Removal and ecotoxicological impact of wastewater discharges and sludge disposal." *Journal of Hazardous Materials* 239: 40-47.
- Martini, F., J. V. Tarazona and M. V. Pablos (2012). "Are fish and standardized FETAX assays protective enough for amphibians? A case study on *Xenopus laevis* larvae assay with biologically active substances present in livestock wastes." *Scientific World Journal*.
- Martins, N., R. Pereira, N. Abrantes, J. Pereira, F. Goncalves and C. R. Marques (2012). "Ecotoxicological effects of ciprofloxacin on freshwater species: data integration and derivation of toxicity thresholds for risk assessment." *Ecotoxicology* 21(4): 1167-1176.
- Maruya, K. A., D. E. Vidal-Dorsch, S. M. Bay, J. W. Kwon, K. Xia and K. L. Armbrust (2012).
 "Organic contaminants of emerging concern in sediments and flatfish collected near outfalls discharging treated wastewater effluent to the Southern California Bight." *Environmental Toxicology and Chemistry* 31(12): 2683-2688.
- Massarsky, A., V. L. Trudeau and T. W. Moon (2011). "Beta-blockers as endocrine disruptors: The potential effects of human beta-blockers on aquatic organisms." *Journal of Experimental Zoology Part a-Ecological Genetics and Physiology* 315A(5): 251-265.

- Matamoros, V., C. A. Arias, L. X. Nguyen, V. Salvado and H. Brix (2012). "Occurrence and behavior of emerging contaminants in surface water and a restored wetland." *Chemosphere* 88(9): 1083-1089.
- Matamoros, V., L. X. Nguyen, C. A. Arias, V. Salvado and H. Brix (2012). "Evaluation of aquatic plants for removing polar microcontaminants: a microcosm experiment." *Chemosphere* 88(10): 1257-1264.
- Materne, E.-M., A. G. Tonevitsky and U. Marx (2013). "Chip-based liver equivalents for toxicity testing organotypicalness versus cost-efficient high throughput." *Lab on a Chip* 13(18): 3481-3495.
- Mathews, S. and D. Reinhold (2013). "Biosolid-borne tetracyclines and sulfonamides in plants." *Environmental Science and Pollution Research* 20(7): 4327-4338.
- Matozzo, V., A. Formenti, G. Donadello and M. G. Marin (2012). "A multi-biomarker approach to assess effects of triclosan in the clam *Ruditapes philippinarum*." *Marine Environmental Research* 74: 40-46.
- Matozzo, V., S. Rova and M. G. Marin (2012). "The nonsteroidal anti-inflammatory drug, ibuprofen, affects the immune parameters in the clam *Ruditapes philippinarum*." *Marine Environmental Research* 79: 116-121.
- Meanwell, N. A. (2011). "Improving drug candidates by design: A focus on physicochemical properties as a means of improving compound disposition and safety." *Chemical Research in Toxicology* 24(9): 1420-1456.
- Meina, E. G., A. Lister, T. Bosker, M. Servos, K. Munkittrick and D. MacLatchy (2013). "Effects of 17 alpha-ethinylestradiol (EE2) on reproductive endocrine status in mummichog (*Fundulus heteroclitus*) under differing salinity and temperature conditions." *Aquatic Toxicology* 134: 92-103.
- Meinertz, J. R., T. M. Schreier and J. A. Bernardy (2011). "Chronic toxicity of erythromycin thiocyanate to *Daphnia magna* in a flow-through, continuous exposure test system." *Bulletin of Environmental Contamination and Toxicology* 87(6): 621-625.
- Memmert, U., A. Peither, R. Burri, K. Weber, T. Schmidt, J. P. Sumpter and A. Hartmann (2013). "Diclofenac: new data on chronic toxicity and bioconcentration in fish." *Environmental Toxicology and Chemistry* 32(2): 442-452.
- Mennigen, J. A., P. Stroud, J. M. Zamora, T. W. Moon and V. L. Trudeau (2011).
 "Pharmaceuticals as neuroendocrine disruptors: Lessons learned from fish on prozac." *Journal of Toxicology and Environmental Health-Part B-Critical Reviews* 14(5-7): 387-412.
- Meredith-Williams, M., L. J. Carter, R. Fussell, D. Raffaelli, R. Ashauer and A. B. A. Boxall (2012). "Uptake and depuration of pharmaceuticals in aquatic invertebrates." *Environmental Pollution* 165: 250-258.
- Michelini, L., F. Meggio, N. La Rocca, S. Ferro and R. Ghisi (2012). "Accumulation and effects of sulfamethoxine in *Salix fragalis* L. plants: a preliminary study on phytoremediation purposes." *International Journal of Phytoremediation* 14(4): 388-402.
- Milan, M., M. Pauletto, T. Patarnello, L. Bargelloni, M. G. Marin and V. Matozzo (2013). "Gene transcription and biomarker responses in the clam *Ruditapes philippinarum* after exposure to ibuprofen." *Aquatic Toxicology* 126: 17-29.

- Miller, H. D., B. W. Clark, D. E. Hinton, A. Whitehead, S. Martin, K. W. Kwok and S. W. Kullman (2012). "Anchoring ethinylestradiol induced gene expression changes with testicular morphology and reproductive function in the medaka." *PloS one* 7(12): e52479.
- Mishra, N. K. (2011). "Computational modeling of P450s for toxicity prediction." *Expert Opinion* on Drug Metabolism & Toxicology 7(10): 1211-1231.
- Mompelat, S., A. Jaffrezic, E. Jarde and B. Le Bot (2013). "Storage of natural water samples and preservation techniques for pharmaceutical quantification." *Talanta* 109: 31-45.
- Montanha, F. P., L. A. G. Galeb, J. D. Mikos, L. N. Ganeco, T. P. Pereira, A. Tanaka, P. G. Kirschnik and C. T. Pimpao (2012). "Pyrethroid toxicity in silver catfish, *Rhamdia quelen*." *Pesquisa Veterinaria Brasileira* 32(12): 1297-1303.
- Munari, M., V. Matozzo, G. Chemello and M. G. Marin (2012). "Combined effects of seawater acidification and diclofenac on immune parameters of the clam *Ruditapes philippinarum*." *Comparative Biochemistry and Physiology a-Molecular & Integrative Physiology* 163(1): S5-S5.
- Na, G., X. Fang, Y. Cai, L. Ge, H. Zong, X. Yuan, Z. Yao and Z. Zhang (2013). "Occurrence, distribution, and bioaccumulation of antibiotics in coastal environment of Dalian, China." *Marine Pollution Bulletin* 69(1-2): 233-237.
- Nallani, G. C., P. M. Paulos, L. A. Constantine, B. J. Venables and D. B. Huggett (2011).
 "Bioconcentration of ibuprofen in fathead minnow (*Pimephales promelas*) and channel catfish (*Ictalurus punctatus*)." *Chemosphere* 84(10): 1371-1377.
- Narvaez, J. F. V. and C. Jimenez C (2012). "Pharmaceutical products in the environment: sources, effects and risks." *Vitae-Revista De La Facultad De Quimica Farmaceutica* 19(1): 93-108.
- Neamtu, M., M. Bobu, A. Kettrup and I. Siminiceanu (2013). "Ozone photolysis of paracetamol in aqueous solution." *Journal of Environmental Science and Health Part a-Toxic/Hazardous Substances & Environmental Engineering* 48(10): 1264-1271.
- Nie, X. P., B. Y. Liu, H. J. Yu, W. Q. Liu and Y. F. Yang (2013). "Toxic effects of erythromycin, ciprofloxacin and sulfamethoxazole exposure to the antioxidant system in *Pseudokirchneriella subcapitata*." *Environmental Pollution* 172: 23-32.
- Notch, E. G. and G. D. Mayer (2013). "Impact of environmental estrogens on nucleotide excision repair gene expression in embryonic zebrafish." *Comparative biochemistry and physiology. Toxicology & pharmacology : CBP* 157(4): 361-365.
- Oganesova, Y. V. and O. F. Filenko (2012). "Comparative toxicity of azine dyes to gastropod mollusks (*Planorbis corneus*)." *Toksikologicheskii Vestnik*(6): 43-48.
- Oldenkamp, R., M. A. J. Huijbregts, A. Hollander, A. Versporten, H. Goossens and A. M. J. Ragas (2013). "Spatially explicit prioritization of human antibiotics and antineoplastics in Europe." *Environment International* 51: 13-26.
- Oliveira, C., J. Almeida, L. Guilhermino, A. M. V. M. Soares and C. Gravato (2012). "Acute effects of deltamethrin on swimming velocity and biomarkers of the common prawn *Palaemon serratus*." *Aquatic Toxicology* 124: 209-216.
- Opris, O., F. Copaciu, M. L. Soran, D. Ristoiu, U. Niinemets and L. Copolovici (2012). "Influence of nine antibiotics on key secondary metabolites and physiological characteristics in *Triticum aestivum*: Leaf volatiles as a promising new tool to assess toxicity." *Ecotoxicology and Environmental Safety* 87: 70-79.

- Orias, F. and Y. Perrodin (2013). "Characterisation of the ecotoxicity of hospital effluents: a review." *Science of the Total Environment* 454: 250-276.
- Oskarsson, H., A. K. E. Wiklund, K. Lindh and L. Kumblad (2012). "Effect studies of human pharmaceuticals on *Fucus vesiculosus* and *Gammarus spp.*" *Marine Environmental Research* 74: 1-8.
- Overturf, M. D., C. L. Overturf, D. Baxter, D. N. Hala, L. Constantine, B. Venables and D. B. Huggett (2012). "Early life-stage toxicity of eight pharmaceuticals to the fathead minnow, *Pimephales promelas*." *Archives of Environmental Contamination and Toxicology* 62(3): 455-464.
- Ozdemir, A., M. Duran and A. Sen (2011). "Potential use of the oligochaete *Limnodrilus profundicola* V., as a bioindicator of contaminant exposure." *Environmental Toxicology* 26(1): 37-44.
- Ozkan, O. and O. Ustuner (2012). "Investigations about genotoxicity of deltamethrin." *Kafkas Universitesi Veteriner Fakultesi Dergisi* 18(1): 69-74.
- Paiga, P., L. Santos, C. G. Amorim, A. N. Araujo, M. Montenegro, A. Pena and C. Delerue-Matos (2013). "Pilot monitoring study of ibuprofen in surface waters of north of Portugal." *Environmental Science and Pollution Research* 20(4): 2410-2420.
- Park, K. and I.-S. Kwak (2012). "Gene expression of ribosomal protein mRNA in *Chironomus riparius*: Effects of endocrine disruptor chemicals and antibiotics." *Comparative Biochemistry and Physiology C-Toxicology & Pharmacology* 156(2): 113-120.
- Parolini, M. and A. Binelli (2012). "Sub-lethal effects induced by a mixture of three non-steroidal anti-inflammatory drugs (NSAIDs) on the freshwater bivalve *Dreissena polymorpha*." *Ecotoxicology* 21(2): 379-392.
- Parolini, M., A. Binelli and A. Provini (2011). "Chronic effects induced by ibuprofen on the freshwater bivalve *Dreissena polymorpha*." *Ecotoxicology and Environmental Safety* 74(6): 1586-1594.
- Parolini, M., A. Pedriali and A. Binelli (2013). "Application of a biomarker response index for ranking the toxicity of five pharmaceutical and personal care products (PPCPs) to the bivalve *Dreissena polymorpha*." *Archives of Environmental Contamination and Toxicology* 64(3): 439-447.
- Parolini, M., B. Quinn, A. Binelli and A. Provini (2011). "Cytotoxicity assessment of four pharmaceutical compounds on the zebra mussel (*Dreissena polymorpha*) haemocytes, gill and digestive gland primary cell cultures." *Chemosphere* 84(1): 91-100.
- Parrott, J., S. Tabe, V. Pileggi, N. Feisthauer, S. Kleywegt, J. Schroeder, P. Seto and S. Backus (2011). "Fathead minnow lifecycle exposure to three Canadian municipal wastewater effluents." *Canadian Technical Report of Fisheries and Aquatic Sciences* 2949: 92-93.
- Pereira, L. A., I. C. Sales Fontes Jardim, A. H. Fostier and S. Rath (2012). "Occurrence, behavior and environemtnal impacts caused by the presence of veterinary antimicrobials in soils." *Quimica Nova* 35(1): 159-169.
- Perez, M. R., J. I. Fernandino, P. Carriquiriborde and G. M. Somoza (2012). "Feminization and altered gonadal gene expression profile by ethinylestradiol exposure to pejerrey, *Odontesthes bonariensis*, a South American teleost fish." *Environmental Toxicology and Chemistry* 31(5): 941-946.

- Perron, M. M., K. T. Ho, M. G. Cantwell, R. M. Burgess and M. C. Pelletier (2012). "Effects of triclosan on marine benthic and epibenthic organisms." *Environmental Toxicology and Chemistry* 31(8): 1861-1866.
- Peuravuori, J. (2012). "Aquatic photochemistry of paracetamol in the presence of dissolved organic chromophoric material and nitrate." *Environmental Science and Pollution Research* 19(6): 2259-2270.
- Pile, J. C. (2011). "Antimicrobial stewardship: optimizing antibiotic use in an era of increasing resistance and rising costs." *Journal of Hospital Medicine* 6: S1-S3.
- Plumlee, M. H., C. J. Gurr and M. Reinhard (2012). "Recycled water for stream flow augmentation: Benefits, challenges, and the presence of wastewater-derived organic compounds." *Science of the Total Environment* 438: 541-548.
- Praskova, E., E. Voslarova, Z. Siroka, S. Macova, L. Plhalova, I. Bedanova, P. Marsalek, V. Pistekova and Z. Svobodova (2011). "Comparison of acute toxicity of ketoprofen to juvenile and embryonic stages of *Danio rerio*." *Neuroendocrinology Letters* 32: 117-120.
- Praskova, E., E. Voslarova, Z. Siroka, L. Plhalova, S. Macova, P. Marsalek, V. Pistekova and Z. Svobodova (2011). "Assessment of diclofenac LC50 reference values in juvenile and embryonic stages of the zebrafish (*Danio rerio*)." *Polish Journal of Veterinary Sciences* 14(4): 545-549.
- Praskova, E., D. Zivna, S. Stepanova, M. Sevcikova, J. Blahova, P. Marsalek, Z. Siroka, E. Voslarova and Z. Svobodova (2012). "Acute toxicity of acetylsalicylic acid to juvenile and embryonic stages of *Danio rerio.*" *Neuro endocrinology letters* 33 Suppl 3: 72-76.
- Proia, L., S. Morin, M. Peipoch, A. M. Romani and S. Sabater (2011). "Resistance and recovery of river biofilms receiving short pulses of triclosan and diuron." *Science of the Total Environment* 409(17): 3129-3137.
- Proia, L., V. Osorio, S. Soley, M. Kock-Schulmeyer, S. Perez, D. Barcelo, A. M. Romani and S. Sabater (2013). "Effects of pesticides and pharmaceuticals on biofilms in a highly impacted river." *Environmental pollution (Barking, Essex : 1987)* 178: 220-228.
- Pruvot, B., Y. Quiroz, A. Voncken, N. Jeanray, A. Piot, J. A. Martial and M. Muller (2012). "A panel of biological tests reveals developmental effects of pharmaceutical pollutants on late stage zebrafish embryos." *Reproductive Toxicology* 34(4): 568-583.
- Quinlan, E. L., C. T. Nietch, K. Blocksom, J. M. Lazorchak, A. L. Batt, R. Griffiths and D. J. Klemm (2011). "Temporal dynamics of periphyton exposed to tetracycline in stream mesocosms." *Environmental Science & Technology* 45(24): 10684-10690.
- Quinn, B. (2012). "Chronic toxicity of pharmaceutical compounds on bivalve mollusc primary cultures." *Toxicology letters* 211: S89-S89.
- Quinn, B., W. Schmidt, K. O'Rourke and R. Hernan (2011). "Effects of the pharmaceuticals gemfibrozil and diclofenac on biomarker expression in the zebra mussel (*Dreissena polymorpha*) and their comparison with standardised toxicity tests." *Chemosphere* 84(5): 657-663.
- Ragugnetti, M., M. L. Adams, A. T. B. Guimaraes, G. Sponchiado, E. C. de Vasconcelos and C. M.
 R. de Oliveira (2011). "Ibuprofen genotoxicity in aquatic environment: An experimental model using *Oreochromis niloticus.*" *Water Air and Soil Pollution* 218(1-4): 361-364.
- Ramos, A. S., F. Goncalves, S. C. Antunes and B. Nunes (2012). "Cholinesterase characterization in *Corbicula fluminea* and effects of relevant environmental contaminants: A pesticide

(chlorfenvinphos) and a detergent (SDS)." *Journal of Environmental Science and Health Part B-Pesticides Food Contaminants and Agricultural Wastes* 47(6): 512-519.

- Rana, S. M., M. R. Asi, F. Niazi, S. Sultana, Ghazala and K. A. Al-Ghanim (2011). "Determination of organochlorine and nitrogen containing pesticide residues in *Labeo rohita*." *Toxicological and Environmental Chemistry* 93(10): 1851-1855.
- Reh, R., T. Licha, T. Geyer, K. Nodler and M. Sauter (2013). "Occurrence and spatial distribution of organic micro-pollutants in a complex hydrogeological karst system during low flow and high flow periods, results of a two-year study." *Science of the Total Environment* 443: 438-445.
- Reyhanian, N., K. Volkova, S. Hallgren, T. Bollner, P. E. Olsson, H. Olsen and I. P. Hallstrom (2011). "17 alpha-Ethinyl estradiol affects anxiety and shoaling behavior in adult male zebra fish (*Danio rerio*)." *Aquatic Toxicology* 105(1-2): 41-48.
- Ribeiro, A. R., P. M. L. Castro and M. E. Tiritan (2012). "Chiral pharmaceuticals in the environment." *Environmental Chemistry Letters* 10(3): 239-253.
- Ribeiro, C., R. Urbatzka, L. F. C. Castro, J. Carrola, A. Fontainhas-Fernandes, R. A. F. Monteiro, E. Rocha and M. J. Rocha (2012). "In vitro exposure of Nile tilapia (*Oreochromis niloticus*) testis to estrogenic endocrine disrupting chemicals: mRNA expression of genes encoding steroidogenic enzymes." *Toxicology Mechanisms and Methods* 22(1): 47-53.
- Richards, N. L., G. Cook, V. Simpson, S. Hall, N. Harrison and K. S. Scott (2011). "Qualitative detection of the NSAIDs diclofenac and ibuprofen in the hair of Eurasian otters (*Lutra lutra*) occupying UK waterways with GC-MS." *European Journal of Wildlife Research* 57(5): 1107-1114.
- Rico, A., Y. Geng, A. Focks and P. J. Van den Brink (2013). "Modeling environmental and human health risks of veterinary medicinal products applied in pond aquaculture." *Environmental Toxicology and Chemistry* 32(5): 1196-1207.
- Riebeling, C., R. Pirow, K. Becker, R. Buesen, D. Eikel, J. Kaltenhauser, F. Meyer, H. Nau, B. Slawik, A. Visan, J. Volland, H. Spielmann, A. Luch and A. Seiler (2011). "The embryonic stem cell test as tool to assess structure-dependent teratogenicity: The case of valproic acid." *Toxicological Sciences* 120(2): 360-370.
- Riva, C., S. Cristoni and A. Binelli (2012). "Effects of triclosan in the freshwater mussel *Dreissena polymorpha*: a proteomic investigation." *Aquatic Toxicology* 118: 62-71.
- Rivadeneira, P. R., M. Agrelo, S. Otero and G. Kristoff (2013). "Different effects of subchronic exposure to low concentrations of the organophosphate insecticide chlorpyrifos in a freshwater gastropod." *Ecotoxicology and Environmental Safety* 90: 82-88.
- Rocco, L., G. Frenzilli, G. Zito, A. Archimandritis, C. Peluso and V. Stingo (2012). "Genotoxic effects in fish induced by pharmacological agents present in the sewage of some Italian water-treatment plants." *Environmental Toxicology* 27(1): 18-25.
- Rocco, L., C. Peluso and V. Stingo (2012). "Micronucleus test and comet assay for the evaluation of zebrafish genomic damage induced by erythromycin and lincomycin." *Environmental Toxicology* 27(10): 598-604.
- Rosi-Marshall, E. J., D. W. Kincaid, H. A. Bechtold, T. V. Royer, M. Rojas and J. J. Kelly (2013). "Pharmaceuticals suppress algal growth and microbial respiration and alter bacterial communities in stream biofilms." *Ecological Applications* 23(3): 583-593.

- Salgado, R., V. J. Pereira, G. Carvalho, R. Soeiro, V. Gaffney, C. Almeida, V. V. Cardoso, E. Ferreira, M. J. Benoliel, T. A. Ternes, A. Oehmen, M. A. M. Reis and J. P. Noronha (2013).
 "Photodegradation kinetics and transformation products of ketoprofen, diclofenac and atenolol in pure water and treated wastewater." *Journal of Hazardous Materials* 244: 516-527.
- Saravanan, M., K. U. Devi, A. Malarvizhi and M. Ramesh (2012). "Effects of Ibuprofen on hematological, biochemical and enzymological parameters of blood in an Indian major carp, *Cirrhinus mrigala*." *Environmental Toxicology and Pharmacology* 34(1): 14-22.
- Saravanan, M., S. Karthika, A. Malarvizhi and M. Ramesh (2011). "Ecotoxicological impacts of clofibric acid and diclofenac in common carp (*Cyprinus carpio*) fingerlings: Hematological, biochemical, ionoregulatory and enzymological responses." *Journal of Hazardous Materials* 195: 188-194.
- Sarria, M. P., M. M. Santos, M. A. Reis-Henriques, N. M. Vieira and N. M. Monteiro (2011). "The unpredictable effects of mixtures of androgenic and estrogenic chemicals on fish early life." *Environment International* 37(2): 418-424.
- Sarria, M. P., J. Soares, M. N. Vieira, L. Filipe, C. Castro, M. M. Santos and N. M. Monteiro (2011). "Rapid-behaviour responses as a reliable indicator of estrogenic chemical toxicity in zebrafish juveniles." *Chemosphere* 85(10): 1543-1547.
- Schmidt, W., K. O'Rourke, R. Hernan and B. Quinn (2011). "Effects of the pharmaceuticals gemfibrozil and diclofenac on the marine mussel (*Mytilus spp.*) and their comparison with standardized toxicity tests." *Marine Pollution Bulletin* 62(7): 1389-1395.
- Scholz, S., P. Renner, S. E. Belanger, F. Busquet, R. Davi, B. A. Demeneix, J. S. Denny, M. Leonard, M. E. McMaster, D. L. Villeneuve and M. R. Embry (2013). "Alternatives to *in vivo* tests to detect endocrine disrupting chemicals (EDCs) in fish and amphibians screening for estrogen, androgen and thyroid hormone disruption." *Critical Reviews in Toxicology* 43(1): 45-72.
- Schultz, M. M., M. M. Painter, S. E. Bartell, A. Logue, E. T. Furlong, S. L. Werner and H. L. Schoenfuss (2011). "Selective uptake and biological consequences of environmentally relevant antidepressant pharmaceutical exposures on male fathead minnows." *Aquatic Toxicology* 104(1-2): 38-47.
- Selderslaghs, I. W. T., R. Blust and H. E. Witters (2012). "Feasibility study of the zebrafish assay as an alternative method to screen for developmental toxicity and embryotoxicity using a training set of 27 compounds." *Reproductive Toxicology* 33(2): 142-154.
- Senta, I., S. Terzic and M. Ahel (2013). "Occurrence and fate of dissolved and particulate antimicrobials in municipal wastewater treatment." *Water Research* 47(2): 705-714.
- Shen, M.-F., A. Kumar, S.-Y. Ding and S. Grocke (2012). "Comparative study on the toxicity of pyrethroids, alpha-cypermethrin and deltamethrin to *Ceriodaphnia dubia*." *Ecotoxicology and Environmental Safety* 78: 9-13.
- Shi, H. H., Z. Sun, Z. Liu and Y. G. Xue (2012). "Effects of clotrimazole and amiodarone on early development of amphibian (*Xenopus tropicalis*)." *Toxicological and Environmental Chemistry* 94(1): 128-135.
- Shono, F., M. Kumamoto and K. Inoue (2013). "Trends of activities in international organization and authorities in each country regarding risk management of nanomaterials." *Yakugaku Zasshi-Journal of the Pharmaceutical Society of Japan* 133(2): 157-167.

- Siegrist, H. and A. Joss (2012). "Review on the fate of organic micropollutants in wastewater treatment and water reuse with membranes." *Water Science and Technology* 66(6): 1369-1376.
- Silva, L. J. G., C. M. Lino, L. M. Meisel and A. Pena (2012). "Selective serotonin re-uptake inhibitors (SSRIs) in the aquatic environment: an ecopharmacovigilance approach." *Science of the Total Environment* 437: 185-195.
- Silva, P., M. J. Rocha, C. Cruzeiro, F. Malhao, B. Reis, R. Urbatzka, R. A. F. Monteiro and E. Rocha (2012). "Testing the effects of ethinylestradiol and of an environmentally relevant mixture of xenoestrogens as found in the Douro River (Portugal) on the maturation of fish gonads-A stereological study using the zebrafish (*Danio rerio*) as model." *Aquatic Toxicology* 124: 1-10.
- Skolness, S. Y., E. J. Durhan, K. M. Jensen, M. D. Kahl, E. A. Makynen, D. L. Villeneuve and G. T. Ankley (2012). "Effects of gemfibrozil on lipid metabolism, steroidogenesis, and reproduction in the fathead minnow (*Pimephales promelas*)." *Environmental Toxicology and Chemistry* 31(11): 2615-2624.
- Slana, M. and M. S. Dolenc (2013). "Environmental risk assessment of antimicrobials applied in veterinary medicine- A field study and laboratory approach." *Environmental Toxicology and Pharmacology* 35(1): 131-141.
- Smith, E. M., F. I. Iftikar, S. Higgins, A. Irshad, R. Jandoc, M. Lee and J. Y. Wilson (2012). "*In vitro* inhibition of cytochrome P450-mediated reactions by gemfibrozil, erythromycin, ciprofloxacin and fluoxetine in fish liver microsomes." *Aquatic Toxicology* 109: 259-266.
- Soffker, M. and C. R. Tyler (2012). "Endocrine disrupting chemicals and sexual behaviors in fish a critical review on effects and possible consequences." *Critical Reviews in Toxicology* 42(8): 653-668.
- Sponchiado, G., E. Reynaldo, A. C. B. de Andrade, E. C. de Vasconcelos, M. L. Adam and C. M. R. de Oliveira (2011). "Genotoxic effects in erythrocytes of *Oreochromis niloticus* exposed to nanograms-per-liter concentration of 17 beta-estradiol (E-2): An assessment using micronucleus test and comet assay." *Water Air and Soil Pollution* 218(1-4): 353-360.
- Stalter, D., A. Magdeburg, K. Quednow, A. Botzat and J. Oehlmann (2013). "Do contaminants originating from state-of-the-art treated wastewater impact the ecological quality of surface waters?" *PloS one* 8(4).
- Stange, D. and J. Oehlmann (2012). "Identification of oestrogen-responsive transcripts in *Potamopyrgus antipodarum.*" *Journal of Molluscan Studies* 78: 337-342.
- Stange, D., A. Sieratowicz, R. Horres and J. Oehlmann (2012). "Freshwater mudsnail (*Potamopyrgus antipodarum*) estrogen receptor: Identification and expression analysis under exposure to (xeno-)hormones." *Ecotoxicology and Environmental Safety* 75: 94-101.
- Stasinakis, A. S., S. Mermigka, V. G. Samaras, E. Farmaki and N. S. Thomaidis (2012).
 "Occurrence of endocrine disrupters and selected pharmaceuticals in Aisonas River (Greece) and environmental risk assessment using hazard indexes." *Environmental Science and Pollution Research* 19(5): 1574-1583.
- Steinbach, C., V. Burkina, G. Federova, K. Grabicova, J. Velisek, V. Zlabek, T. Randak, H. Schmidt-Posthaus, R. Grabic and H. K. Kroupova (2013). "The sub-lethal toxic effects and bioconcentration of the human pharmaceutical atenolol in rainbow trout (*Oncorhynchus mykiss*)." *Toxicology Letters* 221: S60-S60.

- Stepanova, S., E. Praskova, L. Chromcova, L. Plhalova, M. Prokes, J. Blahova and Z. Svobodova (2013). "The effects of diclofenac on early life stages of common carp (*Cyprinus carpio*)." *Environmental Toxicology and Pharmacology* 35(3): 454-460.
- Stuart, M., D. Lapworth, E. Crane and A. Hart (2012). "Review of risk from potential emerging contaminants in UK groundwater." *Science of the Total Environment* 416: 1-21.
- Subedi, B., M. A. Mottaleb, C. K. Chambliss and S. Usenko (2011). "Simultaneous analysis of select pharmaceuticals and personal care products in fish tissue using pressurized liquid extraction combined with silica gel cleanup." *Journal of Chromatography A* 1218(37): 6278-6284.
- Suga, N., M. Ogo and S. Suzuki (2013). "Risk assessment of oxytetracycline in water phase to major sediment bacterial community: a water-sediment microcosm study." *Environmental Toxicology and Pharmacology* 36(1): 142-148.
- Sui, Q., J. Huang, S. B. Deng, W. W. Chen and G. Yu (2011). "Seasonal variation in the occurrence and removal of pharmaceuticals and personal care products in different biological wastewater treatment processes." *Environmental Science & Technology* 45(8): 3341-3348.
- Sun, J. P., W. H. Li, P. Zheng and J. Zhu (2012). "Toxicity evaluation of antibiotics in piggery wastewater by luminescent bacteria." *Polish Journal of Environmental Studies* 21(3): 741-747.
- Szatmari, I., T. Barcza, P. S. Kormoczy and P. Laczay (2012). "Ecotoxicological assessment of doxycycline in soil." *Journal of Environmental Science and Health Part B-Pesticides Food Contaminants and Agricultural Wastes* 47(2): 129-135.
- Tapia-Morales, S. and Z. Garcia-Esquivel (2012). "Effect of chloramphenicol and gentamycin on the larval development of the geoduck clam, *Panopea generosa*" *Journal of Shellfish Research* 31(1): 352-353.
- Teixido, E., E. Pique, J. Gomez-Catalan and J. M. Llobet (2013). "Assessment of developmental delay in the zebrafish embryo teratogenicity assay." *Toxicology in Vitro* 27(1): 469-478.
- Tewari, S., R. Jindal, Y. L. Kho, S. Eo and K. Choi (2013). "Major pharmaceutical residues in wastewater treatment plants and receiving waters in Bangkok, Thailand, and associated ecological risks." *Chemosphere* 91(5): 697-704.
- Thienpont, B., A. Tingaud-Sequeira, E. Prats, C. Barata, P. J. Babin and D. Raldua (2011). "Zebrafish eleutheroembryos provide a suitable vertebrate model for screening chemicals that impair thyroid hormone synthesis." *Environmental Science & Technology* 45(17): 7525-7532.
- Thomas, M. A., P. P. Joshi and R. D. Klaper (2012). "Gene-class analysis of expression patterns induced by psychoactive pharmaceutical exposure in fathead minnow (*Pimephales promelas*) indicates induction of neuronal systems." *Comparative Biochemistry and Physiology C-Toxicology & Pharmacology* 155(1): 109-120.
- Thomas, M. A. and R. D. Klaper (2012). "Psychoactive pharmaceuticals induce fish gene expression profiles associated with human idiopathic autism." *PloS one* 7(6).
- Togunde, O. P., K. D. Oakes, M. R. Servos and J. Pawliszyn (2012). "Optimization of solid phase microextraction for non-lethal *in vivo* determination of selected pharmaceuticals in fish muscle using liquid chromatography-mass spectrometry." *Journal of Chromatography A* 1261: 99-106.

- Toumi, H., M. Boumaiza, M. Millet, C. M. Radetski, V. Felten, C. Fouque and J. F. Ferard (2013).
 "Effects of deltamethrin (pyrethroid insecticide) on growth, reproduction, embryonic development and sex differentiation in two strains of *Daphnia magna* (Crustacea, Cladocera)." *Science of the Total Environment* 458: 47-53.
- Tralau, T. and A. Luch (2012). "Drug-mediated toxicity: illuminating the 'bad' in the test tube by means of cellular assays?" *Trends in Pharmacological Sciences* 33(7): 353-364.
- Tripathi, R., S. S. Pancholi and P. Tripathi (2012). "Genotoxicity of ibuprofen in mouse bone marrow cells *in vivo*." *Drug and Chemical Toxicology* 35(4): 389-392.
- Trombini, C., M. Hampel and J. Blasco (2012). "Effects of the exposure of the clam *Ruditapes philippinarum* to carbamazepine, diclofenac and ibuprofen. Evaluation of enzymatic and molecular endpoints." *Comparative Biochemistry and Physiology a-Molecular & Integrative Physiology* 163(1): S22-S22.
- Trovo, A. G., R. F. P. Nogueira, A. Aguera, A. R. Fernandez-Alba and S. Malato (2011). "Degradation of the antibiotic amoxicillin by photo-Fenton process - Chemical and toxicological assessment." *Water Research* 45(3): 1394-1402.
- Tsai, C. W., C. S. Lin and W. H. Wang (2012). "Multi-residue determination of sulfonamide and quinolone residues in fish tissues by high performance liquid chromatography-tandem mass spectrometry (LC-MS/MS)." *Journal of Food and Drug Analysis* 20(3): 674-680.
- Ucan-Marin, F., W. Ernst, R. K. O'Dor and J. Sherry (2012). "Effects of food borne ivermectin on juvenile Atlantic salmon (*Salmo salar* L.): Survival, growth, behavior, and physiology." *Aquaculture* 334: 169-175.
- Uchea, C., S. Sarda, T. Schulz-Utermoehl, S. Owen and K. J. Chipman (2013). "*In vitro* models of xenobiotic metabolism in trout for use in environmental bioaccumulation studies." *Xenobiotica* 43(5): 421-431.
- Underwood, J. C., R. W. Harvey, D. W. Metge, D. A. Repert, L. K. Baumgartner, R. L. Smith, T. M. Roane and L. B. Barber (2011). "Effects of the antimicrobial sulfamethoxazole on groundwater bacterial enrichment." *Environmental Science & Technology* 45(7): 3096-3101.
- Unkiewicz-Winiarczyk, A. and K. Gromysz-Kalkowska (2012). "Effect of temperature on toxicity of deltamethrin and oxygen consumption by *Porcellio scaber* Latr (Isopoda)." *Bulletin of Environmental Contamination and Toxicology* 89(5): 960-965.
- Valcarcel, Y., S. G. Alonso, J. L. Rodriguez-Gil, A. Gil and M. Catala (2011). "Detection of pharmaceutically active compounds in the rivers and tap water of the Madrid Region (Spain) and potential ecotoxicological risk." *Chemosphere* 84(10): 1336-1348.
- van Leeuwen, J. S., B. Unlu, N. P. E. Vermeulen and J. C. Vos (2012). "Differential involvement of mitochondrial dysfunction, cytochrome P450 activity, and active transport in the toxicity of structurally related NSAIDs." *Toxicology in Vitro* 26(2): 197-205.
- Vani, T., N. Saharan, S. Mukherjee, R. Ranjan, R. Kumar and R. K. Brahmchari (2011).
 "Deltamethrin induced alterations of hematological and biochemical parameters in fingerlings of *Catla catla* (Ham.) and their amelioration by dietary supplement of vitamin C." *Pesticide Biochemistry and Physiology* 101(1): 16-20.
- Vazquez-Roig, P., V. Andreu, C. Blasco and Y. Pico (2012). "Risk assessment on the presence of pharmaceuticals in sediments, soils and waters of the Pego-Oliva Marshlands (Valencia, eastern Spain)." *Science of the Total Environment* 440: 24-32.

- Vazquez-Roig, P., C. Blasco and Y. Pico (2013). "Advances in the analysis of legal and illegal drugs in the aquatic environment." *Trac-Trends in Analytical Chemistry* 50: 65-77.
- Veach, A., M. J. Bernot and J. K. Mitchell (2012). "The influence of six pharmaceuticals on freshwater sediment microbial growth incubated at different temperatures and UV exposures." *Biodegradation* 23(4): 497-507.
- Velasco-Santamaria, Y. M., B. Korsgaard, S. S. Madsen and P. Bjerregaard (2011). "Bezafibrate, a lipid-lowering pharmaceutical, as a potential endocrine disruptor in male zebrafish (*Danio rerio*)." *Aquatic Toxicology* 105(1-2): 107-118.
- Vendrell, D., L. Serarols, J. L. Balcazar, I. de Blas, O. Girones, J. L. Muzquiz and I. Ruiz-Zarzuela (2012). "Accumulation and depletion kinetics of erythromycin in rainbow trout (*Oncorhynchus mykiss*)." *Preventive Veterinary Medicine* 105(1-2): 160-163.
- Verlicchi, P., M. Al Aukidy and E. Zambello (2012). "Occurrence of pharmaceutical compounds in urban wastewater: Removal, mass load and environmental risk after a secondary treatment-A review." *Science of the Total Environment* 429: 123-155.
- Vidal-Dorsch, D. E., S. M. Bay, K. Maruya, S. A. Snyder, R. A. Trenholm and B. J. Vanderford (2012). "Contaminants of emerging concern in municipal wastewater effluents and marine receiving water." *Environmental Toxicology and Chemistry* 31(12): 2674-2682.
- Vosges, M., O. Kah, N. Hinfray, E. Chadili, Y. Le Page, Y. Combarnous, J. M. Porcher and F. Brion (2012). "17 alpha-Ethinylestradiol and nonylphenol affect the development of forebrain GnRH neurons through an estrogen receptors-dependent pathway." *Reproductive Toxicology* 33(2): 198-204.
- Vulliet, E., C. Cren-Olive and M. F. Grenier-Loustalot (2011). "Occurrence of pharmaceuticals and hormones in drinking water treated from surface waters." *Environmental Chemistry Letters* 9(1): 103-114.
- Vystavna, Y., F. Huneau, V. Grynenko, Y. Vergeles, H. Celle-Jeanton, N. Tapie, H. Budzinski and P. Le Coustumer (2012). "Pharmaceuticals in rivers of two regions with contrasted socioeconomic conditions: occurrence, accumulation, and comparison for Ukraine and France." *Water Air and Soil Pollution* 223(5): 2111-2124.
- Vystavna, Y., P. Le Coustumer and F. Huneau (2013). "Monitoring of trace metals and pharmaceuticals as anthropogenic and socio-economic indicators of urban and industrial impact on surface waters." *Environmental Monitoring and Assessment* 185(4): 3581-3601.
- Waiser, M. J., D. Humphries, V. Tumber and J. Holm (2011). "Effluent-dominated streams. Part
 2: Presence and possible effects of pharmaceuticals and personal care products in Wascana creek, Saskatchewan, Canada." *Environmental Toxicology and Chemistry* 30(2): 508-519.
- Wang, H.-c., H.-q. Zhao, D. Qian and Q.-q. Pan (2011). "Effect of common antibiotics in aquaculture on phenoloxidase activity in mud crab." *Fisheries Science (Liaoning)* 30(1): 38-41.
- Wang, H. P., J. J. Wang, T. T. Wu, F. Qin, X. Q. Hu, L. H. Wang and Z. Z. Wang (2011).
 "Molecular characterization of estrogen receptor genes in *Gobiocypris rarus* and their expression upon endocrine disrupting chemicals exposure in juveniles." *Aquatic Toxicology* 101(1): 276-287.
- Wang, J. and P. R. Gardinali (2012). "Analysis of selected pharmaceuticals in fish and the fresh water bodies directly affected by reclaimed water using liquid chromatography-tandem mass spectrometry." *Analytical and Bioanalytical Chemistry* 404(9): 2711-2720.

- Wang, L., G. G. Ying, J. L. Zhao, S. Liu, B. Yang, L. J. Zhou, R. Tao and H. C. Su (2011). "Assessing estrogenic activity in surface water and sediment of the Liao River system in northeast China using combined chemical and biological tools." *Environmental Pollution* 159(1): 148-156.
- Wang, L. Y., N. F. Y. Tam and X. H. Zhang (2011). "Assimilation of 17 alpha-ethinylestradiol by sludge and its stress on microbial communities under aerobic and anaerobic conditions." *Journal of Environmental Science and Health Part a-Toxic/Hazardous Substances & Environmental Engineering* 46(3): 242-247.
- Wang, S., K. D. Oakes, L. M. Bragg, J. Pawliszyn, G. Dixon and M. R. Servos (2011). "Validation and use of *in vivo* solid phase micro-extraction (SPME) for the detection of emerging contaminants in fish." *Chemosphere* 85(9): 1472-1480.
- Wang, S. Y. and C. K. Gunsch (2011). "Effects of selected pharmaceutically active compounds on the ammonia oxidizing bacterium *Nitrosomonas europaea*." *Chemosphere* 82(4): 565-572.
- Wang, X.-N., Z.-T. Liu, Z.-G. Yan, C. Zhang, W.-L. Wang, J.-L. Zhou and S.-W. Pei (2013).
 "Development of aquatic life criteria for triclosan and comparison of the sensitivity between native and non-native species." *Journal of Hazardous Materials* 260: 1017-1022.
- Wassnig, M. and P. C. Southgate (2011). "The effects of egg stocking density and antibiotic treatment on survival and development of winged pearl oyster (*Pteria penguin*, Roding 1798) embryos." *Journal of Shellfish Research* 30(1): 103-107.
- Waye, A. and V. L. Trudeau (2011). "Neuroendocrine disruption: More than hormones are upset." *Journal of Toxicology and Environmental Health-Part B-Critical Reviews* 14(5-7): 270-291.
- Weigt, S., N. Huebler, R. Strecker, T. Braunbeck and T. H. Broschard (2011). "Zebrafish (*Danio rerio)* embryos as a model for testing proteratogens." *Toxicology* 281(1-3): 25-36.
- Wen, B., Y. Liu, P. Wang, T. Wu, S. Z. Zhang, X. Q. Shan and J. F. Lu (2012). "Toxic effects of chlortetracycline on maize growth, reactive oxygen species generation and the antioxidant response." *Journal of Environmental Sciences-China* 24(6): 1099-1105.
- Wiklund, A. K. E., H. Oskarsson, G. Thorsen and L. Kumblad (2011). "Behavioural and physiological responses to pharmaceutical exposure in macroalgae and grazers from a Baltic Sea littoral community." *Aquatic Biology* 14(1): 29-39.
- Wilde, M. L., W. M. M. Mahmoud, K. Kummerer and A. F. Martins (2013). "Oxidationcoagulation of beta-blockers by (K2FeO4)-O-VI in hospital wastewater: Assessment of degradation products and biodegradability." *Science of the Total Environment* 452: 137-147.
- Wong, J., S. Bromand, G. Saffi, S. Bloch, L. Lee, P. Hodson and N. Bols (2011). "Cellular vacuolization is induced in fish cells by environmental pharmaceuticals of the amine drug class (PO)." *Canadian Technical Report of Fisheries and Aquatic Sciences* 2949: 12.
- Wu, N., H. Wei, H. Shen, T. T. Wu and M. Guo (2012). "Acute toxic effects of deltamethrin on red swamp crayfish, *Procambrus clarkii* (Decapoda, Cambaridae)." *Crustaceana* 85(8): 993-1005.
- Wu, S. J., L. L. Zhang and J. M. Chen (2012). "Paracetamol in the environment and its degradation by microorganisms." *Applied Microbiology and Biotechnology* 96(4): 875-884.
- Xie, X. J., Q. X. Zhou, Q. H. Bao, Z. C. He and Y. Y. Bao (2011). "Genotoxicity of tetracycline as an emerging pollutant on root meristem cells of wheat (*Triticum aestivum* L.)." *Environmental Toxicology* 26(4): 417-423.

- Xie, X. J., Q. X. Zhou, D. S. Lin, J. M. Guo and Y. Y. Bao (2011). "Toxic effect of tetracycline exposure on growth, antioxidative and genetic indices of wheat (*Triticum aestivum* L.)." *Environmental Science and Pollution Research* 18(4): 566-575.
- Xiong, Z.-H., W.-L. Yu and P. Hu (2010). "Degradation of the mixed pharmaceuticals with low concentration in aqueous by solar/ferrioxalate system and process evaluation through toxicity bioassay." *Huan jing ke xue= Huanjing kexue / [bian ji, Zhongguo ke xue yuan huan jing ke xue wei yuan hui "Huan jing ke xue" bian ji wei yuan hui.*/31(10): 2336-2343.
- Xu, H., Y. Lin, Q. Sun and C. Yu (2012). "Ecological effects of triclosan and its transport and transformation in the environment." *Asian Journal of Ecotoxicology* 7(3): 225-233.
- Yan, C. X., Y. Yang, J. L. Zhou, M. Liu, M. H. Nie, H. Shi and L. J. Gu (2013). "Antibiotics in the surface water of the Yangtze Estuary: Occurrence, distribution and risk assessment." *Environmental Pollution* 175: 22-29.
- Yan, H., D. D. Wang, B. Dong, F. F. Tang, B. C. Wang, H. Fang and Y. L. Yu (2011). "Dissipation of carbendazim and chloramphenicol alone and in combination and their effects on soil fungal: bacterial ratios and soil enzyme activities." *Chemosphere* 84(5): 634-641.
- Yan, Z. H., G. H. Lu, J. C. Liu and S. G. Jin (2012). "An integrated assessment of estrogenic contamination and feminization risk in fish in Taihu Lake, China." *Ecotoxicology and Environmental Safety* 84: 334-340.
- Yan, Z. H., G. H. Lu and X. F. Yang (2013). "Single and combined effects of estrone and 17 betaestradiol on male goldfish." *Biomedical and Environmental Sciences* 26(3): 176-184.
- Yang, J.-L., Y.-F. Li, W.-Y. Bao, C. G. Satuito and H. Kitamura (2011). "Larval metamorphosis of the mussel *Mytilus galloprovincialis* Lamarck, 1819 in response to neurotransmitter blockers and tetraethylammonium." *Biofouling* 27(2): 193-199.
- Yang, W. W., Z. P. Tang, F. Q. Zhou, W. H. Zhang and L. R. Song (2013). "Toxicity studies of tetracycline on *Microcystis aeruginosa* and *Selenastrum capricornutum*." *Environmental Toxicology and Pharmacology* 35(2): 320-324.
- Yergeau, E., S. Sanschagrin, M. J. Waiser, J. R. Lawrence and C. W. Greer (2012). "Sub-inhibitory concentrations of different pharmaceutical products affect the meta-transcriptome of river biofilm communities cultivated in rotating annular reactors." *Environmental Microbiology Reports* 4(3): 350-359.
- Yildiz, H. Y. and S. Altunay (2011). "Physiological stress and innate immune response in gilthead sea bream (*Sparus aurata*) and sea bass (*Dicentrarchus labrax*) exposed to combination of trimethoprim and sulfamethoxazole (TMP-SMX)." *Fish Physiology and Biochemistry* 37(3): 401-409.
- Yonar, M. E. and F. Sakin (2011). "Ameliorative effect of lycopene on antioxidant status in *Cyprinus carpio* during pyrethroid deltamethrin exposure." *Pesticide Biochemistry and Physiology* 99(3): 226-231.
- Yonar, S. M., F. Sakin, M. E. Yonar, U. Ispir and M. Kirici (2011). "Oxidative stress biomarkers of exposure to deltamethrin in rainbow trout fry (*Oncorhynchus mykiss*)." *Fresenius Environmental Bulletin* 20(8): 1931-1935.
- Yu, Y., L. S. Wu and A. C. Chang (2013). "Seasonal variation of endocrine disrupting compounds, pharmaceuticals and personal care products in wastewater treatment plants." *Science of the Total Environment* 442: 310-316.

- Yu, Y. Y., Q. X. Huang, Z. F. Wang, K. Zhang, C. M. Tang, J. L. Cui, J. L. Feng and X. Z. Peng (2011). "Occurrence and behavior of pharmaceuticals, steroid hormones, and endocrine-disrupting personal care products in wastewater and the recipient river water of the Pearl River Delta, South China." *Journal of Environmental Monitoring* 13(4): 871-878.
- Yuan, H., Y. Zhang and X. Zhou (2012). "Degradation of bezafibrate with UV/H2O2 in surface water and wastewater treatment plant effluent." *Clean-Soil Air Water* 40(3): 239-245.
- Zaleska-Radziwill, M., M. Lebkowska, K. Affek and A. Zarzeczna (2011). "Environmental risk assessment of selected pharmaceuticals present in surface waters in relation to animals." *Archives of Environmental Protection* 37(3): 31-42.
- Zellner, D., B. Padnos, D. L. Hunter, R. C. MacPhail and S. Padilla (2011). "Rearing conditions differentially affect the locomotor behavior of larval zebrafish, but not their response to valproate-induced developmental neurotoxicity." *Neurotoxicology and Teratology* 33(6): 674-679.
- Zhang, J., V. W. C. Chang, A. Giannis and J.-Y. Wang (2013). "Removal of cytostatic drugs from aquatic environment: a review." *Science of the Total Environment* 445: 281-298.
- Zhang, R. J., J. H. Tang, J. Li, Z. N. Cheng, C. Chaemfa, D. Y. Liu, Q. Zheng, M. K. Song, C. L. Luo and G. Zhang (2013). "Occurrence and risks of antibiotics in the coastal aquatic environment of the Yellow Sea, North China." *Science of the Total Environment* 450: 197-204.
- Zhang, R. J., J. H. Tang, J. Li, Q. Zheng, D. Liu, Y. J. Chen, Y. D. Zou, X. X. Chen, C. L. Luo and G. Zhang (2013). "Antibiotics in the offshore waters of the Bohai Sea and the Yellow Sea in China: Occurrence, distribution and ecological risks." *Environmental Pollution* 174: 71-77.
- Zhang, R. J., G. Zhang, Q. Zheng, J. H. Tang, Y. J. Chen, W. H. Xu, Y. D. Zou and X. X. Chen (2012). "Occurrence and risks of antibiotics in the Laizhou Bay, China: Impacts of river discharge." *Ecotoxicology and Environmental Safety* 80: 208-215.
- Zhang, W., M. Zhang, K. F. Lin, W. F. Sun, B. Xiong, M. J. Guo, X. H. Cui and R. B. Fu (2012). "Eco-toxicological effect of carbamazepine on *Scenedesmus obliquus* and *Chlorella pyrenoidosa*." *Environmental Toxicology and Pharmacology* 33(2): 344-352.
- Zhang, X., D. D. Zhang, H. Zhang, Z. X. Luo and C. Z. Yan (2012). "Occurrence, distribution, and seasonal variation of estrogenic compounds and antibiotic residues in Jiulongjiang River, South China." *Environmental Science and Pollution Research* 19(5): 1392-1404.
- Zhang, X. Y. and Z. Y. Gong (2013). "Fluorescent transgenic zebrafish Tg(nkx2.2a:mEGFP) provides a highly sensitive monitoring tool for neurotoxins." *PloS one* 8(2).
- Zhang, Y., X. Y. Cai, X. M. Lang, X. L. Qiao, X. H. Li and J. W. Chen (2012). "Insights into aquatic toxicities of the antibiotics oxytetracycline and ciprofloxacin in the presence of metal: Complexation versus mixture." *Environmental Pollution* 166: 48-56.
- Zhang, Y. Y., H. P. Wang, F. Qin, S. Z. Liu, T. T. Wu, M. Li, P. Xu, X. Zhang, X. Q. Wang, G. J. Hu, C. Yuan and Z. Z. Wang (2012). "Molecular characterization of estrogen receptor genes in loach *Paramisgurnus dabryanus* and their expression upon 17 alpha-ethinylestradiol exposure in juveniles." *General and Comparative Endocrinology* 178(2): 194-205.
- Zhao, J.-L., G.-G. Ying, B. Yang, S. Liu, L.-J. Zhou, Z.-F. Chen and H.-J. Lai (2011). "Screening of multiple hormonal activities in surface water and sediment from the Pearl River system, South China, using effect-directed *in vitro* bioassays." *Environmental Toxicology and Chemistry* 30(10): 2208-2215.

- Zheng, Q., R. J. Zhang, Y. H. Wang, X. H. Pan, J. H. Tang and G. Zhang (2012). "Occurrence and distribution of antibiotics in the Beibu Gulf, China: Impacts of river discharge and aquaculture activities." *Marine Environmental Research* 78: 26-33.
- Zhou, L. J., G. G. Ying, S. Liu, J. L. Zhao, B. Yang, Z. F. Chen and H. J. Lai (2013). "Occurrence and fate of eleven classes of antibiotics in two typical wastewater treatment plants in South China." *Science of the Total Environment* 452: 365-376.
- Ziylan, A. and N. H. Ince (2011). "The occurrence and fate of anti-inflammatory and analgesic pharmaceuticals in sewage and fresh water: Treatability by conventional and non-conventional processes." *Journal of Hazardous Materials* 187(1-3): 24-36.
- Zounkova, R., Z. Kliemesova, L. Nepejchalova, K. Hilscherova and L. Blaha (2011). "Complex evaluation of ecotoxicity and genotoxicity of antimicrobials oxytetracycline and flumequine used in aquaculture." *Environmental Toxicology and Chemistry* 30(5): 1184-1189.
- Zuloaga, O., P. Navarro, E. Bizkarguenaga, A. Iparraguirre, A. Vallejo, M. Olivares and A. Prieto (2012). "Overview of extraction, clean-up and detection techniques for the determination of organic pollutants in sewage sludge: a review." *Analytica Chimica Acta* 736: 7-29.